



## An assessment of contaminant concentrations in toothed whale species of the NW Iberian Peninsula: Part I. Persistent organic pollutants



Paula Méndez-Fernandez<sup>a,b,c,\*</sup>, Lynda Webster<sup>d</sup>, Tiphaine Chauvelon<sup>a</sup>, Paco Bustamante<sup>a</sup>, Marisa Ferreira<sup>c</sup>, Angel F. González<sup>e</sup>, Alfredo López<sup>b</sup>, Colin F. Moffat<sup>d</sup>, Graham J. Pierce<sup>f,g</sup>, Fiona L. Read<sup>e</sup>, Marie Russell<sup>d</sup>, Maria B. Santos<sup>h</sup>, Jérôme Spitz<sup>i</sup>, José V. Vingada<sup>c,g</sup>, Florence Caurant<sup>a</sup>

<sup>a</sup> Littoral Environment et Sociétés (LIENSs), UMRI 7266 CNRS-Université de La Rochelle, 2 Rue Olympe de Gouges, 17042 La Rochelle Cedex 01, France

<sup>b</sup> Coordinadora para o Estudo dos Mamíferos Mariños (CEMMA), Apdo. 15, Pontevedra 36380, Spain

<sup>c</sup> Centro de Biología Molecular e Ambiental (CBMA)/Sociedade Portuguesa de Vida Selvagem (SPVS), Dep. de Biologia, Universidade do Minho, Campus de Gualtar, Braga 4710-057, Portugal

<sup>d</sup> Marine Scotland, Marine Laboratory, Victoria Road, Aberdeen AB11 9DB, United Kingdom

<sup>e</sup> Instituto de Investigaciones Marinas (C.S.I.C), Eduardo Cabello 6, 36208 Vigo, Spain

<sup>f</sup> Oceanlab, University of Aberdeen Main Street, Newburgh, Aberdeenshire AB41 6AA, United Kingdom

<sup>g</sup> Centre for Environmental and Marine Studies (CESAM), Universidade de Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

<sup>h</sup> Instituto Español de Oceanografía, Centro Oceanográfico de Vigo, P.O. Box 1552, Vigo 36200, Spain

<sup>i</sup> Marine Mammal Research Unit, Fisheries Centre, University of British Columbia, 2202 Main Mall, Vancouver, British Columbia V6T 1Z4, Canada

### HIGHLIGHTS

- POPs were measured in toothed whales from the NW Iberian Peninsula.
- Bottlenose dolphin and harbour porpoise showed the greatest PCB concentrations.
- The POP levels were higher than in the South Atlantic and Pacific Oceans.

### ARTICLE INFO

#### Article history:

Received 26 July 2013

Received in revised form 11 February 2014

Accepted 13 February 2014

#### Keywords:

PCBs

PBDEs

Bioaccumulation

Patterns

Toothed whales

Northwest Iberian Peninsula

### ABSTRACT

Concentrations and patterns of polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in the blubber of the five most common toothed whales off the Northwest Iberian Peninsula (NWIP), specifically common dolphin, long-finned pilot whale, harbour porpoise, striped dolphin and bottlenose dolphin, were investigated. The study revealed that differences in PCB and PBDE concentrations among the species are highly dependent on age and sex but also on ecological factors such as trophic level, prey type and habitat. Of the five species studied, bottlenose dolphin and harbour porpoise showed the greatest concentrations of PCBs. Both species exceed the toxic threshold of  $17 \mu\text{g g}^{-1}$  lipid weight (PCB Aroclor equivalent) for health effects on marine mammals, for 100% and 75% of the individuals tested, respectively. Overall, the PCB and PBDE levels observed in the NWIP toothed whales were of the same order of magnitude or lower than those reported by previous studies in areas of the NE Atlantic. However, they are often higher than those for toothed whales from the southern Atlantic and Pacific Ocean.

© 2014 Elsevier B.V. All rights reserved.

### 1. Introduction

Human activities in marine and coastal environments have intensified since the 1950s. Furthermore, reliance of human populations on coastal areas for urban development and exploitation of marine

resources is predicted to keep increasing in the near future. The Northwest Iberian Peninsula (NWIP), situated at the northern limit of the NW African upwelling system (Figueiras et al., 2002), is a good example of such processes. Over the last fifty to sixty years, industrial development and an increase in other human activities in the area have increased the pressures on the marine environment. In this context, and to realise the ambition of clean, productive and biologically diverse seas, the European Community developed the Marine Strategy Framework Directive (MSFD, Directive, 2008/56/EC of the European Parliament and of the Council of 17 June 2008) the main objective of which is to

\* Corresponding author at: Littoral Environment et Sociétés (LIENSs), UMRI 7266 CNRS-Université de La Rochelle, 2 Rue Olympe de Gouges, 17042 La Rochelle Cedex 01, France. Tel.: +33 5 46458388; fax: +33 5 46496711.

E-mail address: [paula.mendez\\_fernandez@univ-lr.fr](mailto:paula.mendez_fernandez@univ-lr.fr) (P. Méndez-Fernandez).

deliver “Good Environmental Status” of European marine ecosystems by 2020. To achieve this, better knowledge of the contamination status of marine populations is needed, specifically in connection with both Descriptor 8 and Descriptor 9 of the MSFD.

The persistent organic pollutants (POPs), including polychlorinated biphenyls (PCBs) and pesticides (e.g. dichlorodiphenyltrichloroethane, DDT) are among the primary pollutants of concern in marine ecosystems cited on the OSPAR list of Chemicals for Priority Action (OSPAR, 2010); they are lipophilic synthetic organic compounds that have been produced for industrial and agricultural purposes since the 1940s, or are by-products of other industrial processes developed over a similar period of time. Although their production has been banned since the end of the 1970s, PCBs can still be found in wildlife and other environmental components e.g. sediments (OSPAR, 2010). Other classes of organic chemicals are also of concern nowadays, notably the brominated diphenyl ether formulations (PBDEs) (de Boer et al., 1998) and the hexabromocyclododecanes (HBCDs) another brominated flame retardant (e.g. Zegers et al., 2005). Marine mammals, as long-lived apex predators, are at risk from these toxic compounds, since they have a high bioaccumulation potential and biomagnify through food webs (Aguilar et al., 1999). Due to their lipophilic nature, POPs reach their highest concentrations in fatty tissues and, particularly, in the hypodermic fat or blubber. Compared to most terrestrial mammals, marine mammals appear to have a lower capacity to metabolize and excrete lipophilic organochlorine compounds (Boon et al., 1992; Duinker et al., 1989; Tanabe et al., 1988). This capacity is lower in toothed whales than in pinnipeds (seals and sea lions) (Tanabe et al., 1988), which makes them especially vulnerable to POPs. Although information on the actual effect of POPs on the health of marine mammals is scarce (Reijnders et al., 1999), results from laboratory feeding studies and field investigations have allowed the determination of several threshold values for adverse effects (e.g. Kannan et al., 2000). The concentration of contaminants in marine mammal tissues primarily varies in relation to prey consumption, but there is also a function of their specific capacity to transform these compounds to metabolised forms and/or ultimately excrete the native form or the associated metabolites (Aguilar et al., 1999). Other biological factors have also been found to be responsible for variation in POP concentrations in marine mammals. These include body size and composition, nutritive condition, age, sex, health status, duration of lactation, transfer from mother to offspring during both pregnancy and lactation (Aguilar et al., 1999). Thus, since the uptake of contaminants in marine mammals depends on the diet, feeding habitat and biological factors, any interpretation of concentrations or comparison between species would be incomplete without considering as many of the factors as possible.

For many years, the concentration of contaminants in the NWIP has been routinely monitored through the analysis of samples of sediments, seawater and commercial species such as shellfish (e.g. Carro et al., 2002; Prego and Cobelo-García, 2003). Potentially toxic substances have also occasionally been investigated in marine mammals since the 1980s, as part of the European funded BIOCET project (Murphy et al., 2010; Pierce et al., 2008; Zegers et al., 2005) among others (Borrell et al., 2001, 2006; Tornero et al., 2006), although to a lesser extent than in other marine organisms from this area.

The overall objective of this study is to assess the contamination status of the five most common marine mammals in the NWIP: the common dolphin (*Delphinus delphis*), the long-finned pilot whale (*Globicephala melas*), the harbour porpoise (*Phocoena phocoena*), the striped dolphin (*Stenella coeruleoalba*) and the bottlenose dolphin (*Tursiops truncatus*). This paper constitutes the first of a two part study. In this first part we report on the PCB and PBDE concentrations and patterns in these species, and evaluate their contamination status in comparison with threshold values for health effects on marine mammals as well as making comparisons with concentrations found in other geographical areas. In Part II of this study, that will be subsequently reported (Méndez-Fernández et al., in press), we investigate the

concentrations of trace elements, which is another group of potential contaminants in the NWIP, in the context of biological and ecological factors.

## 2. Materials and methods

### 2.1. Sampling and study area


Sampling was carried out in the NWIP, from the northern limit of the Galician coast in Spain (43°3'N, 7°2'W) to Nazaré on the Portuguese coast (39°36'N, 9°3'W) (Fig. 1). Experienced members of the Spanish (*Coordinadora para o Estudo dos Mamíferos Mariños*, CEMMA) and Portuguese (*Sociedade Portuguesa de Vida Salvagem*, SPVS) stranding networks have been attending stranded and by-caught cetaceans for over twenty years and over ten years, respectively. Animals were identified to species, measured, sexed and, if the decomposition state of the carcass allowed, full necropsies were performed and samples collected whenever possible. All procedures followed the standard protocol defined by the European Cetacean Society (ECS), as did the coding of decomposition state and condition (Kuiken and García Hartmann, 1991).

A total of 172 stranded and by-caught individuals was selected for this study, covering five toothed whale species (common dolphin,  $n = 114$ ; long-finned pilot whale,  $n = 9$ ; harbour porpoise,  $n = 19$ ; striped dolphin,  $n = 21$  and bottlenose dolphin,  $n = 9$ ) over the period 2004 to 2008. The common dolphin is the cetacean species stranded in the greatest numbers; this is believed to reflect the relatively high abundance in the area (Santos et al., 2013c) and the large number of individuals being by-caught in NWIP fisheries (López et al., 2002, 2003). The animals recovered in a “fresh” state (a score of 1 to 3 from the ECS protocol, i.e. originally stranded alive, freshly dead or mildly decomposed) were selected. Teeth were extracted for age determination, gonads collected for determination of reproductive status and blubber samples for POP analyses. All blubber samples were taken from the left side in front of the dorsal fin. Samples were entire vertical cross-sections of the blubber so as to prevent any possible effects of stratification of the blubber. The samples were wrapped separately in aluminium foil and after the necropsies, all samples were stored frozen at  $-20\text{ }^{\circ}\text{C}$  until required for analysis.

Analysis of POPs is costly and the present study was budget-limited. As such, effort was focused on the best sample sets (i.e. individuals for which most data on other variables were available). Thus, in this part of the study 120 blubber samples, out of a possible 172, were analysed for PCBs (common dolphin,  $n = 81$ ; long-finned pilot whale,  $n = 3$ ; harbour porpoise,  $n = 12$ ; striped dolphin,  $n = 15$  and bottlenose dolphin,  $n = 7$ ) and 20 for PBDEs (common dolphin,  $n = 19$ ; harbour porpoise,  $n = 1$ ).

### 2.2. Age estimation and reproductive status

Age was estimated by analysing growth layer groups (GLGs) in the dentine and cementum of teeth, following adapted methods based on Lockyer (1993) and Hohn and Lockyer (1995). Teeth were decalcified and sectioned at  $25\text{ }\mu\text{m}$  using a cryostat. The most central and complete sections (including the whole pulp cavity) were selected from each tooth, stained with Mayer's haematoxylin (modified by Grue) and ‘blued’ in a weak ammonia solution, mounted on glass slides and allowed to dry. GLGs were counted under a binocular microscope. All readings were made blind (without access to individual biological data), and replicate counts were made by two readers. If the age estimates obtained by the two readers differed by more than 1 year, readings were repeated. If the increments were difficult to count, both readers discussed the interpretation and either reached an agreed age or judged the tooth to be unreadable.



**Fig. 1.** Map of the study area including the 100 and 200 m isobaths. The 200 m isobath was taken as the limit for the shelf-break. The black line contouring the coast represents the length of the coastline which constituted the sampling area.

### 2.2.1. Female reproductive status

Females were examined at the time of necropsy for evidence of pregnancy and/or lactation. Formalin-fixed ovaries were weighed, measured and sectioned at 2 mm intervals along the broad ligament. The presence of mature follicles, *corpora lutea* and *corpora albicantia* were recorded. Microscopic examination was conducted to confirm macroscopic findings. Portions of ovary were paraffin-embedded and sectioned at 5–8  $\mu\text{m}$ , and sections were then stained with Mayer's haematoxylin and eosin and examined by microscopy. Females were classified as adults (mature) and juveniles (immature) based on the presence/absence of ovarian structures.

### 2.2.2. Male reproductive status

Testes with attached epididymis were weighed, and a central cross-section was formalin-fixed. Standard histological analysis of paraffin-embedded sections was conducted. Sections were cut at 5–8  $\mu\text{m}$  and stained with Mayer's haematoxylin and eosin. Microscopic analysis was conducted to measure the diameter of seminiferous tubules and to record cell activity (Sertoli cells, interstitial tissue, and germinal cells such as spermatogonia, spermatocytes, spermatids and spermatozoa). Males were classified as adults (mature) and juveniles (immature) based on seminiferous tubule diameter and cell activity.

## 2.3. Determination of persistent organic pollutants

### 2.3.1. Lipid determination

The total lipid content was determined using a modified Folch et al. (1957) method. The samples were weighed (100–150 mg), homogenised, and then extracted three times with a mixture of chloroform:methanol (1:2, 2:1 and 4:1, v/v). A volume of 6.5 mL of 1% sodium chloride was added and the mixture separated into two phases. The lower layer, containing the lipid and lipophilic compounds, was collected and traces of water removed by addition of dry sodium sulphate. These extracts were shaken and stored at 4 °C for 1 h. Centrifugation at 3000 g for 10 min was used to separate the organic extract

from the particulate material and the solvent was removed under a stream of nitrogen in a water bath at 40 °C. When all solvent had evaporated, the weight of residue was determined and the lipid content calculated by gravimetry.

### 2.3.2. Extraction and clean-up for PCB and PBDE analyses

Samples were extracted by Pressurised Liquid Extraction (PLE) (Walsham et al., 2006). For each extraction, approximately 200 mg of blubber was cut (in vertical sections), homogenised, and mixed with sodium sulphate (~20 g). This mixture was spiked with appropriate internal standards (PCBs by GC-MS:  $^{13}\text{C}$ -CB28,  $^{13}\text{C}$ -CB52,  $^{13}\text{C}$ -CB101,  $^{13}\text{C}$ -CB153,  $^{13}\text{C}$ -CB138,  $^{13}\text{C}$ -CB156,  $^{13}\text{C}$ -CB180,  $^{13}\text{C}$ -CB189,  $^{13}\text{C}$ -CB194 and  $^{13}\text{C}$ -CB209; PBDEs: FBDE160<sup>1</sup>). Samples were then refrigerated overnight before being ground to a fine powder using a mortar and pestle. Solvent-washed PLE cells (100 mL) were packed as follows: solvent-washed filter paper, pre-washed sodium sulphate (10 g), 5% deactivated alumina (30 g), solvent-washed filter paper and the samples/sodium sulphate mixture prepared as above.

Samples were extracted by PLE using an ASE 300 (Dionex Ltd., Camberley, Surrey, UK) under elevated temperature (100 °C) and pressure (1500 psi). Five minutes of heating was followed by 2  $\times$  5 min static cycles. The cell flush was 50% total cell volume (i.e. 25% of the cell volume for each flush = 25 mL per flush) with a 120 s purge (using nitrogen) at the end of each sample extraction. The extraction solvent was iso-hexane.

Special precautions were required when analysing PBDEs due to their sensitivity to UV light. Specifically, incoming light was minimized in the laboratory by placing UV filters over the windows.

Following PLE, the extract for PCB analyses was concentrated by Syncore Analyst R-12 (fitted with flushback module) (Buchi UK Ltd, Oldham, UK) to ~0.5 mL and passed through silica columns, before transferring with washing to amber glass GC vials. For 20 samples the extract was split in two, before being concentrated, one half for PBDE analysis and the other one for PCB analysis. The concentrated extracts were analysed for PCBs by Gas Chromatography Electron Impact Mass

Spectrometry (GC–EIMS) and for PBDEs by Gas Chromatography Electron Capture Negative Ionization Mass Spectrometry (GC–ECNIMS).

### 2.3.3. Determination of PCBs by GC–EIMS

The concentrations of 32 PCB congeners (IUPAC PCB numbers 28, 31, 52, 49, 44, 74, 70, 101, 99, 97, 110, 123, 118, 105, 114, 149, 153, 132, 137, 138, 158, 128, 156, 167, 157, 187, 183, 180, 170, 189, 194, 209) were determined by GC–EIMS using a HP6890 series gas chromatograph interfaced with an HP5975 Mass Selective Detector, fitted with a cool on-column injector and a 50 m × 0.22 mm × 25 μm SGE HT-8 column (SGE, Milton Keynes, UK). The initial oven temperature was 80 °C, which was held for 1 min. The temperature was raised by 20 °C min<sup>-1</sup> up to 170 °C and held at this temperature for 7.5 min. This was followed by a ramp of 3 °C min<sup>-1</sup> up to a final temperature of 290 °C which was maintained for 10 min. The MSD was set for selective ion monitoring (SIM) with a dwell time of 50 ms. Calibration standards containing all 32 PCB congeners and covering the concentration range 0.6–500 ng mL<sup>-1</sup> were analysed in triplicate, and the average response used to compute the calibration curve. Correlation coefficients of at least 0.99 were achieved for all PCBs.

### 2.3.4. Determination of PBDE by GC–ECNIMS

PBDEs were analysed and the concentrations of nine congeners, specifically BDE 28, 47, 66, 85, 99, 100, 153, 154 and 183, were determined by GC–ECNIMS using an HP6890 Series gas chromatograph interfaced with an HP5973N MSD, fitted with a cool on-column injector.

A Thames Restek STX-500 column (STX-500, 30 m × 0.25 mm i.d., 0.15 μm film thickness, Thames Restek, Buckinghamshire, UK) was utilised, fitted with a Thames Restek Siltek (0.53 mm i.d.) 5 m guard column. The injector temperature was initially 120 °C and after 2 min the temperature was elevated by 100 °C min<sup>-1</sup> up to 300 °C at which it was maintained until the end of the run. The carrier gas was helium, set at a constant pressure of 15 psi. Methane was used as the reagent gas at a pressure of 1.6 bar. The transfer line was held at 280 °C and the ion source at 150 °C. Injections were made at 120 °C and the oven temperature held constant for 2 min. Thereafter, the temperature was raised by 15 °C min<sup>-1</sup> up to 205 °C. This was followed by a ramp of 6 °C min<sup>-1</sup> up to a final temperature of 330 °C. The MSD was set for selective ion monitoring with a dwell time of 50 ms. The ions monitored were m/z 78.9 and 80.9 (ions equating to bromine) for all PBDEs.

### 2.3.5. Quality control

The methods employed were validated by the replicate analysis of standards and samples, and through spiking experiments or analysis of certified reference materials (CRMs). The limits of detection (LODs) were determined through the repeated analysis of a low spiked sample and calculated from  $4.65 \times SD$  (standard deviation) of the mean concentration (Cheeseman and Wilson, 1989). LODs were dependent on the sample size. The replicate analysis of standards on separate days gave coefficient of variation (CV%) of ~3% for PCBs analysed by GC–EIMS. Recoveries greater than 75% were achieved for PCBs and PBDEs spiked samples and CRMs. Internal quality control procedures incorporated the use of a laboratory reference material (cod liver oil; LRM) for all determinants, and also a CRM for PCBs, in each batch of samples. Procedural blanks were performed with each batch of samples, and the final concentration adjusted accordingly. The data obtained from the LRM were transferred onto NWA Quality Analyst. Thus Shewhart charts were produced with warning and action limits (i.e.  $\pm 2 \times$  and  $\pm 3 \times$  the standard deviation of the mean, respectively). CRM data were accepted if recoveries were between 70% and 120% of the certified concentration. Quality assurance was further demonstrated through successful participation in the Quality Assurance of Information for Marine Environmental Monitoring in Europe (QUASIMEME) Laboratory Performance Studies. Finally, all POP concentrations were normalized to the lipid content (%) of the blubber.

## 2.4. Data treatment

All data submitted to statistical tests were first checked for normality (Shapiro–Wilk test) and for homogeneity of variances (Bartlett test). Non-parametric tests were applied since the distributions of response variables were found to be non-normal and/or the homogeneity criterion was not satisfied.

Prior to the treatment of the POP concentration data (μg g<sup>-1</sup> lipid weight), an age–gender classification of the individuals based upon their sexual maturity was carried out for each species. The individuals were then divided into four groups: adult male, adult female, juvenile male and juvenile female. Hence, differences in the sum of the 32 PCB congeners (ΣPCBs) were tested between species and age–gender groups using the Kruskal–Wallis test followed by pairwise comparison tests, with the exception of pilot whales and age–gender groups with less than 2 individuals, which were excluded from the statistical treatment. For the same reason, no statistical test was performed for the sum of the 9 PBDE (ΣPBDEs) congeners.

We also verified the number of animals exceeding the toxic threshold concentration (17 μg g<sup>-1</sup> lipid wt.) for total PCBs determined for adverse health effects in marine mammals (Kannan et al., 2000). Since this value was based on comparison with the main peaks in the commercial PCB mixture Aroclor 1254, the PCB concentrations in the samples cannot be compared directly to this limit and had to be converted. Aroclor equivalent concentrations were estimated from the concentration of the seven ICES PCBs (i.e. CB28, 52, 101, 118, 138, 153 and 180 as recommended by the EU Community Bureau of Reference), by multiplying the sum of the concentrations for the seven ICES PCBs by 3 (i.e. total PCB concentration [as Aroclor 1254] =  $3.0 \times$  sum of seven ICES congeners in lipid weight) (Jepson et al., 2005).

The levels of significance for statistical analyses were always set at  $\alpha = 0.05$  and analysis was performed using R version 3.0.1 (R Development Core Team, 2010).

## 3. Results and discussion

### 3.1. Persistent organic pollutant concentrations and patterns

Persistent organic pollutants enter marine mammal tissues almost exclusively via their food and the amounts in tissues vary greatly with intake factors (Aguilar, 1989), i.e. trophic level, prey type, and with the local environmental pollution. Thus, all these factors must be taken into consideration when interpreting POP concentrations in marine mammals. The five toothed whale species studied here have rather similar trophic levels in the NWIP, ranging from 4.3 to 5.3 (Méndez-Fernández et al., 2012). However, stomach contents and stable isotope analyses revealed that these species feed on different prey types and forage in different habitats (Méndez-Fernández et al., 2012, 2013; Santos et al., 2007, 2013a, 2013b, 2013c). In addition, marine mammal organochlorine loads tend to increase with age during the juvenile stage of both genders, because the uptake of pollutants usually exceeds metabolism and excretion. In adult males, this pattern continues throughout their life, while in adult females, the transfer of pollutants to offspring during gestation and lactation progressively reduces pollutant concentrations with age (e.g. Borrell et al., 1996; Covaci et al., 2002; Wolkers et al., 2004). In this study, PCBs were detected in all five species across age–gender groups and significant differences were found for ΣPCB mean concentrations (μg g<sup>-1</sup> lipid wt.) among them (Kruskal–Wallis,  $H = 16.13$ ,  $P < 0.01$ ), with bottlenose dolphin showing the highest ΣPCB concentrations followed by harbour porpoise. Pilot whales, common dolphins and striped dolphins exhibited very similar mean age/gender-specific PCB concentrations (Table 1), and as expected the adult males (not represented in bottlenose dolphins and pilot whales) exhibited the highest ΣPCB concentrations (Table 1). However, there was no significant between-species variation in their concentrations (Kruskal–Wallis,  $H = 0.45$ ,  $P = 0.5$ ). In contrast, for adult females

**Table 1**


Age-/gender-specific PCB concentrations (mean  $\pm$  SD,  $\mu\text{g g}^{-1}$  lipid weight), lipid content (%) and age values expressed in years (mean  $\pm$  SD) in blubber of common dolphin (*Delphinus delphis*), pilot whale (*Globicephala melas*), harbour porpoise (*Phocoena phocoena*), striped dolphin (*Stenella coeruleoalba*) and bottlenose dolphin (*Tursiops truncatus*) from the North West Iberian Peninsula. AF: adult female, AM: adult male, JF: juvenile female and JM: juvenile male, n = sample size.

Species and age-gender group	n	Age	Lipid content	CB153	$\Sigma\text{PCB}^a$
<i>Delphinus delphis</i>	81	6.6 $\pm$ 5.4	60.6 $\pm$ 16.2	5.2 $\pm$ 5	17.2 $\pm$ 14.1
AF	11	14.4 $\pm$ 3.1	56.9 $\pm$ 17.5	2.5 $\pm$ 2.5	8.7 $\pm$ 8.1
AM	8	14.3 $\pm$ 3.8	40.5 $\pm$ 11.9	13.1 $\pm$ 8.2	38.9 $\pm$ 22.2
JF	20	3.4 $\pm$ 3.3	66.3 $\pm$ 13.8	3.7 $\pm$ 2.0	13.2 $\pm$ 6.3
JM	42	4.4 $\pm$ 3.1	62.5 $\pm$ 14.1	5 $\pm$ 4.3	16.9 $\pm$ 12.1
<i>Globicephala melas</i>	3	5.2 $\pm$ 5.5	65.2 $\pm$ 10.9	3.6 $\pm$ 4.1	16.2 $\pm$ 19.7
AF	1	11.5	77.3	0.4	2
JF	1	2.0	62.3	2.2	7.9
JM	1	2.0	56	8.3	38.7
<i>Phocoena phocoena</i>	12	7.0 $\pm$ 6.5	77.9 $\pm$ 13.9	6.3 $\pm$ 6.7	20.5 $\pm$ 20.4
AF	3	13.7 $\pm$ 6.2	76.8 $\pm$ 8.1	12.0 $\pm$ 9.7	37.5 $\pm$ 30.8
AM	1	18.0	53.7	16.6 $\pm$ 0	50.8
JF	5	2.8 $\pm$ 1.6	87.0 $\pm$ 13.7	2.9 $\pm$ 0.8	10.8 $\pm$ 2.8
JM	3	3.7 $\pm$ 1.2	71.7 $\pm$ 7.9	2.8 $\pm$ 1	9.4 $\pm$ 3
<i>Stenella coeruleoalba</i>	15	4.9 $\pm$ 5.5	61.1 $\pm$ 19.1	4.2 $\pm$ 5.1	15.7 $\pm$ 18.6
AF	2	14.0	65.6 $\pm$ 8.7	0.3 $\pm$ 0.01	1.8 $\pm$ 0.3
AM	1	15.0	50.1	10.7	36.4
JF	5	3.8 $\pm$ 3.5	65.7 $\pm$ 5.5	2.6 $\pm$ 1.6	9.9 $\pm$ 5.4
JM	7	1.6 $\pm$ 2.0	58.2 $\pm$ 27.6	5.5 $\pm$ 6.6	20.9 $\pm$ 24.4
<i>Tursiops truncatus</i>	7	3.8 $\pm$ 2.0	68.6 $\pm$ 9.3	15.2 $\pm$ 10.1	56.4 $\pm$ 35.2
JF	3	2.3 $\pm$ 1.6	71.8 $\pm$ 0.8	13.3 $\pm$ 8.8	48.9 $\pm$ 30.9
JM	4	4.9 $\pm$ 1.7	66.3 $\pm$ 12.4	16.7 $\pm$ 12.1	62.1 $\pm$ 41.8

<sup>a</sup>  $\Sigma\text{PCB}$  includes 32 congeners. See Materials and methods section for full list.

(not represented in bottlenose dolphins) there were significantly higher concentrations in harbour porpoise than in common and striped dolphin (post-hoc test,  $P < 0.05$ ; Table 1). Significant differences were found between juvenile bottlenose dolphins and juveniles of all the other species (post-hoc, all  $P < 0.05$ ). The high concentrations found in juvenile bottlenose dolphins and adult harbour porpoises are in accordance with their coastal habitat, their proximity to areas with the highest anthropogenic impact from contaminants, and their mainly fish-feeding dietary habits in the NWIP (López et al., 2004; Méndez-Fernández et al., 2012; Pierce et al., 2010). In addition, both species are more frequently seen in the southern part of the study area (López et al., 2004; Pierce et al., 2010), which is more populated and industrialized than the northern part. Thus, these ecological factors may be one of the reasons for the high  $\Sigma\text{PCB}$  concentrations found for these two species in the NWIP waters. Although pilot whales and striped dolphins are observed in neritic habitats in the NWIP, especially for feeding (Méndez-Fernández et al., 2013), these cetacean species are mainly associated with oceanic habitats, remote from land-based sources of contamination (López et al., 2004; Méndez-Fernández et al., 2012; Pierce et al., 2010). In addition, pilot whales mainly feed on cephalopods that generally, and depending on species, contain lower concentrations of PCBs than fish from a similar geographic region (e.g. Storelli, 2008).

Their different ecological feeding patterns are also reflected in the relative contribution of the distinct PCB congeners (Fig. 2). The mainly cephalopod feeders and oceanic species, the pilot whale and the striped dolphin, had a higher proportion of less chlorinated congeners (i.e. tri-, tetra- and penta-chlorobiphenyls) than the other three species, which had a greater proportion of highly chlorinated PCB congeners such as hexa- and hepta-chlorobiphenyls. This finding can be explained by the more efficient long-range transport of low-chlorinated PCBs through both atmosphere and water (Beyer et al., 2000). Nevertheless, all species showed a predominance of PCBs containing 5 or more chlorines. Hexachlorobiphenyls (56.3%) accounting for the highest percentage across all 5 cetaceans, followed by heptachlorobiphenyls (26%) and pentachlorobiphenyls (12.7%) (Fig. 2). In addition, among the pentachlorinated congeners CB153 was the predominant followed by



**Fig. 2.** Relative contribution of PCB congeners, grouped by the number of chlorine atoms in the molecule, to the  $\Sigma\text{PCB}$  in blubber of Dde: common dolphin (*Delphinus delphis*), Gme: pilot whale (*Globicephala melas*), Pph: harbour porpoise (*Phocoena phocoena*), Sco: striped dolphin (*Stenella coeruleoalba*) and Ttr: bottlenose dolphin (*Tursiops truncatus*) from the North West Iberian Peninsula.

CB138, 187 and 180. These results are in accordance with the patterns observed in cetacean species from different regions (e.g. Covaci et al., 2002; Lailson-Brito et al., 2012; Leonel et al., 2012; Wafo et al., 2005).

Intra-specific differences were only found for common dolphin (Kruskal–Wallis,  $H = 16.69$ ,  $P < 0.05$ ), which was the best represented species ( $n = 81$ ; Table 1). Specifically, the adult males showed significantly higher concentrations than females ( $38.9 \pm 22.2$  and  $8.7 \pm 8.1 \mu\text{g g}^{-1}$  lipid wt., respectively; Table 1). This suggests that adult females have a different accumulation pattern, which is consistent with the well-described transfer of POPs from mother to offspring during gestation and lactation discussed above.

The PBDE concentrations were almost 10 times lower than PCBs in blubber for common dolphin and harbour porpoise and for all age-gender groups (Table 2). This agrees with several previous studies on toothed whales (Dorneles et al., 2010; Leonel et al., 2012; Nyman et al., 2002; Pierce et al., 2008; Yogui et al., 2011). Of the 9 congeners analysed, congener BDE85 was not detected in any sample and congeners BDE28 and BDE183 were detected only in juvenile female and in adult male, adult female and juvenile male of common dolphin, respectively (Table 2). On average, BDE47 showed the highest concentrations in both species and in all age-gender groups, following the same pattern of variation as  $\Sigma\text{PBDE}$  concentrations (Table 2). Similar profiles were reported in other cetacean species from around the world (e.g. Dorneles et al., 2010; Leonel et al., 2012; Weijs et al., 2009).

The concentrations of  $\Sigma\text{PBDE}$  of the juvenile males of both species were similar, being slightly higher for harbour porpoise with  $0.57 \mu\text{g g}^{-1}$  lipid wt. ( $n = 1$ ) than for common dolphin with  $0.31 \pm 0.18 \mu\text{g g}^{-1}$  lipid wt. ( $n = 11$ ). For common dolphin, adult males were more contaminated than adult females with the adult male group having the highest ( $0.71 \pm 0.17 \mu\text{g g}^{-1}$  lipid wt.) and the adult female the lowest ( $0.08 \mu\text{g g}^{-1}$  lipid wt.) mean concentrations. Similar to PCBs, this result supports the hypothesis that adult female animals reduce their PBDE concentrations through gestation and lactation. Besides differences of  $\Sigma\text{PBDE}$  concentrations found between juvenile males of both species, their patterns were also slightly different, namely BDE47 > BDE100 > BDE154 > BDE99 > BDE153 > BDE66 > BDE183 for common dolphin and BDE47 > BDE100 > BDE99 > BDE154 > BDE153 > BDE66 (BDE183 was not detected) for harbour porpoise. The profiles found in the different age-gender groups of common dolphin were similar. The pattern reported in harbour porpoise is similar with those from other regions and from other cetacean species (e.g. Boon et al., 2002; Leonel et al., 2012; Weijs et al., 2009). Thus, in

**Table 2**

Age-/gender-specific PBDE concentrations (mean  $\pm$  SD,  $\mu\text{g g}^{-1}$  lipid weight), lipid content (%) and age values expressed in years (mean  $\pm$  SD) in blubber of common dolphin (*Delphinus delphis*) and harbour porpoise (*Phocoena phocoena*) from the North West Iberian Peninsula. AF: adult female, AM: adult male, JF: juvenile female and JM: juvenile male, n = sample size.

Species/ age-gender group	n	Age	Lipid content	BDE28	BDE47	BDE66	BDE100	BDE99	BDE154	BDE153	BDE183	$\Sigma\text{BDE}^a$
<i>Delphinus delphis</i>	19	7.4 $\pm$ 5.1	59.9 $\pm$ 13.8	0.01	0.23 $\pm$ 0.19	0.004 $\pm$ 0.003	0.05 $\pm$ 0.04	0.02 $\pm$ 0.01	0.03 $\pm$ 0.01	0.012	0.001 $\pm$ 0.0002	0.34 $\pm$ 0.25
AF	1	16.0	63.4	nd	0.03	0.004	0.01	0.01	0.01	0.008	0.001	0.08
AM	2	15.5 $\pm$ 0.7	42.3	nd	0.44 $\pm$ 0.12	0.011	0.12 $\pm$ 0.45	0.04 $\pm$ 0.001	0.06 $\pm$ 0.003	0.03 $\pm$ 0.009	0.001 $\pm$ 0.00005	0.71 $\pm$ 0.17
JF	5	4.4 $\pm$ 3.0	70.8 $\pm$ 19.7	0.01	0.2 $\pm$ 0.3	0.003 $\pm$ 0.004	0.04 $\pm$ 0.41	0.01 $\pm$ 0.007	0.02 $\pm$ 0.01	0.007 $\pm$ 0.003	nd	0.33 $\pm$ 0.34
JM	11	6.5 $\pm$ 4.2	57.8 $\pm$ 7.8	nd	0.2 $\pm$ 0.13	0.003 $\pm$ 0.001	0.05 $\pm$ 0.03	0.02 $\pm$ 0.01	0.02 $\pm$ 0.008	0.01 $\pm$ 0.004	0.001	0.31 $\pm$ 0.18
<i>Phocoena phocoena</i>	1	3.0	40.3	nd	0.48	0.004	0.08	0.04	0.02	0.009	nd	0.57
JM	1	3.0	40.3	nd	0.48	0.004	0.08	0.04	0.02	0.009	nd	0.57

nd = values were less than the limit of detection (see [Materials and methods section](#)).

The congener BDE85 was not detected in any individual analysed.

<sup>a</sup>  $\Sigma\text{BDE}$  includes 9 congeners. See [Material and methods](#) section for full list.

general, the  $\Sigma\text{PBDE}$  concentrations were higher in common dolphin than in harbour porpoise and congeners BDE28 and BDE 183 were detected only in common dolphins', indicating that common dolphin has difficulties with metabolizing PBDEs. However, we must consider these results as only indicative due to the low number of samples analysed for PBDEs.

### 3.2. Geographic comparison

The PCB concentrations of Iberian common dolphins from this study are in the range reported by previous published studies in the NWIP, in France and also in England. However these concentrations are much higher than in common dolphins from Ireland and those from the South Atlantic Ocean and from the Pacific Ocean (Table 3). It is important to note that the PCB concentration in the Irish common dolphins ( $2.8 \mu\text{g g}^{-1}$  lipid weight) is the mean value of only 5 PCB congeners: CB118, 138, 153, 180 and 170. In the present study the sum of these 5 PCB congeners is  $10.3 \mu\text{g g}^{-1}$  lipid weight. A different pattern was found when a comparison was made with common dolphins from the Mediterranean Sea and the east coast of the USA (NW Atlantic Ocean); common dolphins from these areas were more highly contaminated with PCBs than those from the NWIP (Table 3). The PBDE concentrations from male and female common dolphins in the NE Atlantic Ocean are lower than in other areas, especially when compared to dolphins from Korea in the Pacific Ocean.

Despite the small sample size for pilot whale, the PCB concentrations found in the NWIP are lower than in the rest of the NE Atlantic and Ligurian Sea, with the exception of one male specimen from Ireland. However, in this study carried out by Troisi et al. (1998) only 5 PCB congeners were analysed and the sum of these is also lower in Iberian pilot whales than in Ireland ( $8.3$  and  $10.2 \mu\text{g g}^{-1}$  lipid weight, respectively). In contrast, when the PCB concentrations in pilot whales from the NW Atlantic and the Pacific Oceans are compared with the data from this study, with the exception of females from Massachusetts (Tilbury et al., 1999), the Iberian pilot whales contain higher concentrations (Table 3).

Harbour porpoise and bottlenose dolphin contained the highest PCB concentrations among the Iberian toothed whales. Comparing with other areas across the world we observed that both species have in general higher concentrations than seen in conspecifics from the Atlantic, Pacific and Indian Oceans. There were exceptions; the North, Baltic and Norwegian Seas for harbour porpoise and the Mediterranean Sea for bottlenose dolphin. This indicates that those seas are highly contaminated with PCBs, as has been demonstrated by previously published studies (Fossi et al., 2013; Pierce et al., 2008; Weijs et al., 2009, 2010). Borrell et al. (2006) reported mean PCB concentrations in male and female bottlenose dolphins stranded in the South of the Iberian Peninsula that were two times higher than in the present study (Table 3). These

high values may be a result of the proximity of the study area used by Borrell et al. (2006) to the Mediterranean Sea. PBDE concentrations in the only male harbour porpoise analysed in the present study were lower than for males from other areas and, as for the PCBs, North Sea porpoises.

Striped dolphins presented the lowest PCB concentrations among the Iberian toothed whales but also when we compare with specimens from other areas of the Northern Hemisphere (Table 3). Only one specimen stranded on the coast of England showed similar concentrations to those of the Iberian striped dolphins (Morris et al., 1989). Regarding data from the Southern Hemisphere, the concentrations found in the present study are higher than those reported in dolphins stranded on the Brazilian coast. This difference in PCB levels between the Northern and Southern Hemispheres is common to all five species studied here, reflecting the highly industrialized development of the Northern Hemisphere. PBDE concentrations were not analysed in striped dolphins from the NWIP. However, previously published studies showed similar values across different areas of the Atlantic and Pacific Oceans, with the exception of one female stranded on the coast of Japan which contained only  $0.08 \mu\text{g g}^{-1}$  lipid wt. of PBDEs and also only  $3.2 \mu\text{g g}^{-1}$  lipid wt. of PCBs (Isobe et al., 2009).

### 3.3. Toxicological aspects

Reliable toxicity data for predatory marine mammals are scarce. Instead, threshold levels are often extrapolated from terrestrial species, since the effects of toxic compounds cannot be tested in free-living animals because such experimental manipulations raise ethical considerations (Das et al., 2003). Thus, although the validity of these extrapolations could be questionable, they can be justified by the current lack of better data. The harmful consequences of the bioaccumulation of POPs in marine mammals include depression of the immune system (e.g. de Swart et al., 1996; Ross, 1995), increased risk of infection and reproductive failure. Specifically, a total PCB concentration of  $17 \mu\text{g g}^{-1}$  lipid wt. has been reported as a threshold level above which there are health effects in mammals (Kannan et al., 2000). This threshold was obtained in laboratory mammals (seals, European otters and mink) fed with field food items. In this study the threshold value was frequently exceeded for all species, often with more than 50% of the individuals (except pilot whale). However, this value was exceeded by all the bottlenose dolphins and 75% of the harbour porpoises. This result is even more important when it is considered in association with the previous study carried out in the Northeast Atlantic, including samples from the NWIP. This showed that almost half of the harbour porpoises for which cause of death was determined as being from pathological causes, had significantly higher concentrations of all classes of POPs than animals dying from other causes (Pierce et al., 2008). PBDE concentrations measured in common dolphins and harbour porpoises from

**Table 3**  
Mean  $\pm$  SD of PCB and PBDE concentrations ( $\mu\text{g g}^{-1}$  lipid weight) in blubber of the five toothed whale species from all over the world. Sample size “n” of each species by sex and area is in brackets.

Species	Area	Sex	$\Sigma\text{PCB}$	$\Sigma\text{PBDE}$	References	
<i>Delphinus delphis</i>	NE Atlantic Ocean					
	NWIP	M	20.4 $\pm$ 16.1 (50)	0.51 $\pm$ 0.17 (13)	This study	
	NWIP	F	11.6 $\pm$ 7.2 (31)	0.205 $\pm$ 0.34 (6)	This study	
	Spain	F	10.9 $\pm$ 11.6 (23)	0.42 $\pm$ 0.18 (23)	Pierce et al. (2008)	
	Spain	M	37.8 $\pm$ 18.9 (33)		Borrell et al. (2001)	
	Spain	F	23.9 $\pm$ 17.7 (23)		Borrell et al. (2001)	
	France	F	13.7 $\pm$ 12.7 (36)	0.61 $\pm$ 0.41 (36)	Pierce et al. (2008)	
	Ireland	M	2.8 <sup>a</sup>		Troisi et al. (1998)	
	Ireland	F	3.6 $\pm$ 3.4 (11)	0.76 $\pm$ 0.5 (11)	Pierce et al. (2008)	
	England	F	20.2 $\pm$ 16.7 (43)		Law et al. (2013)	
	Mediterranean Sea					
	Spain	M	54.3 $\pm$ 22.7 (11)		Borrell et al. (2001)	
	Spain	F	23.8 $\pm$ 32.8 (11)		Borrell et al. (2001)	
	NW Atlantic Ocean					
	USA	M	36.5 $\pm$ 4.0 (4)		Kuehl et al. (1991)	
	SE Atlantic Ocean					
	South Africa	M	5.9 $\pm$ 5.05 (10)		de Kock et al. (1994)	
	South Africa	F	2.8 $\pm$ 1.7 (7)		de Kock et al. (1994)	
	SW Atlantic Ocean					
	Brazil <sup>*a</sup>	M	17.0 (1)		Kajiwara et al. (2004)	
	Brazil <sup>*b</sup>	M	14.6 $\pm$ 15.3 (2)		Lailson-Brito et al. (2012)	
	Brazil <sup>*c</sup>	F	2.2 $\pm$ 0.7 (2)		Lailson-Brito et al. (2012)	
	Pacific Ocean					
	NE Australia	F	0.6 (1)		Vetter et al. (2001)	
	New Zealand	M	0.8 $\pm$ 0.5 (12)		Stockin et al. (2007)	
	New Zealand	F	0.14 $\pm$ 0.13 (7)		Stockin et al. (2007)	
	Korea	M	15.0 $\pm$ 7.6 (12)	1.7 $\pm$ 0.78 (12)	Moon et al. (2010)	
	Korea	F	15.0 $\pm$ 7.8 (10)	1.6 $\pm$ 0.68 (10)	Moon et al. (2010)	
	<i>Globicephala melas</i>	NE Atlantic Ocean				
		NWIP	M	38.7 (1)		This study
		NWIP	F	4.9 $\pm$ 4.2 (2)		This study
		France	nd	189.0 $\pm$ 298.0 (7) <sup>b</sup>		Alzieu and Duguy (1979)
		Ireland	M	10.2 <sup>a</sup>		Troisi et al. (1998)
Faroe Islands		M	48.8 $\pm$ 23.1 (52)	2.4 (21)	Borrell (1993); Lindström et al. (1999)	
Faroe Islands		F	26.3 $\pm$ 23.1 (159)	1.6 (32)	Borrell (1993); Lindström et al. (1999)	
Ligurian Sea		nd	125.0 (1) <sup>b</sup>		Marsili and Focardi (1997)	
NW Atlantic Ocean						
Canada		M	9.0 $\pm$ 3.8 (5)		Muir et al. (1988)	
Canada		F	3.5 $\pm$ 3.3 (9)		Muir et al. (1988)	
Massachusetts		M	12 $\pm$ 2.7 (6) <sup>c</sup>		Tilbury et al. (1999)	
Massachusetts		F	6.1 $\pm$ 1.1 (16) <sup>c</sup>		Tilbury et al. (1999)	
Pacific Ocean						
Tasmania		M	0.41 $\pm$ 0.04 (21)		Weijs et al. (2013)	
Tasmania		F	0.36 $\pm$ 0.26 (33)		Weijs et al. (2013)	
<i>Phocoena phocoena</i>		NE Atlantic Ocean				
		NWIP	M	19.8 $\pm$ 20.8 (4)	0.57 (1)	This study
		NWIP	F	20.8 $\pm$ 21.6 (8)		This study
		NW Spain	F	5.3 $\pm$ 4.2 (3)	0.28 $\pm$ 0.04 (3)	Pierce et al. (2008)
		France	F	13.8 $\pm$ 10.6 (2)	1.4 $\pm$ 0.94 (2)	Pierce et al. (2008)
	Ireland	M	6.2 (1) <sup>a</sup>		Troisi et al. (1998)	
	Ireland	F	0.53 $\pm$ 0.5 (12)	0.66 $\pm$ 0.49 (12)	Pierce et al. (2008)	
	Scotland	M	13.1 (21) <sup>d</sup>		Wells et al. (1994)	
	Scotland	F	10.5 $\pm$ 13.1 (31)	1.4 $\pm$ 1.3 (31)	Pierce et al. (2008)	
	South England	M	23.4 $\pm$ 21.6 (2)	2.1 $\pm$ 1.6 (21) <sup>d</sup>	Law et al. (2006, 2010)	
	East England	M	11.6 $\pm$ 9.7 (23)		Law et al. (2010)	
	Southern North Sea	M	46.4 $\pm$ 30.7 (21)		Law et al. (2010)	
	Southern North Sea	F	15 $\pm$ 8.6 (19)	1.06 $\pm$ 0.8 (19)	Pierce et al. (2008)	
	Baltic Sea	M	31 $\pm$ 18.5 (17)		Berggren et al. (1999)	
	Norwegian Sea	M	15 $\pm$ 11 (8)		Berggren et al. (1999)	
	Faroe Islands	M	13.4 $\pm$ 2.4 (3)		Borrell (1993)	
	Faroe Islands	F	8.8 $\pm$ 1.05 (3)		Borrell (1993)	
	Greenland	M	2.4 (32)		Borrell et al. (2004)	
	Greenland	F	1.7 (43)		Borrell et al. (2004)	
	NW Atlantic Ocean					
	Canada	M	13.2 $\pm$ 9.9 (45)		Westgate et al. (1997)	
	Canada	F	9.3 $\pm$ 5 (43)		Westgate et al. (1997)	
	Boston	M	33.6 (2)		Tilbury et al. (1997)	
	Maine (Canada)	F	15.8 (1)		Tilbury et al. (1997)	
	Pacific Ocean					
	North Pacific	M	18.2 $\pm$ 16.6 (17)		Calambokidis and Barlow (1991)	
	North Pacific	F	11.7 $\pm$ 10.0 (26)		Calambokidis and Barlow (1991)	

Table 3 (continued)

Species	Area	Sex	ΣPCB	ΣPBDE	References
<i>Stenella coeruleoalba</i>	NE Atlantic Ocean				
	NWIP	M	22.8 ± 23.3 (8)		This study
	NWIP	F	7.6 ± 5.9 (7)		This study
	England and Wales		20.0 (1)	0.45 (1)	Morris et al. (1989); Law et al. (2003)
	Mediterranean Sea				
	Spain	M	65.6 ± 40.5 (15)		Borrell and Aguilar (2005)
	Spain	F	90.0 ± 70.7 (5)		Borrell and Aguilar (2005)
	France	M	57.7 ± 41.9 (19)		Wafo et al. (2012)
	France	F	45.3 ± 45.7 (12)		Wafo et al. (2012)
	Italy	M	215.3 (33) <sup>b</sup>		Marsili and Focardi (1997)
	Italy	F	92.8 (26) <sup>b</sup>		Marsili and Focardi (1997)
	NW Atlantic Ocean				
	USA	–	59.0 (3)	0.66 (1)	Taruski et al. (1975); Johnson-Restrepo et al. (2005)
	SW Atlantic Ocean				
	Brazil <sup>*</sup>	M	12.7 ± 9.6 (3)	0.81 ± 0.47 (3)	Leonel et al. (2012)
	Brazil <sup>*</sup>	F	5.3 ± 7.8 (6)	0.54 ± 0.45 (6)	Leonel et al. (2012)
	Pacific Ocean				
	Japan	M	25.3 ± 7.2 (20)	0.36 ± 0.28 (20)	Isobe et al. (2009)
	Japan	F	3.2 (1)	0.08 (1)	Isobe et al. (2009)
	<i>Tursiops truncatus</i>	NE Atlantic Ocean			
NWIP		M	62.1 ± 41.8 (4)		This study
NWIP		F	48.9 ± 30.9 (3)		This study
South Spain		M–F	182.6 ± 90.8 (5)		Borrell et al. (2006)
South Portugal		M–F	75.3 ± 39.4 (7)		Borrell et al. (2006)
Canary Islands		M	12.7 ± 10.7 (6)		Carballo et al. (2008)
Canary Islands		F	2.9 ± 4.3 (3)		Carballo et al. (2008)
West Ireland			5.01 ± 6.0 (6)		Berrow et al. (2002)
Scotland		M	13.6 (1) <sup>d</sup>		SOAFD data <sup>e</sup>
Scotland		F	11.0 (5) <sup>d</sup>		Wells et al. (1994)
Wales		M	290.0 (3)		Morris et al. (1989)
Wales		F	760.0 (1)		Morris et al. (1989)
Mediterranean Sea					
Spain		M–F	167.8 ± 99.5 (13)		Borrell et al. (2006)
Balearic Islands		M–F	117.3 ± 104.1 (7)		Borrell et al. (2006)
Italy		M	1192 (5) <sup>f</sup>		Corsolini et al. (1995)
Italy		F	587 (2) <sup>f</sup>		Corsolini et al. (1995)
NW Atlantic Ocean					
North Carolina		M	53.3 (17)		Hansen et al. (2004)
North Carolina		F	11.6 (14)		Hansen et al. (2004)
South Carolina		M	50.4 (5)	0.98 (35)	Hansen et al. (2004); Fair et al. (2010)
South Carolina		F	8 (6)	5.9 (11)	Hansen et al. (2004); Fair et al. (2010)
Florida		M	20 (9)	1.5 (31)	Hansen et al. (2004); Fair et al. (2010)
Florida		F	12.6 (2)	0.58 (15)	Hansen et al. (2004); Fair et al. (2010)
SE Atlantic Ocean					
Rio de Janeiro		M	11.8 ± 2.4 (2)		Lailson-Brito et al. (2012)
São Paulo		M	5.9 (1)	0.06 (1)	Yogui et al. (2010, 2011)
SW Atlantic Ocean					
South Africa	M	2.6 ± 3.3 (5)		de Kock et al. (1994)	
South Africa	F	1.6 (1)		de Kock et al. (1994)	
Pacific Ocean					
Australia	M–F	0.06 (6)		Kemper et al. (1994)	
Indian Ocean					
Bay of Bengal, India	M	1.2 (2)		Tanabe et al. (1993)	
Bay of Bengal, India	F	0.75 (2)		Tanabe et al. (1993)	

\* *Delphinus delphis* and *Stenella frontalis*.

<sup>a</sup> Sum of 5 PCB congeners: CB118, 138, 153, 180 and 170.

<sup>b</sup> Values expressed on a dry weight basis (from lyophilized tissue).

<sup>c</sup> SEM (standard error of the mean).

<sup>d</sup> As Aroclor 1254 formulation.

<sup>e</sup> SOAFD data (Scottish Office Agriculture and Fisheries Department).

<sup>f</sup> Values expressed on a wet weight basis.

this study are at least 10 times lower than those of PCBs, being slightly higher for harbour porpoise than common dolphin. There is still no information on a toxic threshold for PBDEs in marine mammals, although experimental exposure investigations revealed that PBDEs induce a wide variety of disorders in mammals (e.g. cancer, reproductive and developmental toxicity, endocrine disruption and central nervous system effects; Hana et al., 2004). As such it is not possible to say whether or not such concentrations are likely to impact on the toothed whales. However, what needs to be considered is that the PBDEs are present and have augmented the overall concentration of POPs in these whales.

In future studies consideration should be given to the possible cumulative impacts of the range of contaminants found in these marine animals.

#### 4. Conclusions

The inter-species differences found in the present study, covering the accumulation and patterns of POPs, illustrate the important influence of both the biological and ecological factors of each species in determining the contaminant loading. Overall, the POP concentrations observed in toothed whales from the NWIP were the same order of



magnitude or lower than those reported by previous studies in areas of the NE Atlantic. However, they were higher than those of toothed whales from the southern Atlantic Ocean and Pacific Ocean.

The bottlenose dolphin and harbour porpoise are recorded in the EU Habitats Directive as Species of Special Interest (Directive 92/43/CEE), the protection of which requires the designation of Special Areas of Conservation (SAC) by EU Member States. In the present study both species seem to be the more contaminated with PCBs as shown by their higher concentrations when compared to the other three species. Their PCB concentrations also raise particular concern as they are much higher than the threshold level for PCB concentrations associated with adverse health effects on marine mammals. This study adds to our knowledge of POP concentrations in marine mammals that frequent the NWIP waters, complementing the existing database on these chemical contaminants in this area. However, nowadays there remains a lack of information on concentrations of other organochlorine compounds of special concern, and this should be addressed through further studies. This is particularly relevant given that the European scientific community is developing research on large marine vertebrates as indicators of medium and long term marine environmental change for incorporation under the Marine Strategy Framework Directive. A complete database of chemical contaminants in marine mammals of the NWIP is an important step forward.

## Acknowledgements

The authors gratefully acknowledge the assistance of volunteers from the Galician (CEMMA) and Portuguese (SPVS) stranding networks. They also thank E. Dalgarno, L. Phillips, I. Hussy and J. A. Scurfield for their help with the organochlorine analysis. We would like to thank R. Gallois and C. Trichet, for their participation in the determination of total lipid content. This work was supported through the PhD grant of PMF from the Portuguese Foundation for Science and Technology of the Government of Portugal (SFRH/BD/36766/2007). GJP acknowledges support from the EU under the ANIMATE project (MEXC-CT-2006-042337). MBS was supported by the Spanish Ministry of Education, Programa Nacional de Movilidad de Recursos Humanos de Investigación (PR-2010-0518) and the LOTOPPEL project (Plan Nacional de I + D + I, CTM 2010–16053). Two anonymous reviewers and the associate editor F. Riget are thanked for helpful suggestions and comments on an earlier form of this manuscript.

## References

- Aguilar A. Organochlorine pollutants and cetaceans: a perspective. In: Evans PGH, Smeenk C, editors. European research on cetaceans. E-publishing Inc.; 1989. p. 10–1.
- Aguilar A, Borrell A, Pastor T. Biological factors affecting variability of persistent pollutant levels in cetaceans. *J Cetac Res Manage* 1999;1:83–116.
- Alzieu C, Duguy R. Teneurs en composés organochlorés chez les cétacés et pinnipèdes fréquentant les côtes françaises. *Oceanol Acta* 1979;2:107–20.
- Berggren P, Ishaq R, Zebühr Y, Näf C, Bandh C, Broman D. Patterns and levels of organochlorines (DDTs, PCBs, non-ortho PCBs and PCDD/Fs) in male harbour porpoises (*Phocoena phocoena*) from the Baltic Sea, the Kattegat-Skagerrak Seas and the West Coast of Norway. *Mar Pollut Bull* 1999;38:1070–84.
- Berrow SD, Mchugh B, Glynn D, McGovern E, Parsons KM, Baird RW, et al. Organochlorine concentrations in resident bottlenose dolphins (*Tursiops truncatus*) in the Shannon estuary, Ireland. *Mar Pollut Bull* 2002;44:1296–313.
- Beyer A, Mackay D, Matthies M, Wania F, Webster E. Assessing long-range transport potential of persistent organic pollutants. *Environ Sci Technol* 2000;34:699–703.
- Boon JP, van Arnhem E, Jansen S, Kannan N, Petrick G, Schulz DE, et al. The toxicokinetics of PCBs in marine mammals with special reference to possible interactions of individual congeners with the cytochrome P450-dependent monooxygenase system—an overview. In: Walker CH, Livingstone DR, Lipnick RL, editors. Persistent pollutants in marine ecosystems. Oxford, New York, Seoul, Tokyo: Pergamon Press; 1992. p. 119–59.
- Boon JP, Lewis WE, Tjoen-A-Choy MR, Allchin CR, Law RJ, Jacob de Boer, et al. Levels of polybrominated diphenylether (PBDE) flame retardants in animals representing different trophic levels of the North Sea food web. *Environ Sci Technol* 2002;36:4025–32.
- Borrell A. PCB and DDTs in blubber of cetaceans from the northeastern north Atlantic. *Mar Pollut Bull* 1993;26:146–51.
- Borrell A, Aguilar A. Differences in DDT and PCB residues between common and striped dolphins from the southwestern Mediterranean. *Arch Environ Contam Toxicol* 2005;48:501–8.
- Borrell A, Aguilar A, Corsolini S, Focardi S. Evaluation of toxicity and sex-related variation of PCB levels in Mediterranean striped dolphins affected by an epizootic. *Chemosphere* 1996;32:2359–69.
- Borrell A, Cantos G, Pastor T, Aguilar A. Organochlorine compounds in common dolphin (*Delphinus delphis*) from the Atlantic and Mediterranean waters of Spain. *Environ Pollut* 2001;114:265–74.
- Borrell A, Aguilar A, Cantos G, Lockyer C, Heide-Jørgensen M, Jensen J. Organochlorine residues in harbour porpoises from Southwest Greenland. *Environ Pollut* 2004;128:381–91.
- Borrell A, Aguilar A, Tornero V, Sequeira M, Fernandez G, Alis S. Organochlorine compounds and stable isotopes indicate bottlenose dolphin subpopulation structure around the Iberian Peninsula. *Environ Int* 2006;32:516–23.
- Calambokidis J, Barlow J. Chlorinated hydrocarbon concentrations and their use for describing population discreteness in harbour porpoises from Washington, Oregon, and California. In: Reynolds JE, Oddi DK, editors. Marine mammal strandings in the United States. NOAA Technical Report NMFS 98. Silver Spring, MD, USA: US Department of Commerce, Natural Marine Fisheries Service; 1991.
- Carballo M, Arbelo M, Esperón F, Méndez M, de La Torre A, Muñoz MJ. Organochlorine residues in the blubber and liver of bottlenose dolphins (*Tursiops truncatus*) stranded in the Canary Islands, North Atlantic Ocean. *Environ Toxicol* 2008;23:200–10.
- Carro N, García I, Ignacio M, Mouteira A. Levels and distribution patterns of polychlorinated biphenyl congeners in surface sediments from Galicia coast (Northwestern, Spain) according to granulometric characteristics. *Environ Technol* 2002;23:919–30.
- Cheeseman RV, Wilson AL. A manual on quality control for the water industry. WRc Technical Report No 66; 1989.
- Corsolini S, Focardi S, Kannan K, Tanabe S, Borrell A, Tatsukawa R. Congener profile and toxicity assessment of polychlorinated biphenyls in dolphins, sharks and tuna collected from Italian coastal waters. *Mar Environ Res* 1995;40:33–53.
- Covaci A, Van de Vijver I, De Coen W, Das K, Bouquegneau JM, Blust R, et al. Determination of organohalogenated contaminants in liver of harbour porpoises stranded on the Belgian North Sea coast. *Mar Pollut Bull* 2002;44:1156–64.
- Das K, Debacker V, Pillet S, Bouquegneau JM. Heavy metals in marine mammals. In: Vos JG, Bossart GD, Fournier M, O'Shea TJ, editors. Toxicology of marine mammals volume 3 – systems. New York: Taylor & Francis Inc.; 2003. p. 135–67.
- de Boer J, Wester PG, Klamer HJC, Lewis WE, Boon JP. Do flame retardants threaten ocean life? *Nature* 1998;394:28–9.
- de Kock AC, Best PB, Cockcroft V, Bosma C. Persistent organochlorine residues in small cetaceans from the east and west coasts of southern Africa. *Sci Total Environ* 1994;154:153–62.
- de Swart RL, Ross PS, Vos JG, Osterhaus ADME. Impaired immunity in harbour seals (*Phoca vitulina*) exposed to bioaccumulated environmental contaminants: review of a long-term feeding study. *Environ Health Perspect* 1996;823–8.
- Dorneles PR, Lailson-Brito J, Durtu AC, Weijs L, Azevedo AF, Torres JPM, et al. Anthropogenic and naturally-produced organobrominated compounds in marine mammals from Brazil. *Environ Int* 2010;36:60–7.
- Duinker JC, Hillebrand MTJ, Zeinstra T, Boon JP. Individual chlorinated biphenyls and pesticides in tissues of some cetacean species from the North Sea and the Atlantic Ocean; tissue distribution and biotransformation. *Aquat Mamm* 1989;15:95–124.
- Fair PA, Adams J, Mitchum G, Hulsey TC, Reif JS, Houde M, et al. Contaminant blubber burdens in Atlantic bottlenose dolphins (*Tursiops truncatus*) from two southeastern US estuarine areas: concentrations and patterns of PCBs, pesticides, PBDEs, and PAHs. *Sci Total Environ* 2010;408:1577–97.
- Figueiras FG, Labarta U, Fernández-Reiriz JM. Coastal upwelling, primary production and mussel growth in the Rias Baixas of Galicia. *Hydrobiologia* 2002;484:121–31.
- Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* 1957;226:497–509.
- Fossi MC, Panti C, Marsili L, Maltese S, Spinsanti G, Casini S, et al. The Pelagos Sanctuary for Mediterranean marine mammals: marine protected area (MPA) or marine polluted area? The case study of the striped dolphin (*Stenella coeruleoalba*). *Mar Pollut Bull* 2013;70:64–72.
- Hana RP, Stephen B, Richard JA, Carol JE. Toxicological profile for polybrominated diphenyl ethers Agency for Toxic Substances and Disease Registry (ATSDR). Atlanta, Georgia: Division of Toxicology/Toxicology Information Branch; 2004. p. 619.
- Hansen LJ, Schwacke LH, Mitchum GB, Hohn AA, Wells RS, Zolman ES, et al. Geographic variation in polychlorinated biphenyl and organochlorine pesticide concentrations in the blubber of bottlenose dolphins from the US Atlantic coast. *Sci Total Environ* 2004;319:147–72.
- Hohn AA, Lockyer C. Protocol for obtaining age estimates from harbour porpoise teeth. Appendix 3. Report of the harbour porpoise age determination workshop. In: Bjørge A, Donovan GP, editors. Biology of phocoenids. Cambridge: Rep Int Whal Commn; 1995.
- Isobe T, Ochi Y, Ramu K, Yamamoto T, Tajima Y, Yamada TK, et al. Organohalogen contaminants in striped dolphins (*Stenella coeruleoalba*) from Japan: present contamination status, body distribution and temporal trends (1978–2003). *Mar Pollut Bull* 2009;58:396–401.
- Jepson PD, Bennett PM, Deaville R, Allchin CR, Baker JR, Law RJ. Relationships between polychlorinated biphenyls and health status in harbor porpoises (*Phocoena phocoena*) stranded in the United Kingdom. *Environ Toxicol Chem* 2005;24:238–48.
- Johnson-Restrepo B, Kannan K, Addink R, Adams DH. Polybrominated diphenyl ethers and polychlorinated biphenyls in a marine food web of coastal Florida. *Environ Sci Technol* 2005;39:8243–50.

- Kajiwaru N, Matsuoka S, Iwata H, Tanabe S, Rosas FCW, Fillmann G, et al. Contamination by persistent organochlorines in cetaceans incidentally caught along Brazilian coastal waters. *Arch Environ Contam Toxicol* 2004;46:124–34.
- Kannan K, Blakenship AL, Jones PD, Giesy JP. Toxicity reference values for the toxic effects of polychlorinated biphenyls to aquatic mammals. *Hum Ecol Risk Assess* 2000;6:181–201.
- Kemper C, Gibbs P, Obendorf D, Marvanek S, Lenghaus C. A review of heavy metal and organochlorine levels in marine mammals in Australia. *Sci Total Environ* 1994;154:129–39.
- Kuehl DW, Haebler R, Potter CW. Chemical residues in dolphins from the U.S. Atlantic coast including Atlantic bottlenose obtained during the 1987/88 mass mortality. *Chemosphere* 1991;22:1071–84.
- Kuiken T, Garcia Hartmann M. Proceedings of the first European Cetacean Society workshop on 'Cetacean pathology: dissection techniques and tissue sampling'. *ECS Newsletter* 1991;17:1–39.
- Lailson-Brito J, Dorneles PR, Azevedo-Silva CE, Bisi TL, Vidal LG, Legat LN, et al. Organochlorine compound accumulation in delphinids from Rio de Janeiro State, southeastern Brazilian coast. *Sci Total Environ* 2012;433:123–31.
- Law RJ, Alaei M, Allchin CR, Boon JP, Lebeuf M, Lepom P, et al. Levels and trends of polybrominated diphenylethers and other brominated flame retardants in wildlife. *Environ Int* 2003;29:757–70.
- Law RJ, Jepson PD, Deaville R, Reid RJ, Patterson IAP, Allchin CR, et al. Collaborative UK Marine Mammals Strandings Project: summary of contaminant data for the period 1993–2001. *Cefas Lowestoft: Science Series Technical Report*; 2006.
- Law RJ, Bersuder P, Barry J, Deaville R, Reid RJ, Jepson PD. Chlorobiphenyls in the blubber of harbour porpoises (*Phocoena phocoena*) from the UK: levels and trends 1991–2005. *Mar Pollut Bull* 2010;60:470–3.
- Law RJ, Bersuder P, Barry J, Barber J, Deaville R, Barnett J, et al. Organochlorine pesticides and chlorobiphenyls in the blubber of bycaught female common dolphins from England and Wales from 1992–2006. *Mar Pollut Bull* 2013;69:238–42.
- Leonel J, Taniguchi S, Sasaki DK, Cascaes MJ, Dias PS, Botta S, et al. Contamination by chlorinated pesticides, PCBs and PBDEs in Atlantic spotted dolphin (*Stenella frontalis*) in western South Atlantic. *Chemosphere* 2012;86:741–6.
- Lindström G, Wingfors H, Dam M, Bavel B. Identification of 19 polybrominated diphenyl ethers (PBDEs) in long-finned pilot whale (*Globicephala melas*) from the Atlantic. *Arch Environ Contam Toxicol* 1999;36:355–63.
- Lockyer C. A report on patterns of deposition of dentine and cement in teeth of pilot whales, genus *Globicephala*. *Rep Int Whal Commn Special Issue* 1993;14:138–61.
- López A, Santos MB, Pierce GJ, Gonzalez AF, Valeiras X, Guerra A. Trends in strandings of cetaceans on the Galician coast, NW Spain, during the 1990s. *Mar Biol Assoc UK* 2002;82:3916/1–9.
- López A, Pierce GJ, Santos MB, Gracia J, Guerra A. Fishery by-catches of marine mammals in Galician waters: results from on-board observations and an interview survey of fishermen. *Conserv Biol* 2003;11:25–40.
- López A, Pierce GJ, Valeiras X, Santos MB, Guerra A. Distribution patterns of small cetaceans in Galician waters. *J Mar Biol Assoc UK* 2004;84:1–13.
- Marsili L, Focardi S. Chlorinated hydrocarbon (HCB, DDTs and PCBs) levels in cetaceans stranded along the Italian coasts: an overview. *Environ Monit Assess* 1997;45:129–80.
- Méndez-Fernandez P, Pierce GJ, Bustamante P, Chouvelon T, Ferreira M, López A, et al. Foraging ecology of five toothed whale species in the Northwest Iberian Peninsula, inferred using carbon and nitrogen isotope ratios. *J Exp Mar Biol Ecol* 2012;413:150–8.
- Méndez-Fernandez P, Pierce GJ, Bustamante P, Chouvelon T, Ferreira M, González AF, et al. Ecological niche segregation among five toothed whale species off the NW Iberian Peninsula using ecological tracers as multi-approach. *Mar Biol* 2013:1–16.
- Méndez-Fernandez P, Webster L, Chouvelon T, Bustamante P, Ferreira M, González AF, et al. An assessment of contaminant concentrations in toothed whale species of the NW Iberian Peninsula: Part II. Trace element concentrations. *Sci Total Environ* 2014;484:206–17.
- Moon HB, Kannan K, Choi M, Yu J, Choi HG, An Y-R, et al. Chlorinated and brominated contaminants including PCBs and PBDEs in minke whales and common dolphins from Korean coastal waters. *J Hazard Mater* 2010;179:735–41.
- Morris RJ, Law RJ, Allchin CR, Kelly CA, Fileman CF. Metals and organochlorines in dolphins and porpoises of Cardigan Bay, West Wales. *Mar Pollut Bull* 1989;20:512–23.
- Muir DCG, Wagemann R, Crift NR, Norstrom RJ, Simon M, Lien J. Organochlorine chemical and heavy metal contaminants in white-beaked dolphins (*Lagenorhynchus albirostris*) and pilot whales (*Globicephala melana*) from the coast of Newfoundland, Canada. *Arch Environ Contam Toxicol* 1988;17:613–29.
- Murphy S, Pierce GJ, Law RJ, Bersuder P, Jepson PD, Learmonth JA, et al. Assessing the effect of persistent organic pollutants on reproductive activity in common dolphins and harbour porpoises. *NAFO* 2010;42:153–73.
- Nyman M, Koistinen J, Fant ML, Vartiainen T, Helle E. Current levels of DDT, PCB and trace elements in the Baltic ringed seals (*Phoca hispida baltica*) and grey seals (*Halichoerus grypus*). *Environ Pollut* 2002;119:399–412.
- OSPAR. Quality Status Report 2010. London: OSPAR Commission; 2010.
- Pierce GJ, Santos MB, Murphy S, Learmonth JA, Zuur AF, Rogan E, et al. Bioaccumulation of persistent organic pollutants in female common dolphins (*Delphinus delphis*) and harbour porpoises (*Phocoena phocoena*) from western European seas: geographical trends, causal factors and effects on reproduction and mortality. *Environ Pollut* 2008;153:401–15.
- Pierce DCJ, Caldas M, Cedeira J, Santos MB, Llavona A, Covelo P, et al. Trends in cetacean sightings along the Galician coast, north-west Spain, 2003–2007, and inferences about cetacean habitat preferences. *J Mar Biol Assoc UK* 2010;90:1547–60.
- Prego R, Cobelo-García A. Twentieth century overview of heavy metals in the Galician Ria (NW Iberian Peninsula). *Environ Pollut* 2003;121:425–52.
- R Development Core Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2010 ([www.R-project.org](http://www.R-project.org)).
- Reijnders PJH, Donovan GP, Aguilar A, Bjørge A. Report of the workshop on chemical pollution and cetaceans. *J Cetacean Res Manag* 1999;1:1–42.
- Ross PS. Seals, pollution and disease: environmental contaminant induced immunosuppression (PhD dissertation) Utrecht: Universiteit Utrecht; 1995.
- Santos MB, Fernandez R, López A, Martínez JA, Pierce GJ. Variability in the diet of bottlenose dolphin, *Tursiops truncatus*, in Galician waters, north-western Spain, 1990–2005. *J Mar Biol Assoc UK* 2007;87:231–41.
- Santos MB, German I, Correia D, Read FL, Martínez Cedeira J, Caldas M, et al. Long-term variation in common dolphin diet in relation to prey abundance. *Mar. Ecol. Prog. Ser.* 2013a;481:249–68.
- Santos MB, Monteiro SS, Vingada JV, Ferreira M, López A, Reid RJ, et al. Patterns and trends in the diet of long-finned pilot whales (*Globicephala melas*) in the Northeast Atlantic. *Mar Mamm Sci* 2013b. <http://dx.doi.org/10.1111/mms.12015>.
- Santos MB, Saavedra C, Pierce GJ. Quantifying the predation on sardine and hake by cetaceans in the Atlantic waters of the Iberian Peninsula. *Deep Sea Res II* 2013c. <http://dx.doi.org/10.1016/j.dsr2.2013.09.040>.
- Stockin KA, Law RJ, Duignan PJ, Jones GW, Porter L, Mirimim L, et al. Trace elements, PCBs and organochlorine pesticides in New Zealand common dolphins (*Delphinus* sp.). *Sci Total Environ* 2007;387:333–45.
- Storelli MM. Potential human health risks from metals (Hg, Cd, and Pb) and polychlorinated biphenyls (PCBs) via seafood consumption: estimation of target hazard quotients (THQs) and toxic equivalents (TEQs). *Food Chem Toxicol* 2008;46:2782–8.
- Tanabe S, Watanabe S, Kan H, Tatsukawa R. Capacity and mode of PCB metabolism in small cetaceans. *Mar Mamm Sci* 1988;4:103–24.
- Tanabe S, Subramanian AN, Ramesh A, Kumaran PL, Miyazaki N, Tatsukawa R. Persistent organochlorine residues in dolphins from the Bay of Bengal, south India. *Mar Pollut Bull* 1993;26:311–6.
- Taruski AG, Olney CE, Winn HE. Chlorinated hydrocarbons in cetaceans. *J Fish Res Board Can* 1975;32:2205–9.
- Tilbury KL, Stein JE, Meador JP, Krone CA, Chan SL. Chemical contaminants in harbor porpoise (*Phocoena phocoena*) from the North Atlantic coast: tissue concentrations and intra- and inter-organ distribution. *Chemosphere* 1997;34:2159–81.
- Tilbury KL, Adams NG, Krone CA, Meador JP, Early G, Varanasi U. Organochlorines in stranded pilot whales (*Globicephala melana*) from the coast of Massachusetts. *Arch Environ Contam Toxicol* 1999;37:125–34.
- Tornero V, Borrell A, Aguilar A, Forcada J, Lockyer C. Organochlorine contaminant and retinoid levels in blubber of common dolphins (*Delphinus delphis*) off northwestern Spain. *Environ Pollut* 2006;140:312–21.
- Troisi GM, Haraguchi K, Simmonds MP, Mason CF. Methyl sulphone metabolites of polychlorinated biphenyls (PCBS) in cetaceans from the Irish and the Aegean Seas. *Arch Environ Contam Toxicol* 1998;35:121–8.
- Vetter W, Scholz E, Gaus C, Müller JF, Haynes D. Anthropogenic and natural organohalogen compounds in blubber of dolphins and dugongs (*Dugong dugon*) from Northeastern Australia. *Arch Environ Contam Toxicol* 2001;41:221–31.
- Wafo E, Sarrazin L, Diana C, Dhermain F, Schembri T, Lagadec V, et al. Accumulation and distribution of organochlorines (PCBs and DDTs) in various organs of *Stenella coeruleoalba* and a *Tursiops truncatus* from Mediterranean littoral environment (France). *Sci Total Environ* 2005;348:115–27.
- Wafo E, Risoul V, Schembri T, Lagadec V, Dhermain F, Mama C, et al. PCBs and DDTs in *Stenella coeruleoalba* dolphins from the French Mediterranean coastal environment (2007–2009): current state of contamination. *Mar Pollut Bull* 2012;64:2535–41.
- Walsham P, Webster L, Russell M, Dalgarno EJ, Moffat CF. Development and evaluation of an accelerated solvent extraction (ASE) method for the analysis of organic contaminants in biota. *Organohalogen Compd* 2006;68:2485–8.
- Weijs L, Dirtu AC, Das K, Gheorghe A, Reijnders PJH, Neels H, et al. Inter-species differences for polychlorinated biphenyls and polybrominated diphenyl ethers in marine top predators from the Southern North Sea: part 1. Accumulation patterns in harbour seals and harbour porpoises. *Environ Pollut* 2009;157:437–44.
- Weijs L, Tibax D, Roach AC, Manning TM, Chapman JC, Edge K, et al. Assessing levels of halogenated organic compounds in mass-stranded long-finned pilot whales (*Globicephala melas*) from Australia. *Sci Total Environ* 2013;461–462:117–25.
- Wells DE, Campbell LA, Ross HM, Thompson PM, Lockyer CH. Organochlorine residues in harbour porpoise and bottlenose dolphins stranded on the coast of Scotland, 1988–1991. *Sci Total Environ* 1994;151:77–99.
- Westgate AJ, Muir DCG, Gaskin DE, Kingsley MCS. Concentrations and accumulation patterns of organochlorine contaminants in the blubber of harbour porpoises, *Phocoena phocoena*, from the coast of Newfoundland, the Gulf of St Lawrence and the Bay of Fundy/Gulf of Maine. *Environ Pollut* 1997;95:105–19.
- Wolkers H, Lydersen C, Kovacs KM. Accumulation and lactational transfer of PCBs and pesticides in harbour seals (*Phoca vitulina*) from Svalbard, Norway. *Sci Total Environ* 2004;319:137–46.
- Yogui GT, Santos MCO, Bertozzi CP, Montone RC. Levels of persistent organic pollutants and residual pattern of DDTs in small cetaceans from the coast of São Paulo, Brazil. *Mar Pollut Bull* 2010;60:1862–7.
- Yogui GT, Santos MCO, Bertozzi CP, Sericano JL, Montone RC. PBDEs in the blubber of marine mammals from coastal areas of São Paulo, Brazil, southwestern Atlantic. *Mar Pollut Bull* 2011;62:2666–70.
- Zegers BN, Mets A, Van Bommel R, Minkenberg C, Hamers T, Kamstra JH, et al. Levels of hexabromocyclododecane in harbor porpoises and common dolphins from western European seas, with evidence for stereoisomer-specific biotransformation by cytochrome P450. *Environ Sci Technol* 2005;39:2095–100.