

Trace elements in tissues of white-chinned petrels (*Procellaria aequinoctialis*) from Kerguelen waters, Southern Indian Ocean

Caio V. Z. Cipro · Yves Cherel · Florence Caurant ·
Pierre Miramand · Paula Méndez-Fernandez ·
Paco Bustamante

Received: 22 July 2013/Revised: 24 February 2014/Accepted: 25 February 2014/Published online: 7 March 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract The use of seabirds to assess marine contamination by trace elements in areas remote from pollutant emission points has already been done at various latitudes. Nevertheless, little information is available concerning the Southern Indian Ocean. Determining the contaminants levels, there appears necessary not only due to several deleterious effects reported in literature, but also as previous studies have highlighted elevated concentrations of cadmium (Cd) and mercury (Hg) in mollusks, crustaceans and fish. Within this context, the white-chinned petrel appears as a key species due to its lifespan, diet and trophic position. Thirty-three accidentally killed (collision with lights/bycatch in longline vessels) individuals collected in Kerguelen waters were analysed for Cd, copper (Cu), Hg, selenium (Se) and zinc (Zn) in liver, kidney, pectoral muscle, feathers and for mature males, testis. Elevated Hg concentrations (average $58.4 \mu\text{g g}^{-1}$ dw in liver) are likely due to the presence of mesopelagic prey in the diet of *Procellaria aequinoctialis*. Cd concentrations (average of $65.7 \mu\text{g g}^{-1}$ dw in kidney) can be attributed to a high level of fisheries offal consumption, as well as crustacean and squid ingestion. Correlation of Hg with Se indicates its detoxification by co-precipitation, and correlation of Cd with Zn suggests its displacement by Cd on

metallothioneins binding sites. This work also indirectly confirms ecological data (range and diet composition) from the wintering period of the species, which is rather scarce. Seasonal diet change and moulting accounted more for the obtained results than sex of the birds.

Keywords Heavy metals · Seabirds · Procellariiformes · Southern Ocean · Sub-Antarctic Islands

Introduction

Contamination of the marine environment is a problem of concern at the worldwide scale. Oceanic ecosystems are often considered less impacted than coastal ones because they are not directly submitted to river influxes, runoff and point-source pollution from urban, industrial and agricultural emissions (Mailman 1980). Nonetheless, many of these contaminants, e.g. persistent organic pollutants or mercury (Hg), are transported by the atmosphere all over the world, and they deposit into oceanic environments (Fitzgerald and Engstrom 1998). In the aquatic environment, contaminants such as trace elements are more mobile than in terrestrial ones (Selin 2009), and species foraging in these environments could, therefore, be considered more vulnerable.

Trace elements encompass essential elements involved in physiological and biochemical processes (Abdulla and Chmielnicka 1990) that can, however, be toxic when exposure and assimilation are excessive. Others, such as cadmium (Cd) and Hg, are nonessential elements that are potentially toxic even at relatively low concentrations. Exposure of marine organisms is governed by various factors, including foraging location and trophic position (Anderson et al. 2010). Among trace elements, Cd and Hg

C. V. Z. Cipro (✉) · F. Caurant · P. Miramand ·
P. Méndez-Fernandez · P. Bustamante
Littoral Environnement et Sociétés (LIENSs), UMR 7266,
CNRS-Université de La Rochelle, 2 rue Olympe de Gouges,
17042 La Rochelle Cedex 01, France
e-mail: caiovzc@gmail.com

C. V. Z. Cipro · Y. Cherel
Centre d'Etudes Biologiques de Chizé, UPR 1934 du Centre
National de la Recherche Scientifique, BP 14,
79360 Villiers-en-Bois, France

are known to bioaccumulate in food webs, resulting to high concentrations in long-lived top predator species such as marine birds. This process is particularly exacerbated at high latitudes (Elliott and Scheuhammer 1997; Dietz et al. 1998), where the baseline concentrations are probably higher than at more temperate latitudes (Bocher et al. 2003). Food constitutes the main pathway of exposure of seabirds to contaminants (e.g. Ochoa-acuña et al. 2002). Nevertheless, numerous factors such as phylogeny, moult pattern, sex, life span and diet composition are likely to influence trace element bioaccumulation in seabird tissues, with Cd and Hg concentrations showing the largest variation among seabird species (Walsh 1990; Stewart et al. 1999). Because seabirds are long-lived species situated at the top of the food web, they are also submitted to the process of biomagnification of contaminants (Furness and Camphuysen 1997). For these reasons, marine birds are considered to be useful bioindicators of environmental contamination (Monteiro and Furness 1995; Furness and Camphuysen 1997). They seem to be particularly appropriate to survey the health status of marine ecosystems as they generally display lower coefficients of variation in contaminant levels than fish or marine mammals (Gilbertson et al. 1987). Furthermore, they are reflecting large areas integrating contaminants even from remote places that would be otherwise difficult to sample (Stewart et al. 1999; Carravieri et al. 2013). The use of seabirds to monitor marine contamination by trace elements in areas remote from pollutant emission points has already been done at various latitudes (Fitzgerald and Engstrom 1998; Kojadinovic et al. 2007; Anderson et al. 2010). However, information concerning the Southern Indian Ocean is still scarce. Only one study reported trace element concentrations in liver, kidney and muscle for five small petrels, prions and diving petrels from the Kerguelen Islands, all of them mainly feeding on zooplankton (Bocher et al. 2003). The study highlighted that the diet was a key factor driving the exposure and trace element levels in the tissues of these birds. More recently, Hg concentrations were determined in feathers from a large number of species, revealing a strong influence of the feeding behaviour on Hg concentrations (Blévin et al. 2013; Carravieri et al. 2013). Identification of high levels of Cd and Hg in mollusks, crustaceans and fish from the Kerguelen waters (Bustamante et al. 1998a, b, 2003; Bocher et al. 2003) would result in elevated exposure of seabirds. Moreover, several deleterious effects of such elements have already been reported in birds, for instance: decrease in egg laying, egg hatchability and increase in embryonic mortality, and besides neurotoxicity, there is also evidence for immunotoxic effects, such as decreased immune response and histological changes (Scheuhammer et al. 2007; Goutte et al. 2014) and several endocrine effects as well (e.g. Tan

et al. 2009; Tartu et al. 2013). In this context, the monitoring of the contaminants in the Southern Indian Ocean appears therefore a matter of importance.

Procellaria aequinoctialis, even though one of the most abundant seabirds in the Southern Ocean is classified as vulnerable in the IUCN red list since literature presents historical declines of populations within the Southern Ocean, namely at Crozet (Barbraud et al. 2008) and South Georgia (Berrow et al. 2000), but not conclusively in Kerguelen Islands (Barbraud et al. 2009). Very high rates of incidental mortality in longline fisheries have been recorded in recent years (Birdlife International 2013). The probability that these circumstances will continue and its susceptibility to predation and the degradation of breeding habitat indicate that a rapid and on-going population decline is likely (Birdlife International 2013). Specifically, for the Kerguelen waters, the mortality caused by fisheries constitutes a serious threat for *P. aequinoctialis* at least at the regional scale of the Southern Indian Ocean (Barbraud et al. 2009). A similar conclusion was taken for the Crozet Islands population, even though estimates of the number of birds killed by the longline fishing vessels remained within the threshold levels (estimated around 8,000 individuals per annum in this specific case) after 2003 (Barbraud et al. 2008). Association with commercial fisheries is due whether because *P. aequinoctialis* target the same areas of high productivity than fishermen and/or because fisheries appear to represent an additional potential food source for these opportunistic seabirds in the form of offal and discards of nontarget species (Delord et al. 2010). In their wintering grounds, i.e. the eastern boundary upwelling system of the Benguela current (Péron et al. 2010), *P. aequinoctialis* populations from the Kerguelen waters mainly feed on offal, i.e. fish heads and guts scavenged from commercial trawlers (Jackson 1988). Nevertheless, their diet also includes “natural” pelagic fish prey, crustaceans and cephalopods, in this order. Prey given to chicks during the breeding season is mainly composed of fish, followed by cephalopods and then crustaceans (Delord et al. 2010). In the Crozet Islands (3.2° north from Kerguelen), the diet of *P. aequinoctialis* is somewhat similar with fish prevailing, especially mesopelagic species (Ridoux 1994; Catard et al. 2000; Connan et al. 2007). However, crustaceans and cephalopods are important prey as well.

In this study, we examine the levels of Cd, copper (Cu), Hg, selenium (Se) and zinc (Zn) concentrations in the feathers, liver, kidney and muscle tissues (and also testis in a smaller extent) of the white-chinned petrel *P. aequinoctialis* from Kerguelen waters using only incidentally killed animals. The second objective of this work was to make the relationship between trace element levels in internal tissues, reflecting the risk of toxicity for the birds and their levels in the feathers that, if related to the internal




Fig. 1 Kerguelen location in the Southern Indian Ocean among the French Southern Lands (*left*) and detailed map (*right*). STZ, SAZ and AZ stand for, respectively, the subtropical, sub-Antarctic and Antarctic zones; whereas STF and PF stand for the subtropical and polar fronts

organs, could provide a wider nondestructive monitoring basis (Agusa et al. 2005).

Materials and methods

Sampling

Sample collection occurred from October 1998 to March 2000 at Kerguelen Archipelago, Southern Indian Ocean (Fig. 1). Thirty-three white-chinned petrels were collected as bycatch of longline fishing vessels or found recently dead on land killed following collision with the Scientific Station lights. Only intact or almost intact birds were considered for this study. The petrels were weighed in the field, immediately frozen and kept at -20°C until analyses. Birds were thawed overnight and dissected for liver, kidney, muscle (pectoral), testis (only on three mature males) and body feather removal, as body feathers (mainly pectoral ones, avoiding brood patch areas) give the best measure of the Hg contamination of a bird (Furness and Camphuysen 1997). They were also sexed by visual observation of the gonads: 6 individuals could be identified as females and 12 as males.

Trace element analyses

Analyses were performed as described by Bocher et al. (2003): for Cd, Cu and Zn determination, two aliquots of approximately 200–300 mg of each homogenised dried sample were digested in 5 ml of suprapure 14 N nitric acid at 60°C on a hot plate until the solution was clear. After evaporation, the residues were dissolved in 10 ml of 0.3 N suprapure nitric acid. Trace elements were determined by flame atomic absorption spectrophotometry (AAS) with a Varian spectrophotometer Spectra 250 Plus with a

deuterium background correction (high Cd concentrations, Cu and Zn) and by graphite furnace AAS (Hitachi Z-5000) with Zeeman background correction (low Cd concentrations and Se). Feather analyses included a previous cleaning step in order to remove surface lipids and contaminants, by using a 2:1 chloroform/methanol solution followed by two successive methanol rinses. After cleaning, body feathers were oven-dried for 48 h at 50°C .

Mercury analysis was carried out with an Automatic Mercury Analyser spectrophotometer, ALTEC AMA 254, which does not require an acid digestion of the samples. Aliquots ranging from 10 to 50 mg of dried sample were directly analysed after being inserted in the oven of the apparatus. After drying, the samples were heated under an oxygen atmosphere for 3 min, and the Hg liberated and subsequently amalgamated on an Au-net. The net was then heated to liberate the collected Hg, which was measured by AAS.

Accuracy and reproducibility of the methods were tested using dogfish liver (DOLT-2) and muscle (DORM-2) and lobster hepatopancreas (TORT-2) (National Research Council, Canada) reference standards. Standard and blanks were analysed along with each set of samples, and recoveries of the certified values and recoveries of the metals ranged from 92 to 105 %. Measurements were also validated by the IAEA inter-calibration exercise (Coquery et al. 2001). Concentrations are expressed in dry weight in order to compensate eventual moisture loss during freezing and to facilitate comparison between tissues and with other studies. Blanks were analysed at the beginning of each set of samples, and the detection limit of the method was $0.005\ \mu\text{g g}^{-1}$ dry mass.

Statistical analyses

Tests were performed using Microsoft Excel and Statsoft Statistica. Before analyses, data were checked for

Table 1 Trace element concentrations (mean \pm SD and range; $\mu\text{g g}^{-1}$ dw) in tissues of white-chinned petrels *P. aequinoctialis* from Kerguelen waters

Tissue	<i>n</i>	Cd	<i>n</i>	Cu	<i>n</i>	Hg	<i>n</i>	Se	<i>n</i>	Zn
Kidney	29	65.7 \pm 22.2	29	15.6 \pm 4.4	27	23.6 \pm 15.5	24	16.1 \pm 4.8	29	131 \pm 23
		26.4–121		11.8–35.7		8.91–91.8		7.5–25.2		96–200
Liver	26	20.5 \pm 6.5	26	16.2 \pm 4.6	26	58.4 \pm 33.4	23	6.2 \pm 2.0	26	127 \pm 31
		7.43–37.0		10.6–31.8		17.6–152		3.4–10.4		80–193
Muscle	32	1.80 \pm 0.82	32	12.0 \pm 2.2	32	2.86 \pm 0.80	31	9.7 \pm 3.4	32	80 \pm 24
		0.86–4.04		8.2–15.7		1.28–4.54		5.3–21.0		54–134
Feather	22	0.66 \pm 0.39	22	6.0 \pm 2.6	22	7.63 \pm 3.87	–	–	22	196 \pm 143
		0.18–1.53		2.8–11.8		2.30–14.2				100–797
Testis	3	42.7 \pm 14.4	3	19.2 \pm 13.7	–	–	–	–	3	145 \pm 15
		30.8–58.8		11.3–35.0						128–156

Sampling numbers are given accordingly

normality of distribution and homogeneity of variances using Shapiro–Wilk and Brown–Forsythe tests, respectively. Parametrical (Tukey’s HSD/ANOVA) and non-parametrical tests (Kruskal–Wallis/ANOVA) followed accordingly. Comparison between sexes was performed by means of Wilcoxon–Mann–Whitney test. Statistically significant results were set at $\alpha = 0.05$.

Results

Cadmium, Cu, Hg, Se and Zn concentrations in the tissues of *P. aequinoctialis* are presented in Table 1. No significant correlations (Pearson’s product-moment) were detected between tissue concentrations (both in wet weight and dry weight basis) and total body weight of the birds. The results overall showed that liver and kidney are the main sites for bioaccumulation of nonessential trace elements, with mean values of $58.4 \pm 33.4 \mu\text{g g}^{-1}$ dw of Hg in the liver and $65.7 \pm 22.2 \mu\text{g g}^{-1}$ dw of Cd in the kidney. In these tissues, Cd and Hg concentrations were one order of magnitude higher than those of muscle and feathers, which were the only tissues that did not present statistically significant difference for both contaminants. In contrast, Cd concentrations in the testis were as high as in the kidney.

Copper presented similar concentrations and ranges in liver, kidney, muscle and testis. Nevertheless, Cu variability was always lower than 30 % in the three former tissues, whereas it was above 70 % in the testis, mostly due to the small sampling number ($n = 3$) that precluded further statistical analyses. Feathers showed the lowest concentrations in average, but with a relatively high variability (coefficient of variation in 43 %). Se concentrations were the lowest in the liver and highest in the kidney, with average concentrations of 6.2 ± 2.0 and $16.1 \pm 4.8 \mu\text{g g}^{-1}$ dw, respectively. Nevertheless, the variation in Se concentrations remained always <35 %. For Zn, the concentrations were the lowest in muscles and highest in feathers in which the

variability was relatively high (coefficient of variation in 73 %). Nevertheless, this difference was statistically significant. Only muscle and liver, and liver and kidney (paired) did not present such differences. The other tissues, liver, kidney and testis, showed similar concentrations that were intermediate between those of muscle and feathers.

Males and females have shown significant contaminant differences in only one case: Cd in muscle was higher in males. All other comparisons were not statistically significant.

Inter- and intra-tissue correlations are shown in Table 2. Significant intra-tissue correlations appeared mostly within liver and kidney and to a lesser extent in muscle, whereas no correlation was found in feathers. In liver, Cd was positively correlated with Hg, Se and Zn, Cu was positively correlated with Zn, and Se to Hg. In kidney, Se was positively correlated with Cd, Cu and Zn, and Zn was positively correlated with Cd, Cu, and Se. In muscle, only Cu was negatively correlated with Zn and Se positively correlated with Hg.

Almost no significant inter-tissue correlations were detected for Cu and Zn, with only concentrations in the liver being negatively correlated with those in the feathers. Significant inter-tissue correlations appear mostly between the liver and kidney for nonessential elements but also for Se. Thus, Cd concentrations in the liver correlated positively with those in the kidney but negatively with those in the feathers. Concerning Hg, concentrations in the liver positively correlated with those in kidney and in muscle. Similar correlations were found for Se, as well as a significant correlation between liver and muscle Se concentrations.

In order to better understand detoxification processes, other correlations were also investigated for nonessential elements (Figs. 2, 3). Figure 2 presents the liver Hg concentrations as a function of kidney Cd concentrations. The relationship between both nonessential elements in their main tissues of bioaccumulation was, however, not

Table 2 Correlation (Pearson’s product-moment correlation coefficient) matrix for intra- and inter-tissue comparisons

	Cd	Cu	Hg	Se	Zn		Kidney	Liver	Muscle	Feather
Kidney						Cd				
Cd	1					Kidney	1			
Cu	-0.244	1				Liver	0.650*	1		
Hg	0.044	-0.070	1			Muscle	0.387	0.332	1	
Se	0.413*	0.399*	-0.051	1		Feather	-0.100	-0.635*	-0.188	1
Zn	0.382*	0.624*	0.063	0.552*	1	Cu				
Liver						Kidney				
Cd	1					Liver	0.320	1		
Cu	0.184	1				Muscle	-0.334	0.480	1	
Hg	0.435*	-0.244	1			Feather	0.029	-0.232	0.168	1
Se	0.607*	0.048	0.698*	1		Hg				
Zn	0.533*	0.525*	0.194	0.328	1	Kidney	1			
Muscle						Liver				
Cd	1					Muscle	0.584*	1		
Cu	0.236	1				Feather	0.419	0.579*	1	
Hg	-0.098	-0.050	1			Se				
Se	0.090	0.244	0.469*	1		Kidney	1			
Zn	-0.164	-0.663*	-0.107	-0.225	1	Liver	0.519*	1		
Feather						Muscle				
Cd	1					Feather	0.652*	0.728*	1	
Cu	0.420	1				Zn				
Hg	-0.286	-0.135	1			Kidney	1			
Se	-	-	-	1		Liver	0.296	1	1	
Zn	0.005	-0.038	0.290	-	1	Muscle	0.031	-0.259	0.352	1
						Feather	-0.316	-0.547*		

Significant values at $\alpha = 0.05$ are bold and asterisk marked




Fig. 2 Concentrations of liver Hg versus kidney Cd ($\mu\text{g g}^{-1}$ dw) in white-chinned petrels *P. aequinoctialis* from Kerguelen waters




Fig. 3 Se and Hg molar concentrations ($\mu\text{mol g}^{-1}$ dw) in liver and kidney of white-chinned petrels *P. aequinoctialis* from Kerguelen waters. **Bold line** the 1:1 molar ratio

significant. Figure 3 presents the molar relationship between Se and Hg. It highlighted a significant increase in the number of atoms of Se with those of Hg in the liver, but not in the kidney.

Discussion

Limited information is currently available for trace element concentrations in the tissues of the *P. aequinoctialis* with, to

the best of our knowledge, only three studies reported such levels for feathers (Blévin et al. 2013), for soft tissues (Anderson et al. 2010) and for both (Stewart et al. 1999). The concentrations of trace elements in liver and kidney of *P. aequinoctialis* from Kerguelen waters were overall in the same order of magnitude than those reported for *P. aequinoctialis* collected in 1995 and 1996 as bycatch of tuna fisheries around New Zealand (Stewart et al. 1999). Moreover, it presents the same distribution pattern between liver and kidney tissues for all elements (except Se which was not analysed by these authors). However, Hg concentrations in the feathers of adult birds from the present study were obviously higher (4.2 times) than those in chicks from the same location as a result of longer-term exposure in adults (Blévin et al. 2013).

In birds from Kerguelen waters, Cu concentrations in feathers were half those of *P. aequinoctialis* collected in Bird Island, South Georgia, during the 2001/02 austral summer (Anderson et al. 2010). In contrast, Zn values from the present study were 1.5 times higher when compared to Bird Island seabirds. Even though trace elements contamination is known to have several deleterious effects in birds testis (Scheuhammer 1987), data for analyses of such tissue is scarce. Kim et al. (1998) present data with similar distribution and, in a general way, within the same orders of magnitude as the present work. Elevated Cu variation in this tissue might be linked to body condition, as suggested by Malinga et al. (2010). These authors also suggested that the variation in Cu concentrations could be due to different stages of development of the gonads, which was visually confirmed in the present study. More generally, sex might play a role in trace element distribution in some seabird species (Kim et al. 1998). Sexual size dimorphism or different age composition in each gender (see Bugoni and Furness 2009) could affect contaminant levels, but the main driving sex-related factor is the excretion of potentially toxic materials into the egg that reduce the contaminant burdens of breeding females (Stewart et al. 1999). Combined to the different turnover rates of the different tissues, it is most likely that the significantly higher Cd muscle concentrations in males than in females of white-chinned petrels from the Kerguelen waters could be explained by the muscle turnover possibly matching the egg laying period, since it is long known that this tissue has a slower turnover rate than liver (Hobson and Clark 1992). Moreover, no dilution with growth or emaciation effect was detected, as previously stated. So, further discussion considers the whole grouped dataset, with no sex separation.

Influence of trophic ecology on trace element concentrations

Stewart et al. (1999) classified procellariiform seabirds in function of their pattern of accumulation of Hg in the liver

and Cd in the kidney. White-chinned petrels from Kerguelen fall into the same category as *Procellaria* petrels (including the white-chinned petrel) from New Zealand that are characterised by moderately high liver Hg and high kidney Cd concentrations. No significant relationship between liver Hg and kidney Cd was found, which can be attributed to the distinctive biochemistry of the two metals (Stewart et al. 1999). Elevated Hg concentrations in the tissues of *P. aequinoctialis* are likely due to the presence of mesopelagic prey in its diet that is mainly composed of fish and cephalopods in Kerguelen waters (66 and 17 % by mass, respectively; Delord et al. 2010). In marine predators, trophic position is not always the main factor in determining Hg concentrations, because mesopelagic prey was shown to increase disproportionately their Hg burdens (Thompson et al. 1998; Ochoa-acuña et al. 2002). Indeed, mesopelagic fish have elevated Hg concentrations as the result of enhanced Hg methylation in deep waters (Monteiro et al. 1996; Chauvelon et al. 2012). Cephalopods can also be considered as an important predator Hg source since they mainly contain Hg under organic forms, which is highly bioavailable for the upper trophic levels (Bustamante et al. 2006b). Long-term Hg exposure of *P. aequinoctialis* population from Kerguelen waters is also influenced by the species feeding habits outside the breeding period. For instance, *P. aequinoctialis* winters in the Benguela current, with associated dietary changes (Jackson 1988). In this area, fish is still the main prey class (73 % by mass), but it includes an important proportion (36 %) of offal, mainly fish heads and guts discarded by fishing vessels that keep aboard the commercially interesting part, i.e. the muscle. Muscle is one of the main tissues in fish with the lowest concentrations of trace elements (except Hg and arsenic; Bustamante et al. 2003; Kojadinovic et al. 2007; Metian et al. 2013). Because trace element concentrations in muscle are below concentrations in the whole fish, the consumption of offal could represent a higher intake of trace elements, especially Cd. In Kerguelen fish, liver contains up to 87 % of the whole Cd body burden with concentrations $>10 \mu\text{g g}^{-1} \text{dw}$ (Bustamante et al. 2003). Therefore, interactions with fisheries both during and outside the breeding season would lead to an increase intake of Cd in addition to Cd contamination resulting from squid consumption (Muirhead and Furness 1988; Bustamante et al. 1998a). Indeed, Cd levels in *P. aequinoctialis* tissues were in the same range as those of species with a higher proportion of cephalopods in their diets, such as albatrosses. For example, the Laysan albatross (*Phoebastria immutabilis*) presents average renal Cd levels of $59.7 \mu\text{g g}^{-1} \text{dw}$ (Honda et al. 1990) and Southern Ocean albatrosses (*Diomedea exulans*, *D. epomorpha*, *D. cauta*, *D. bulleri* and *D. melanophris*) show renal Cd levels ranging from 74 to $132 \mu\text{g g}^{-1} \text{dw}$ (Stewart et al. 1999).

Crustaceans are also likely to contribute to Cd exposure in *P. aequinoctalis*. For instance, the amphipods *Themisto* spp. (notably *T. gaudichaudii*) is naturally enriched in Cd (Ritterhoff and Zauke 1997; Bocher et al. 2003) and may represent a significant source of this element via both direct consumption (Catard et al. 2000; Delord et al. 2010) and secondary ingestion, since this crustacean contributes significantly to the diet of several petrel prey (Anderson et al. 2010). Assuming that offal have a Cd concentration of 10, “natural prey” fish of 0.1, cephalopods of 25, *Themisto* spp. of 40 and other crustaceans of $1 \mu\text{g g}^{-1}$ dw, an estimation of prey contribution in the Benguela waters suggests that cephalopods, offal, crustaceans and natural fish prey would be responsible for approximately 51.8, 43.5, 1.2 and 0.4 % of the total Cd intake, respectively. Therefore, the hypothesis of offals influencing greatly the birds’ Cd levels is likely to be verified. Such calculations were made from diet analysis by mass in the Benguela Region (Jackson 1988) together with available Cd concentrations of prey from the Southern Ocean (Ritterhoff and Zauke 1997; Bocher et al. 2003; Bustamante et al. 2003).

Diet is the main pathway of incorporation of nonessential but also essential elements in seabirds. Studies on trace elements in prey showed generally higher values for Cu and Zn in cephalopods when compared to mesopelagic fish (Bocher et al. 2003; Bustamante et al. 2003, 2006a; Anderson et al. 2010). Moreover, the antagonism between Cd, Cu and Zn could modify absorption, retention and distribution of the other two (Underwood 1977). Nevertheless, homeostatic control of internal essential elements plays a major role regulating their absorption in vertebrates, which depends on the individual nutritional requirements (Walsh 1990). In *P. aequinoctalis* from the present study, the correlations of Cu with Zn would reflect their nutritional status. Thus, low intra-specific variation in Cu and Zn concentrations should be expected in the tissues of marine birds in good health. In *P. aequinoctalis* from Kerguelen waters, Cu and Zn averages were close to the majority of other Kerguelen species (authors’ unpublished data), thus highlighting their homeostatic regulation, which is also confirmed by their comparatively low relative standard deviations.

Inter- and intra-tissues relationships between trace elements

Inter- and intra-tissue correlations between elements reflect the different processes of regulation of essential elements and of detoxification of nonessential elements.

In the case of intra-tissue analyses, in the liver, Cd positively correlated with Zn, Se and Hg, and Hg positively correlated with Se, which is consistent with previous data for procellariiformes seabirds such as the black-footed

albatross *Diomedea nigripes* from Ryukyu Islands (Kim et al. 1998). In regard to Cd and Zn co-accumulation, their correlation is likely to reflect Cd detoxification by metallothioneins (MTs) as already reported for different seabird groups (e.g. Ninomiya et al. 2004; Kojadinovic et al. 2007). Indeed, the great affinity of Cd for MTs leads to the competition with Zn for binding sites and to a reduction in the toxic effects of Cd (Klaassen et al. 1999), whereas the Zn displacement would induce MT synthesis, increasing the number of binding sites within the cells. In regard to the co-accumulation of Hg with Se, their correlation (the highest coefficient found in the whole study) is indicative of Hg detoxification mechanism involving the biomineralisation and storage of nontoxic Se–Hg complex and formation of liver granules when elimination through the feathers cannot match the individual’s dietary Hg intake (Nigro and Leonzio 1996). Such a process seems to be operative in *P. aequinoctalis* from Kerguelen waters in the liver but not in the kidney (Fig. 3).

Moreover, in the kidney, correlation of Se with Zn could be associated with their interaction in detoxifying processes, even though according to Norheim (1987) Se plays a more important role. Se correlation with Cu could be inferred from the consumption of some crustacean prey, notably *E. superba* and *T. gaudichaudii* (see Anderson et al. 2010), since the physico-chemical form in which Se is sequestered by different prey may vary and affect its subsequent bioavailability. Hence, it would appear that intra-specific variation in blood Se concentrations in seabirds most likely reflects individual differences in prey selection (see standard deviation in diet composition in Delord et al. 2010).

Besides Hg, Se is involved in detoxification of other heavy metals such as Ag, Cd, Cu and Zn (Ikemoto et al. 2004). In regard to inter-tissue analyses, Se significant positive correlations between liver, muscle and kidney are probably linked to the detoxifying mechanisms it is involved in, which is corroborated by its significant positive correlation with other elements, notably Hg in liver and muscle tissues. Hg significant positive correlation in liver with muscle tissue was already reported for several bird species (e.g. Borgå et al. 2006) and is assumed to result from the bioaccumulation of this element.

The significant positive correlations of Cd and Hg in both the liver and kidney confirm the role of these tissues as the main sites for detoxification and long-term storage (e.g. Bocher et al. 2003). The negative correlation between Cd concentrations in liver and Cd concentrations in feather could be related to a different diet during the nonbreeding period (higher offal proportion) when moulting normally occurs (Anderson et al. 2010) and in the breeding period, when samples were collected. Indeed, liver has a high protein turnover rate (Cherel et al. 1991) that reflects diet a few days before collection. The Zn negative correlation

between liver and feather could result from the same processes, which is confirmed by its significant correlation with Cd in the liver. This shift towards liver co-accumulation can happen in periods as short as one or 2 weeks, as demonstrated by Levengood and Skowron (2007). In other words, Cd and Zn are co-accumulated in the liver (emphasised by the significant intra-tissue correlation) and before being released from the liver to the feathers (emphasised by the negative inter-tissue correlation between liver and feathers). Even though, Zn in feathers presented large variation. This could be due to either molecular interactions between Cd, Cu, and Zn or to Zn affinity for liver MTs (Bocher et al. 2003).

Finally, in regard to the use of feathers as a non-destructive monitoring tool, it is noticeable that they presented significant inter-tissue correlation in only two occasions: Zn and Cd, both with liver and both negative as well. No significant correlation with other trace elements was found for none of the tissues, possibly due to two main factors: firstly, the sampling of the birds occurred in different times of the year and not only that, but consequently in different time shifts in regard to the moult. This might result in a degree of fluctuation in the trace element concentration of the internal tissues, and therefore, the static concentration of the feathers may no longer be correlated with those. The second hypothesised factor is that moult occurs normally during the nonbreeding period (Anderson et al. 2010), when diet is different (Jackson 1988); therefore, the variation in internal tissues could be enhanced due to a variation in exposure via diet. A further study with birds sampled in a narrower time interval and with a homogenous set of animals in regard to the last moult could better answer the question whether feathers can be used to monitor trace elements in this species and under what conditions.

Acknowledgments The authors thank the fieldworkers who helped with collecting the petrels. The present work was supported financially and logistically by the Agence Nationale de la Recherche (programme POLARTOP, O. Chastel), the Institut Polaire Français Paul Emile Victor (IPEV, programme no. 109, H. Weimerskirch) and the Terres Australes et Antarctiques Françaises (TAAF). C.V.Z. Cipro was supported financially through a post-doctoral grant from the University of La Rochelle and the CNRS (French acronym for National Council of Scientific Research).

References

- Abdulla M, Chmielnicka J (1990) New aspects on the distribution and metabolism of essential trace elements after dietary exposure to toxic metals. *Biol Trace Elem Res* 23:25–53
- Agusa T, Matsumoto T, Ikemoto T, Anan Y, Kubota R, Yasunaga G, Kunito T, Tanabe S, Ogi H, Shibata Y (2005) Body distribution of trace elements in black-tailed gulls from Rishiri Island, Japan: age-dependent accumulation and transfer to feather and eggs. *Environ Toxicol Chem* 24:2107–2120
- Anderson ORJ, Phillips RA, Shore RF, McGill RAR, McDonald RA, Bearhop S (2010) Element patterns in albatrosses and petrels: influence of trophic position, foraging range, and prey type. *Environ Pollut* 158:98–107
- Barbraud C, Marteau C, Ridoux V, Delord K, Weimerskirch H (2008) Demographic response of a population of white-chinned petrels *Procellaria aequinoctialis* to climate and longline fishery bycatch. *J Appl Ecol* 45:1460–1467
- Barbraud C, Delord K, Marteau C, Weimerskirch H (2009) Estimates of population size of white-chinned petrels and grey petrels at Kerguelen Islands and sensitivity to fisheries. *Anim Conserv* 12:258–265
- Berrow SD, Croxall JP, Grant SD (2000) Status of white-chinned petrels *Procellaria aequinoctialis* Linnaeus 1758, at Bird Island, South Georgia. *Antarct Sci* 12:399–405
- Birdlife International (2013) White-chinned Petrel *Procellaria aequinoctialis*. <http://www.birdlife.org/datazone/speciesfactsheet.php?id=3922>. Accessed 2 July 2013
- Blévin P, Carravieri A, Jaeger A, Chastel O, Bustamante P, Cherel Y (2013) Wide range of mercury contamination in chicks of Southern Ocean seabirds. *PLoS ONE* 8:e54508
- Bocher P, Caurant F, Miramand P, Cherel Y, Bustamante P (2003) Influence of the diet on the bioaccumulation of heavy metals in zooplankton-eating petrels at Kerguelen archipelago, Southern Indian Ocean. *Polar Biol* 26:759–767
- Borgå K, Campbell L, Gabrielsen GW, Norstrom RJ, Muir DCG, Fisk AT (2006) Regional and species specific bioaccumulation of major and trace elements in Arctic seabirds. *Environ Toxicol Chem* 25:2927–2936
- Bugoni L, Furness R (2009) Age composition and sexual size dimorphism of albatrosses and petrels off Brazil. *Mar Ornithol* 260:253–260
- Bustamante P, Caurant F, Fowler SW, Miramand P (1998a) Cephalopods as a vector for the transfer of cadmium to top marine predators in the north-east Atlantic Ocean. *Sci Total Environ* 220:71–80
- Bustamante P, Cherel Y, Caurant F, Miramand P (1998b) Cadmium, copper and zinc in octopuses from Kerguelen Islands, Southern Indian Ocean. *Polar Biol* 19:264–271
- Bustamante P, Bocher P, Cherel Y, Miramand P, Caurant F (2003) Distribution of trace elements in the tissues of benthic and pelagic fish from the Kerguelen Islands. *Sci Total Environ* 313:25–39
- Bustamante P, Bertrand M, Boucaud-Camou E, Miramand P (2006a) Subcellular distribution of Ag, Cd Co, Cu, Fe, Mn, Pb, and Zn in the digestive gland of the common cuttlefish *Sepia officinalis*. *J Shellfish Res* 25:987–993
- Bustamante P, Lahaye V, Durnez C, Churlaud C, Caurant F (2006b) Total and organic Hg concentrations in cephalopods from the North Eastern Atlantic waters: influence of geographical origin and feeding ecology. *Sci Total Environ* 368:585–596
- Carravieri A, Bustamante P, Churlaud C, Cherel Y (2013) Penguins as bioindicators of mercury contamination in the Southern Ocean: birds from the Kerguelen Islands as a case study. *Sci Total Environ* 454(455):141–148
- Catard A, Weimerskirch H, Cherel Y (2000) Exploitation of distant Antarctic waters and close shelf-break waters by white-chinned petrels rearing chicks. *Mar Ecol Prog Ser* 194:249–261
- Cherel Y, Attaix D, Rosolowska-Huszcz D, Belkhou R, Robin JR, Arnal M, Le Maho Y (1991) Whole-body and tissue protein synthesis during brief and prolonged fasting in the rat. *Clin Sci* 81:611–619
- Chouvelon T, Spitz J, Caurant F, Mèndez-Fernandez P, Autier J, Lassus-Débat A, Chappuis A, Bustamante P (2012) Enhanced bioaccumulation of mercury in deep-sea fauna from the Bay of Biscay (north-east Atlantic) in relation to trophic positions identified by analysis of carbon and nitrogen stable isotopes. *Deep-Sea Res Part I* 65:113–124

- Connan M, Cherel Y, Mabile G, Mayzaud P (2007) Trophic relationships of white-chinned petrels from Crozet Islands: combined stomach oil and conventional dietary analyses. *Mar Biol* 152:95–107
- Coquery M, Azemard S, Mora S (2001) The analytical performance study for the Medpol region: determination of trace elements and methylmercury in estuarine sediment sample IAEA-405
- Delord K, Cotté C, Péron C, Marteau C, Pruvost P, Gasco N, Duhamel G, Cherel Y, Weimerskirch H (2010) At-sea distribution and diet of an endangered top predator: relationship between white-chinned petrels and commercial longline fisheries. *Endanger Species Res* 13:1–16
- Dietz R, Noorgard J, Hansen JC (1998) Have arctic marine mammals adapted to high cadmium levels? *Mar Pollut Bull* 36:490–492
- Elliott J, Scheuhammer A (1997) Heavy metal and metallothionein concentrations in seabirds from the Pacific coast of Canada. *Mar Pollut Bull* 34:794–801
- Fitzgerald W, Engstrom D (1998) The case for atmospheric mercury contamination in remote areas. *Environ Sci Technol* 32:1–7
- Furness R, Camphuysen K (1997) Seabirds as monitors of the marine environment. *ICES J Mar Sci* 54:726–737
- Gilbertson M, Elliott JE, Peakall DB (1987) Seabirds as indicators of marine pollution. In: Diamond AW, Filion F (eds) *The value of birds*. International Council for Bird Preservation, Cambridge, pp 231–248
- Goutte A, Bustamante P, Barbraud C, Delord K, Weimerskirch H, Chastel O (2014) Demographic responses to mercury exposure in two closely-related Antarctic top predators. *ESA Ecol*. doi:10.1890/13-1229.1
- Hobson KA, Clark RG (1992) Assessing avian diets using stable isotopes I: turnover of ^{13}C in tissues. *Condor* 94:181–188
- Honda K, Marcovecchio JE, Kan S, Tatsukawa R, Ogi H (1990) Metal concentrations in pelagic seabirds from the North Pacific Ocean. *Arch Environ Contam Toxicol* 19:704–711
- Ikemoto T, Kunito T, Tanaka H, Baba N, Miyazaki N, Tanabe S (2004) Detoxification mechanism of heavy metals in marine mammals and seabirds: interaction of selenium with mercury, silver, copper, zinc, and cadmium in liver. *Arch Environ Contam Toxicol* 47:402–413
- Jackson S (1988) Diets of the white-chinned petrel and sooty shearwater in the southern Benguela region, South Africa. *Condor* 90:20–28
- Kim E, Goto R, Tanabe S (1998) Distribution of 14 elements in tissues and organs of oceanic seabirds. *Arch Environ Contam Toxicol* 35:638–645
- Klaassen CD, Liu J, Choudhuri S (1999) Metallothionein: an intracellular protein to protect against cadmium toxicity. *Annu Rev Pharmacol Toxicol* 39:267–294
- Kojadinovic J, Le Corre M, Cosson RP, Bustamante P (2007) Trace elements in three marine birds breeding on Reunion Island (Western Indian ocean): part I-factors influencing their bioaccumulation. *Arch Environ Contam Toxicol* 52:418–430
- Levengood JM, Skowron LM (2007) Coaccumulation of cadmium and zinc in tissues of sentinel mallards (*Anas platyrhynchos*) using a former dredge-disposal impoundment. *Arch Environ Contam Toxicol* 53:281–286
- Mailman RB (1980) Heavy metals. In: Perry JJ (ed) *Introduction to environmental toxicology*. Elsevier, New York, pp 34–43
- Malinga M, Szefer P, Gabrielsen GW (2010) Age, sex and spatial dependent variations in heavy metals levels in the Glaucous Gulls (*Larus hyperboreus*) from the Bjørnøya and Jan Mayen, Arctic. *Environ Monit Assess* 169:407–416
- Metian M, Warnau M, Chouvelon T, Pedraza F, Rodriguez Y, Baena AM, Bustamante P (2013) Trace element bioaccumulation in reef fish from New Caledonia: influence of trophic groups and risk assessment for consumers. *Mar Environ Res* 87(88):26–36
- Monteiro LR, Furness RW (1995) Seabirds as monitors of mercury in the marine environment. *Water Air Soil Pollut* 80:851–870
- Monteiro LR, Costa V, Furness RW, Santos RS (1996) Mercury concentrations in prey fish indicate enhanced bioaccumulation in mesopelagic environments. *Mar Ecol Prog Ser* 141:21–25
- Muirhead S, Furness RW (1988) Heavy metal concentrations in the tissues of seabirds from Gough Island, South Atlantic Ocean. *Mar Pollut Bull* 19:278–283
- Nigro M, Leonzio C (1996) Intracellular storage of mercury and selenium in different marine vertebrates. *Mar Ecol Prog Ser* 135:137–143
- Ninomiya R, Koizumi N, Murata K (2004) Metal concentrations in the liver and kidney of aquatic mammals and penguins. *Biol Trace Elem Res* 97:135–148
- Norheim G (1987) Levels and interactions of heavy metals in sea birds from Svalbard and the Antarctic. *Environ Pollut* 47:83–94
- Ochoa-acuña H, Sepúlveda MS, Gross TS (2002) Mercury in feathers from Chilean birds: influence of location, feeding strategy, and taxonomic affiliation. *Mar Pollut Bull* 44:340–345
- Péron C, Delord K, Phillips RA, Charbonnier Y, Marteau C, Louzao M, Weimerskirch H (2010) Seasonal variation in oceanographic habitat and behaviour of white-chinned petrels *Procellaria aequinoctialis* from Kerguelen Island. *Mar Ecol Prog Ser* 416:267–284
- Ridoux V (1994) The diets and dietary segregation of seabirds at the subantarctic Crozet Islands (part 1). *Mar Ornithol* 22:1–64
- Ritterhoff J, Zauke GP (1997) Influence of body length, life-history status and sex on trace metal concentrations in selected zooplankton collectives from the Greenland Sea. *Mar Pollut Bull* 34:614–621
- Scheuhammer AM (1987) The chronic toxicity of aluminium, cadmium, mercury, and lead in birds: a review. *Environ poll* 46:263–295
- Scheuhammer AM, Meyer MW, Sandheinrich MB, Murray MW (2007) Effects of environmental methylmercury on the health of wild birds, mammals, and fish. *Ambio* 36:12–18
- Selin NE (2009) Global biogeochemical cycling of mercury: a review. *Annu Rev Environ Resour* 34:43–63
- Stewart F, Phillips R, Bartle J (1999) Influence of phylogeny, diet, moult schedule and sex on heavy metal concentrations in New Zealand Procellariiformes. *Mar Ecol Prog Ser* 178:295–305
- Tan SW, Meiller JC, Mahaffey KR (2009) The endocrine effects of mercury in humans and wildlife. *Crit Rev Toxicol* 39:228–269
- Tartu S, Goutte A, Bustamante P, Angelier F, Moe B, Clément-Chastel C, Bech C, Gabrielsen GW, Bustness JO, Chastel O (2013) To breed or not to breed: endocrine response to mercury contamination by an arctic seabird. *Biol Lett* 9:20130317
- Thompson DR, Furness RW, Monteiro LR (1998) Seabirds as biomonitors of mercury inputs to epipelagic and mesopelagic marine food chains. *Sci Total Environ* 213:299–305
- Underwood EJ (1977) *Trace elements in human and animal nutrition*. Academic ed, New York
- Walsh PM (1990) The use of seabirds as monitors of heavy metals in the marine environment. In: Furness RW, Rainbow PS (eds) *heavy metals in the marine environment*. CRC, New York, pp 183–204