

Delineation of ^{134}Cs uptake pathways (seawater and food) in the variegated scallop *Mimachlamys varia*



Simon Pouil ^{a, b}, Paco Bustamante ^b, Michel Warnau ^a, François Oberhänsli ^a,
Jean-Louis Teysié ^a, Marc Metian ^{a, *}

^a International Atomic Energy Agency, Environment Laboratories (IAEA-EL), 4a Quai Antoine 1er, MC-98000 Principality of Monaco, Monaco

^b Littoral Environnement et Sociétés (LIENSs), UMR 7266, CNRS-Université de La Rochelle, 2 rue Olympe de Gouges, F-17000 La Rochelle, France

ARTICLE INFO

Article history:

Received 25 March 2015
Received in revised form
1 June 2015
Accepted 19 June 2015
Available online xxx

Keywords:

Bivalve
Scallop
Cesium
Kinetics
Depuration

ABSTRACT

Among bivalves, scallops have been shown to be good bioindicator species for radionuclide monitoring. The present paper looked at the Cs bioaccumulation capacities of the variegated scallop *Mimachlamys varia* exposed separately via seawater and food under laboratory conditions. Results were compared with data previously obtained for the king scallop *Pecten maximus*, the only Pectinid species for which Cs accumulation has been studied in laboratory. Results indicated that *M. varia* has higher uptake capacity (CF: 1.86 ± 0.08) but lower absorption efficiency (A_{01} : $33 \pm 5\%$) than *P. maximus* when exposed to waterborne Cs (CF of *P. maximus*: 0.94 ± 0.05 and A_{01} : $45 \pm 3\%$). When scallops were fed radiolabeled phytoplankton, the assimilation efficiency of Cs was similar for the two species (AE: $24 \pm 3\%$ for *M. varia* and $28 \pm 4\%$ for *P. maximus*). Interspecific differences in terms of accumulation and retention, can be explained by physiological factors (including size of individuals) and/or difference in storage mechanisms. Indeed, organotropism differed between the two scallop species, suggesting the occurrence of specific redistribution mechanisms towards the tissues involved in Cs storage, excretion and detoxification. Finally, the present study examined the relative contribution of the different exposure pathways (seawater and food) to global ^{134}Cs bioaccumulation for *M. varia*. Results showed that food constitutes the main accumulation pathway, contributing for 77% of the global ^{134}Cs bioaccumulation.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Tracking radionuclide contamination using organisms (i.e. bio-monitoring) helps understanding the levels as well as the variation (in time and space) of radionuclides that are bioavailable in the environment (Philips and Rainbow, 1993). Since 1950, marine ecosystems were sporadically subjected to radionuclide contamination from industries, nuclear accidents and fallout from nuclear weapon testing and uses. Recently, the accident that occurred in the Daiichi Nuclear Power Plant at Fukushima in Japan, released massive amount of radioactive Cs in the marine environment (Chino et al., 2011; Bailly du Bois et al., 2012).

Among marine organisms, Pectinids have been proposed as good bioindicator candidates of radionuclide contamination (Marzano et al., 2000; Bustamante et al., 2002). This has been

confirmed experimentally for the king scallop *Pecten maximus* (Metian et al., 2011). Field surveys (e.g. Marzano et al., 2000) have also documented the capacity of the family of Pectinids to concentrate radionuclides, in particular Cs. However, levels of radionuclides in organisms are influenced by various factors such as geographical origin, season, sexual maturity or species considered (Bryan, 1963; Mauri et al., 1990; Bustamante and Miramand, 2004, 2005). Laboratory characterization of Cs bioaccumulation by Pectinids is currently limited to one species, *P. maximus* (Metian et al., 2011) and since phylogenetically-close species may have different accumulation capacities (as shown for metals in scallops, e.g. Metian et al., 2007 and in oysters, Hédouin et al., 2010a), it is worth comparing Cs bioaccumulation capacities among scallops species.

Therefore, the objective of the present study was to investigate ^{134}Cs bioaccumulation capacities of the variegated scallop *Mimachlamys varia*, exposed via the dissolved and trophic pathways. This species is an interesting model to assess Cs contamination in coastal area given the fact that it settles, with its byssus, on rocky substrates (Shumway and Parsons, 2006) whereas *P. maximus* lives

* Corresponding author. Radioecology Laboratory, IAEA Environment Laboratories, 4a Quai Antoine 1er, MC-98000 Principality of Monaco, Monaco.

E-mail address: m.metian@iaea.org (M. Metian).

buried in the sediment. The same protocol as described in Metian et al. (2011) was used in order to allow easy comparison between the two scallop species. In addition, the relative contribution of the different exposure pathways (dissolved vs. dietary) of Cs bioaccumulation in *M. varia* was determined.

2. Materials and methods

2.1. Origin and acclimation of organisms

In spring 2004 and 2005, one hundred variegated scallops *M. varia* were collected on the French Atlantic coast (Pertuis Breton, Charente-Maritime) by SCUBA diving. They were transported to IAEA-EL premises in Monaco and were acclimated to laboratory conditions for 4 weeks (constantly aerated, open-circuit 700 L aquarium; flux: 50 L h⁻¹; salinity: 36 p.s.u.; temperature: 17 ± 0.5 °C; pH: 8.0 ± 0.1; light/dark cycle: 12 h/12 h) prior to experimentation. During this period, scallops were fed daily an algal diet (*Isochrysis galbana*).

2.2. Radiotracer and counting

Uptake and depuration kinetics of ¹³⁴Cs in scallops were determined using high-specific activity ¹³⁴Cs purchased from Isotope Product Lab (¹³⁴CsCl 0.1 N, T_{1/2} = 2 years). ¹³⁴Cs was counted using a high-resolution γ -spectrometer system composed of four Germanium (N- or P-type) detectors (EGNC 33–195-R, Canberra® and Eurysis®) connected to a multi-channel analyzer and a computer equipped with a spectra analysis software (Interwinner® 6). The radioactivity was determined by comparison with standards of known activity and of appropriate geometry. Measurements were corrected for counting efficiency and physical radioactive decay. The counting time was adjusted to obtain a propagated counting error less than 5% (Rodriguez y Baena et al., 2006).

2.3. Seawater exposure

Twenty three *M. varia* (30 ± 7 g) were placed in a 70-L aquarium (constantly aerated, closed-circuit aquarium; salinity: 36 p.s.u.; temperature: 17 °C ± 0.5 °C; pH: 8.0 ± 0.1; light/dark cycle: 12 h/12 h) and exposed for 7 d to ¹³⁴Cs dissolved in 0.45 μ m filtered seawater (1.4 kBq L⁻¹). Spiked seawater was renewed regularly in order to keep radioactivity constant. The ¹³⁴Cs in seawater was checked before and after each spike renewal. Nine tag-identified scallops were collected at different time intervals and were whole-body radioanalyzed alive. At the end of the 7 d exposure period, four scallops were collected and dissected into their body compartments (digestive gland, gills, kidneys, intestine, gonad, foot, mantle, adductor muscle and remaining tissues). Samples were radioanalyzed in order to assess the radionuclide body distribution. The remaining scallops were then placed in uncontaminated water in the same conditions for 36 d and the nine identified individuals were regularly radioanalyzed alive in order to follow the depuration kinetics of ¹³⁴Cs. Three non-exposed individuals were introduced into the aquarium in order to control possible tracer recycling from the contaminated scallops. During the 36 d depuration period, scallops were fed daily with *I. galbana* (5.10⁴ cells mL⁻¹). At the end of the depuration period, four contaminated scallops were collected and dissected.

2.4. Dietary exposure

¹³⁴Cs transfer to scallops through their diet was studied using the Haptophyceae *I. galbana*. Phytoplankton cells were exposed to 7 kBq ¹³⁴Cs L⁻¹ during their exponential growing phase (7 d). After

that period, the phytoplankton medium was filtered (1 μ m-mesh size; Osmonic® filters), and the phytoplankton cells re-suspended in a 70 L aquarium in the same conditions as previously described for the scallops. Cell density was 5.10⁴ cells mL⁻¹ to avoid pseudofeces production by the molluscs. Six *M. varia* (17 ± 5 g) had been placed in the aquarium for one week before the start of feeding experiment. Scallops were fed with radiolabelled phytoplankton for 2 h in closed-circuit. After the feeding period, all scallops were γ -counted and flow restored in the aquarium. Depuration was followed during 30 d as described previously (see section 2.3). Three non-exposed individuals were used as controls of tracer recycling. During the 30-d depuration period, scallops were fed daily with *I. galbana* (5.10⁴ cells mL⁻¹). Four contaminated individuals were collected after 30 d and dissected to determine the radionuclide body distribution.

2.5. Data analysis

Whole-body uptake and depuration kinetics were fitted using nonlinear regression routines and iterative adjustment (Statistica® 7) and statistical methods described by Warnau et al. (1996a, 1996b) and Metian et al. (2011). The level of significance for statistical analyses was always set at $\alpha = 0.05$. The relative contribution of each pathway (seawater and food) was determined using a global bioaccumulation model described in Hédouin et al. (2010a). In addition to the parameters obtained from each biokinetic models, we have used the ingestion rate of *M. varia* (g g⁻¹ d⁻¹, wet wt basis) and the K_d, partition coefficients for phytoplankton calculated experimentally (Metian et al., 2009).

3. Results

3.1. Seawater exposure

Whole-body uptake of ¹³⁴Cs in *M. varia* followed exponential kinetics, reaching a steady-state equilibrium after approximately 4 days (R² = 0.79; Fig. 1A and Table 1). The kinetic parameters and their statistics are shown in Table 1. The whole-body concentration factor of ¹³⁴Cs (i.e. ratio between the radioisotope activity in organisms – Bq g⁻¹ wet wt – and time-integrated activity in the seawater – Bq g⁻¹) estimated at steady state (CF_{ss}) was 1.86 ± 0.08 (Fig. 1A and Table 1).

Calculated CF_{7d} for the different body compartments are shown in Table 2. Among the tissues and organs, the foot displayed the highest CF closely followed by the kidney and the digestive gland (Table 2). In term of distribution, Cs was mainly located in the adductor muscle, the digestive gland and the mantle with, respectively, 26 ± 3, 24 ± 10 and 23 ± 5% of the total Cs body burden.

After the exposure period, non-contaminating conditions were restored and depuration of ¹³⁴Cs was followed for 36 d. The whole-body depuration kinetics of ¹³⁴Cs was best described by a two-component exponential model (Fig. 1B and Table 1). An important part of ¹³⁴Cs was efficiently absorbed (A₀₁: 33 ± 5%) with an associated long-term biological half-life (T_{b1/21}) of 178 d (not significantly different from infinite; Table 1). After 36 d of the depuration, ¹³⁴Cs was mainly located in the soft body parts (72 ± 7%; Table 2). Distribution of ¹³⁴Cs among the soft tissues displayed a different pattern than that observed at the end of the exposure period (Table 2). Indeed, the digestive gland contained most of ¹³⁴Cs (59 ± 13%) and the relative proportion of the total Cs found in the intestine was much higher than the one in the same organ after the exposure phase (10 vs. <1%).




Fig. 1. (A) Uptake kinetics of ^{134}Cs in *M. varia* exposed for 7 d to dissolved radiotracers ($n = 9$); (B) subsequent 36-d depuration kinetics ($n = 9$) and (C) 30-d depuration kinetics after ^{134}Cs -labeled food ingestion ($n = 6$). All values are mean \pm SD.

Table 1
Parameters of whole-body uptake and depuration kinetics of ^{134}Cs in *M. varia* 1) exposed for 7 d to waterborne radionuclide ($n = 9$) and followed by 36 d of depuration ($n = 9$) and 2) after a 2-h feeding on radiolabelled *I. galbana* followed by 30 d of depuration ($n = 6$).

Experiment	Uptake			Depuration				
	$\text{CF}_{\text{ss}} \pm \text{ASE}$	$k_u \pm \text{ASE}$	R^2	$A_{0s} \pm \text{ASE}$	$T_{b/2s} \pm \text{ASE}$	$A_{0l} \pm \text{ASE}$	$T_{b/2l}$	R^2
1) Seawater	1.86 ± 0.08^a	1.77 ± 0.33^a	0.79	66.41 ± 6.92^a	0.6 ± 0.2	33.09 ± 4.63^a	178	0.50
2) Feeding	–	–	–	75.46 ± 5.48^a	0.3 ± 0.1	24.45 ± 2.92^a	115	0.80

Uptake parameters: CF_{ss} : concentration factors at steady state; k_u : uptake rate constant (d^{-1}). Depuration parameters: A_{0s} and A_{0l} : remaining activity (%) according to the short- and the long-lived exponential component, respectively; $T_{b/2s}$: biological half-life (d). ASE: asymptotic standard error; R^2 : determination coefficient of the uptake or depuration kinetics.

^a Probability of the model adjustment: $p < 0.0001$.

Table 2
Concentration Factors (mean CF \pm SD) and body distribution (mean % \pm SD) of ^{134}Cs in *M. varia* during seawater (after 7 d of exposure and after 36 d of depuration) and feeding (after 30 d of depuration) experiments.

Compartments	Seawater experiment		Food experiment ^a	
	Uptake (7 d, $n = 4$)		Loss (30d, $n = 4$)	
	Concentration factor	Distribution (%)	Distribution (%)	Distribution (%)
Digestive gland	6 ± 4	24 ± 10	59 ± 13	57 ± 16
Gills	4 ± 0.4	15 ± 4	2 ± 1	7 ± 6
Kidneys	6 ± 1	4 ± 2	3 ± 1	6 ± 5
Intestine	2 ± 0.3	<1	10 ± 3	2 ± 1
Gonad	3 ± 1	4 ± 1	4 ± 1	4 ± 2
Foot	7 ± 2	3 ± 1	3 ± 4	7 ± 6
Mantle	3 ± 0.4	23 ± 5	2 ± 1	8 ± 7
Adductor muscle	3 ± 1	26 ± 3	14 ± 9	9 ± 9
Remaining tissues	4 ± 1	3 ± 1	3 ± 1	2 ± 1
Whole soft parts	3 ± 1	41 ± 9	72 ± 7	n.a.
Shell	2 ± 1	59 ± 9	28 ± 7	n.a.

^a For this experiment, counting errors are comprises between 5 and 15% due to the low activities measured.

3.2. Dietary exposure

The depuration kinetics of the ^{134}Cs ingested with food in *M. varia* was best fitted by a double exponential model (Fig. 1C and Table 1). *M. varia* displayed an assimilation efficiency of $24 \pm 3\%$. The depuration rate constant, k_{el} ($0.006 \pm 0.006 \text{ d}^{-1}$) was not significantly different from 0 ($p > 0.05$), and therefore the derived $T_{b/2}$ is not significantly different from infinite (115 d). At the end of the depuration period, the digestive gland contained the major part of ^{134}Cs ($57 \pm 16\%$, Table 2).

3.3. Bioaccumulation model

The relative contribution of each exposure pathway to the global bioaccumulation of ^{134}Cs was determined using the different kinetic parameters determined in the two experiments, as well as other parameters such as the ^{134}Cs K_{df} in phytoplankton ($1.32 \cdot 10^2$ for *I. galbana*; present study) and the ingestion rate of phytoplankton in scallops ($IR = 0.0869 \text{ g g}^{-1} \text{ d}^{-1}$; Metian et al., 2009). Results of the computations indicated that feeding pathway was the major contributor (77%) to the global bioaccumulation of ^{134}Cs in *M. varia* whereas seawater contributed for 23%.

4. Discussion

Pectinids have been used in biomonitoring programmes to assess radionuclide contamination in the marine environment (JCAC, 2002) given their high accumulation capacity shown during field surveys (Marzano et al., 2000; Bustamante et al., 2002). However, the behavior of radionuclides in scallops and their mode of uptake and depuration are less known in comparison to other bivalve families such as Ostreidae or Mytilidae (Cranmore and Harrison, 1975; Ke et al., 2000; Ryan, 2002). To the best of our knowledge, experimental investigations of ^{134}Cs bioaccumulation have been limited to one species *P. maximus* exposed to three pathways. Unlike *P. maximus*, which lives buried in the bottom sediment, *M. varia* settles, with its byssus, on rocky substrates (Shumway and Parsons, 2006); therefore, the exposure of *M. varia* to radionuclides occurs through seawater and food pathways.

Our results show that there are interspecific differences in Pectinids in terms of accumulation and retention of dissolved Cs. Indeed, the concentration factor (CF_{ss}) in *M. varia* was two times higher than CF_{ss} of *P. maximus* (Metian et al., 2011). Difference can be attributed to variation of physiological factors (including size of individuals) and/or difference in storage mechanisms (as highlighted by the contrasting body distribution patterns of Cs in both species at the end of exposure; Metian et al., 2011). Similar interspecific differences of CF_{ss} between these two species have already been highlighted for other radionuclides (^{110m}Ag , ^{109}Cd , ^{54}Mn , ^{210}Pb , ^{65}Zn ; Metian et al., 2007, 2009a, 2008, 2009b, 2009c). Interspecific differences also concerned Cs depuration. Indeed, the percentage of long-term retention of ^{134}Cs absorbed by *M. varia* after 36 d of depuration (following a ^{134}Cs exposure through seawater) did not exceed $33 \pm 5\%$ (Table 1) whereas $45 \pm 3\%$ of ^{134}Cs was retained by *P. maximus* (Metian et al., 2011). Differences observed in term of retention were radionuclide-dependent: A_{01} were higher in *P. maximus* for ^{110m}Ag , ^{57}Co , ^{109}Cd and ^{65}Zn than in *M. varia* whereas no differences were observed between the two species for ^{54}Mn and ^{210}Pb (Metian et al., 2007, 2009a, 2008, 2009b, 2009c). For Cs, difference in CF between the two species exposed to the same contamination conditions is related to a higher Cs uptake rate (uptake rate constant: 1.77 d^{-1} vs. 0.55 d^{-1}) in *M. varia* compared to *P. maximus* (Metian et al., 2011, Table 1).

Interestingly, interspecific differences can be identified upon a dissolved exposure but not when Pectinids are exposed to ^{134}Cs

through food. The present results show that the retention of Cs from food is similar for both scallop's species. Indeed, when *M. varia* were fed with radiolabeled phytoplankton cells (*I. galbana*), ^{134}Cs was similarly assimilated (AE: $24 \pm 3\%$, Table 1) compared to AE of $28 \pm 4\%$ obtained for *P. maximus* fed in the same conditions (Metian et al., 2011). This similarity in Cs AE between these two Pectinids was not observed for other elements: the AEs of ^{110m}Ag , ^{57}Co , and ^{210}Pb were higher in *M. varia* whereas ^{54}Mn AE was higher for the *P. maximus* (Metian et al., 2007, 2008, 2009a, 2009b, 2009c). To confirm our results for ^{134}Cs , it would be appropriate to test the AE between these two scallop species with other radiolabelled phytoplankton species since Hédouin et al. (2010b) have already demonstrated that the quality of food may affect radionuclide accumulation (^{57}Co , ^{54}Mn and ^{65}Zn) in bivalves.

Globally, our results confirm that Cs is well retained in Pectinids. Fig. 2 put these results in perspectives with previous studies on Cs accumulation in bivalves. Whereas ^{134}Cs CF in Pectinids is similar (or lower in some cases) than other bivalve species, retention is always higher in Pectinids (Fig. 2). One has however to keep in mind that some of the observed differences in CF can be partly due to others factor than interspecific bioaccumulation properties, such as difference in ambient salinity. Indeed it is well documented that CF for a given species generally increases with decreasing salinity (e.g. Ke et al., 2000). However, most of the species compared here (see Fig. 2) were living in a narrow range of salinity (between 35 and 38 p.s.u.; Nolan and Dahlgard, 1991; Bustamante et al., 2006; Qureshi et al., 2007; Lacoue-Labarthe et al., 2010; Metian et al., 2011).

Percentage of Cs retention from seawater (A_{01}) and food (AE) is higher in Pectinids than other bivalves. For example, Mytilidae showed low AEs for Cs with values (ranging between 3 and 13% for *Perna viridis*; Table 2, e.g. Wang et al., 2000). However, Fig. 2 also shows the reported AEs of Cs in other molluscs (predators) and they can be higher than Pectinids (ranging between 19 and 55% for gastropods and cephalopods; e.g. Wang et al., 2000; Bustamante et al., 2006). These results suggest that food source is a key pathway for the transfer of Cs towards molluscs.

In the natural environment, *M. varia* may be exposed to radioactive Cs through seawater and food and each pathway contribute simultaneously to the global bioaccumulation of ^{134}Cs in the organism. Using the kinetic parameters that we obtained experimentally and K_{df} and IR that we measured, the relative contributions of the two exposure pathways were delineated, using the model developed by Landrum et al. (1992) and adapted more recently (Hédouin et al., 2010a). Results indicate that food has a major role in Cs accumulation in *M. varia* in the field (relative contribution of 77% in average). The importance of the trophic pathway in the Cs accumulation has been suggested in taxa such as cephalopods (Bustamante et al., 2006) and fish (Pentreath and Jefferies, 1971; Hewett and Jefferies, 1978) although without direct evidences. In Pectinids, food has been shown to be the major contamination pathway for ^{109}Cd (*M. varia* and *P. maximus* respectively $>99\%$ and 84% of the total contribution; Metian et al., 2007) and for ^{110m}Ag (*P. maximus*: contribution over 98%; Metian et al., 2008) although there is a non-negligible effect of the type of food used in the model.

5. Conclusion

The pectinids, *M. varia* included, stand out from bivalves group when one considers their ability to retain ^{134}Cs . This characteristic makes them very good candidates to be used in biomonitoring programmes. However, our study highlighted interspecific differences among Pectinids regarding accumulation of waterborne radionuclides: *M. varia* shown a higher capacity to concentrate ^{134}Cs




Fig. 2. Comparison of concentration factors (CF) and percentages of long-term component absorption efficiency (A_{0t}) for different taxa of molluscs in experimental conditions. Values are mean \pm SD. When the values were not directly available, a graphical estimate was performed from uptake and depuration kinetics. Bars are ranges of values. Light grey and dark grey bars represent respectively seawater and food pathways. Black vertical lines indicate means calculated from all the considered values. The numbers next to grey bars represent the number of values below and above the calculated means. ^aPresent study and Metian et al. (2011), ^b(Ueda et al., 1978; Nolan and Dahlgaard, 1991; Hutchins et al., 1998; Ke et al., 2000; Wang et al., 2000; Güngör et al., 2001; Qureshi et al., 2007; Kalaycı et al., 2013), ^c(Bryan, 1963; Bustamante et al., 2006; Lacoue-Labarthe et al., 2010).

but lower absorption efficiency than *P. maximus*. A global bioaccumulation model indicated that ingestion with food is the major uptake pathway of Cs in *M. varia*. In light with the general recommendation to use multiple species in biomonitoring (e.g. Philips and Rainbow, 1993; Warnau et al., 1996), the present work strongly suggests that the use of *M. varia* in such programmes would be an excellent way to assess the trophically-available Cs and would ideally complement information provided by other species accumulation Cs predominantly from the dissolved phase.

Acknowledgments

The IAEA is grateful for the support provided to its Environment Laboratories by the Government of the Principality of Monaco. MW is an Honorary Senior Research Associate of the National Fund for Scientific Research (NFSR, Belgium). The authors thank L. Hédouin (CRNS-LABEX) for assistance on the contribution model and M. de Ré for his assistance on the scallop collection.

References

- Bailly du Bois, P., Laguionie, P., Boust, D., Korsakissok, I., Didier, D., Fiévet, B., 2012. Estimation of marine source-term following Fukushima Dai-ichi accident. *J. Environ. Radioact.* 114, 2–9.
- Bryan, G.W., 1963. The accumulation of radioactive caesium by marine invertebrates. *J. Mar. Biol. Assoc. U. K.* 43, 519–539.
- Bustamante, P., Germain, P., Leclerc, G., Miramand, P., 2002. Concentration and distribution of ²¹⁰Po in the tissues of the scallop *Chlamys varia* and the mussel *Mytilus edulis* from the coasts of Charente-Maritime (France). *Mar. Pollut. Bull.* 44, 997–1002.
- Bustamante, P., Miramand, P., 2004. Interspecific and geographical variations of trace element concentrations in Pectinidae from European waters. *Chemosphere* 57, 1355–1362.
- Bustamante, P., Miramand, P., 2005. Subcellular and body distributions of 17 trace elements in the variegated scallop *Chlamys varia* from the French coast of the Bay of Biscay. *Sci. Total Environ.* 337, 59–73.
- Bustamante, P., Teyssié, J.-L., Fowler, S.W., Warnau, M., 2006. Assessment of the exposure pathway in the uptake and distribution of americium and cesium in cuttlefish (*Sepia officinalis*) at different stages of its life cycle. *J. Exp. Mar. Biol. Ecol.* 331, 198–207.
- Chino, M., Nakayama, H., Nagai, H., Terada, H., Katata, G., Yamazawa, H., 2011. Preliminary estimation of release amounts of ¹³¹I and ¹³⁷Cs accidentally discharged from the Fukushima Daiichi nuclear power plant into the atmosphere. *J. Nucl. Sci. Technol.* 48, 1129–1134.
- Cranmore, G., Harrison, F.L., 1975. Loss of ¹³⁷Cs and ⁶⁰Co from the oyster *Crassostrea gigas*. *Health Phys.* 28 (4), 319–333.
- Güngör, N., Tuğrul, B., Topcuoğlu, S., Güngör, E., 2001. Experimental studies on the biokinetics of ¹³⁴Cs and ²⁴¹Am in mussels (*Mytilus galloprovincialis*). *Environ. Int.* 27, 259–264.
- Hédouin, L., Metian, M., Teyssié, J.-L., Fichez, R., Warnau, M., 2010a. Delineation of heavy metal contamination pathways (seawater, food and sediment) in tropical oysters from New Caledonia using radiotracer techniques. *Mar. Pollut. Bull.* 61, 542–553.
- Hédouin, L., Metian, M., Lacoue-Labarthe, T., Fichez, R., Teyssié, J.-L., Bustamante, P., Warnau, M., 2010b. Influence of food on the assimilation of selected metals in tropical bivalves from the New Caledonia lagoon: qualitative and quantitative aspects. *Mar. Pollut. Bull.* 61, 568–575.
- Hewett, C.J., Jefferies, D.F., 1978. The accumulation of radioactive caesium from food by the plaice (*Pleuronectes platessa*) and the brown trout (*Salmo trutta*). *J. Fish. Biol.* 13, 143–153.
- Hutchins, D.A., Stupakoff, L., Hook, S., Luoma, S.N., Fisher, N.S., 1998. Effects of arctic temperatures on distribution and retention of the nuclear waste radionuclides ²⁴¹Am, ⁵⁷Co, and ¹³⁷Cs in the bioindicator bivalve *Macoma balthica*. *Mar. Environ. Res.* 45, 17–28.
- Japan Chemical Analysis Center (JCAC), 2002. Radioactivity Survey Data in Japan (Part 2: Dietary Materials). NIRS-RSD No 133. National Institute of Radiological Sciences, Chiba, Japan, p. 34.
- Kalaycı, G., Belivermiş, M., Kılıç, Ö., Topcuoğlu, S., Çotuk, Y., 2013. Investigation of radiocesium biokinetics in Manila clam (*Ruditapes philippinarum*). *J. Radioanal. Nucl. Chem.* 295, 239–244.
- Ke, C., Yu, K.N., Lam, P.K.S., Wang, W.-X., 2000. Uptake and depuration of cesium in the green mussel *Perna viridis*. *Mar. Biol.* 137, 567–575.
- Lacoue-Labarthe, T., Warnau, M., Oberhansli, F., Teyssié, J.-L., Bustamante, P., 2010. Contrasting accumulation biokinetics and distribution of ²⁴¹Am, Co, Cs, Mn and Zn during the whole development time of the eggs of the common cuttlefish, *Sepia officinalis*. *J. Exp. Mar. Biol. Ecol.* 382, 131–138.
- Landrum, P.F., Lydy, M.J., Lee, H., 1992. Toxicokinetics in aquatic systems: model comparisons and use in hazard assessment. *Environ. Toxicol. Chem.* 11, 1709–1725.
- Marzano, F.N., Fiori, F., Jia, G., Chiantore, M., 2000. Anthropogenic radionuclides bioaccumulation in Antarctic marine fauna and its ecological relevance. *Polar Biol.* 23, 753–758.
- Mauri, M., Orlando, E., Nigro, M., Regoli, F., 1990. Heavy metals in the Antarctic scallop *Adamussium colbecki*. *Mar. Ecol. Prog. Ser.* 67, 27–33.
- Metian, M., Warnau, M., Oberhansli, F., Teyssié, J.-L., Bustamante, P., 2007. Interspecific comparison of Cd bioaccumulation in European Pectinidae (*Chlamys varia* and *Pecten maximus*). *J. Exp. Mar. Biol. Ecol.* 353, 58–67.
- Metian, M., Bustamante, P., Cosson, R.P., Hédouin, L., Warnau, M., 2008. Investigation of Ag in the king scallop *Pecten maximus* using field and laboratory approaches. *J. Exp. Mar. Biol. Ecol.* 367, 53–60.
- Metian, M., Bustamante, P., Hédouin, L., Oberhansli, F., Warnau, M., 2009a. Delineation of heavy metal uptake pathways (seawater and food) in the variegated scallop *Chlamys varia* using radiotracer techniques. *Mar. Ecol. Prog. Ser.* 375, 161–171.
- Metian, M., Warnau, M., Hédouin, L., Bustamante, P., 2009b. Bioaccumulation of essential metals (Co, Mn and Zn) in the king scallop *Pecten maximus*: seawater, food and sediment exposures. *Mar. Biol.* 156, 2063–2075.

- Metian, M., Warnau, M., Oberh ansli, F., Bustamante, P., 2009c. Delineation of Pb contamination pathways in two Pectinidae: the variegated scallop *Chlamys varia* and the king scallop *Pecten maximus*. *Sci. Total Environ.* 407, 3503–3509.
- Metian, M., Warnau, M., Teyssi , J.-L., Bustamante, P., 2011. Characterization of ^{241}Am and ^{134}Cs bioaccumulation in the king scallop *Pecten maximus*: investigation via three exposure pathways. *J. Environ. Radioact.* 102, 543–550.
- Nolan, C., Dahlgaard, H., 1991. Accumulation of metal radiotracers by *Mytilusedulis*. *Mar. Ecol. Prog. Ser.* 70, 165–174.
- Pentreath, R.J., Jefferies, D.F., 1971. The uptake of radionuclides by I-group plaice (*Pleuronectes platessa*) off the Cumberland coast, Irish sea. *J. Mar. Biol. Assoc. U. K.* 51, 963–976.
- Phillips, D.J.H., Rainbow, P.S., 1993. *Biomonitoring of Trace Aquatic Contaminants*. Elsevier Applied Science, London.
- Qureshi, R.M., Mashiatullah, A., Yaqoob, N., Akhtar, P., Chaghtai, F., Jabbar, A., Warnau, M., 2007. Bioaccumulation of ^{137}Cs , Zn and Cr[VI] in the green mussel *Perna viridis*: influence of salinity and temperature. *Environ. Bioindic.* 2, 245–252.
- Rodr guez y Baena, A.M., Metian, M., Teyssi , J.-L., De Broyer, C., Warnau, M., 2006. Experimental evidence for ^{234}Th bioaccumulation in three Antarctic crustaceans: potential implications for particle flux studies. *Mar. Chem.* 100, 354–365.
- Ryan, T.P., 2002. Transuranic biokinetic parameters for marine invertebrates: a review. *Environ. Int.* 28, 83–96.
- Shumway, S.E., Parsons, J.G.J., 2006. *Scallops: Biology, Ecology and Aquaculture*, second ed. Elsevier, Amsterdam.
- Ueda, T., Nakamura, R., Suzuki, Y., 1978. Comparison of Influences of Sediments and Sea Water on Accumulation of Radionuclides by Marine Organisms.
- Wang, W.-X., Ke, C., Yu, K.N., Lam, P.K.S., 2000. Modeling radiocesium bioaccumulation in a marine food chain. *Mar. Ecol. Prog. Ser.* 208, 41–50.
- Warnau, M., Fowler, S.W., Teyssi , J.-L., 1996a. Biokinetics of selected heavy metals and radionuclides in two marine macrophytes: the seagrass *Posidonia oceanica* and the alga *Caulerpa taxifolia*. *Mar. Environ. Res.* 41, 343–362.
- Warnau, M., Teyssi , J.-L., Fowler, S.W., 1996b. Biokinetics of selected heavy metals and radionuclides in the common Mediterranean echinoid *Paracentrotus lividus*: sea water and food exposures. *Mar. Ecol. Prog. Ser.* 141, 83–94.