



## Deleterious effects of mercury contamination on immunocompetence, liver function and egg volume in an antarctic seabird

Andrés E. Ibañez <sup>a,\*</sup>, William F. Mills <sup>b</sup>, Paco Bustamante <sup>c</sup>, Lara M. Morales <sup>a</sup>, Diego S. Torres <sup>a</sup>, Beatriz D' Astek <sup>a</sup>, Rocío Mariano-Jelicich <sup>d</sup>, Richard A. Phillips <sup>e</sup>, Diego Montalti <sup>a,f</sup>

<sup>a</sup> Sección Ornitológia, Div. Zool. Vert. Museo de la Plata (FCNyM-UNLP-CONICET), La Plata, Buenos Aires, Argentina

<sup>b</sup> Department of Geography and Environmental Science, University of Reading, Reading, UK

<sup>c</sup> Littoral Environnement et Sociétés (LIENSS), UMR 7266, CNRS-La Rochelle Université, 2 rue Olympe de Gouges, 17000, La Rochelle, France

<sup>d</sup> Instituto de Investigaciones Marinas y Costeras (IIMyC), UNMdP-CONICET, Universidad Nacional de Mar del Plata, Mar del Plata, Argentina


<sup>e</sup> British Antarctic Survey, Natural Environment Research Council, Cambridge, CB3 0ET, UK

<sup>f</sup> Instituto Antártico Argentino, San Martín, Buenos Aires, Argentina

### HIGHLIGHTS

- Blood THg concentrations were measured in brown skuas on the Antarctic Peninsula.
- Higher blood THg concentrations had deleterious effects on physiology.
- Higher blood THg concentrations disrupted immune and liver function.
- Higher blood THg concentrations were associated with lower egg volume.

### GRAPHICAL ABSTRACT



### ARTICLE INFO

Handling Editor: Michael Bank

**Keywords:**  
Pollution  
Mercury  
Seabirds  
Physiology  
Reproduction  
Antarctica

### ABSTRACT

Mercury (Hg) is a globally important pollutant that can negatively impact metabolic, endocrine and immune systems of marine biota. Seabirds are long-lived marine top predators and hence are at risk of bioaccumulating high Hg concentrations from their prey. Here, we measured blood total mercury (THg) concentrations and relationships with physiology and breeding parameters of breeding brown skuas (*Sturnarius antarcticus*) ( $n = 49$  individuals) at Esperanza/Hope Bay, Antarctic Peninsula. Mean blood THg concentrations were similar in males and females despite the differences in body size and breeding roles, but differed between study years. Immune markers (hematocrit, Immunoglobulin Y [IgY] and albumin) were negatively correlated with blood THg concentrations, which likely indicates a disruptive effect of Hg on immunity. Alanine aminotransferase (GPT)

\* Corresponding author. Div. Zool. Vert. Museo de La Plata (FCNyM-UNLP), Paseo del Bosque s/n, (B1900FWA), La Plata, Buenos Aires, Argentina.  
E-mail address: [aeibanez@fcnym.unlp.edu.ar](mailto:aeibanez@fcnym.unlp.edu.ar) (A.E. Ibañez).

activity, reflecting liver dysfunction, was positively associated with blood THg. Additionally, triacylglycerol and albumin differed between our study years, but did not correlate with Hg levels, and so were more likely to reflect changes in diet and nutritional status rather than Hg contamination. Egg volume correlated negatively with blood THg concentrations. Our study provides new insights into the sublethal effects of Hg contamination on immunity, liver function and breeding parameters in seabirds. In this Antarctic species, exposure to sublethal Hg concentrations reflects the short-term risks which could make individuals more susceptible to environmental stressors, including ongoing climatic changes.

## 1. Introduction

Mercury (Hg) is a global environmental pollutant that can have deleterious consequences for humans and wildlife (Corsolini, 2009; Tan et al., 2009; Driscoll et al., 2013). Although Hg occurs naturally, its availability has increased in marine and terrestrial environments in different organic and inorganic forms (MeHg, inorganic Hg, etc) (Aronson et al., 2011). The gaseous, elemental form of Hg ( $Hg^0$ ) can spread through atmospheric transport from emission sources to distant regions (Calle et al., 2015). Once deposited in marine environments, the inorganic form of Hg ( $Hg^{II}$ ) is methylated by microorganisms to the more toxic form, methyl-Hg (MeHg,  $[CH_3Hg]^+$ ). MeHg bioaccumulates within the tissues of marine organisms and biomagnifies through marine food webs (Bargagli, 2008; Driscoll et al., 2013; Ibañez et al., 2022a; Matias et al., 2022). Seabirds, which are typically long-lived and feed at high trophic positions, are potentially at risk of accumulating high Hg levels in their tissues from dietary exposure (Bearhop et al., 2000; Bargagli, 2008; Tavares et al., 2013; McKenzie et al., 2021; Mills et al., 2020, 2022; Ibañez et al., 2022a).

Hg in Antarctic ecosystems derives from natural and human sources, and the Antarctic continent constitutes a sink for Hg, which condenses in colder regions after evaporation and long-range transportation from lower latitudes (Angot et al., 2016). Artisanal and small-scale gold mining, which releases large quantities of Hg into the environment, is mostly concentrated in the Southern Hemisphere (Keane et al., 2023). There are also local sources of Hg, including volcanic activity and the release of Hg stored in sea ice which increase its bioavailability for microbial methylation (Cossa, 2013; de Ferro et al., 2014; Gionfriddo et al., 2016). Human activity that is associated with nearby research stations on the Antarctic Peninsula increases during spring and summer, which results in the release of pollutants (heavy metals and organic compounds) from waste-disposal sites, construction materials and compounds used for treating effluent (Acero et al., 1996).

Hg is a neurological, immune and endocrine disruptor (Tartu et al., 2013; Whitney and Cristol, 2017), and, ultimately, can Hg contamination can have short- or long-term fitness consequences for seabirds (Bustnes et al., 2007; Roos et al., 2012; Dietz et al., 2019; Chételat et al., 2020; Mills et al., 2020; Goutte et al., 2014a, 2014b). Hg contamination may also negatively impact body condition (Tan et al., 2009; Ackerman et al., 2016; Chételat et al., 2020), although, in the wild, body condition indices are unreliable indicators of Hg sublethal effects (Caravari et al., 2022). Although many studies have examined inter- and intraspecific variation in Hg contamination of seabirds in the Southern Ocean (Caravari et al., 2014a; 2016, 2017; Mills et al., 2020, 2022; Quillfeldt et al., 2023), few have tested for relationships with fitness parameters. Indeed, these have only been investigated for the wandering albatross (*Diomedea exulans*) (Tavares et al., 2013; Caravari et al., 2014b, 2014c; Goutte et al., 2014a; Bustamante et al., 2016), Antarctic petrel (*Thalassula antarctica*) (Caravari et al., 2018, 2021), grey-headed albatross (*Thalassarche chrysostoma*) (Mills et al., 2020), and brown skua (*Stercorarius antarcticus*) (Ibañez et al., 2022a) and south polar skua (*S. maccormicki*) (Goutte et al., 2014b).

Brown skuas breed on the Antarctic continent and sub-Antarctic islands and are opportunistic predators that feed on a wide diversity of prey in both terrestrial and marine environments (Reinhardt et al., 2000; Phillips et al., 2004; Ritz et al., 2008; Carneiro et al., 2015; Graña

Grilli and Montalti, 2015; Borghello et al., 2019; Ibañez et al., 2022b). As the brown skua is a migratory seabird, birds are exposed to pollutants not just during the breeding season but also in the nonbreeding season when they visit regions with higher anthropogenic pressure (Carneiro et al., 2016; Albert et al., 2022). At Bahía Esperanza/Hope Bay—located on the Antarctic Peninsula—brown skuas breed close to large colonies of Adélie penguins (*Pygoscelis adeliae*) and gentoo penguins (*Pygoscelis papua*). They feed mainly on penguins and to a lesser extent on marine prey (e.g., fishes and invertebrates) (Borghello et al., 2019; Ibañez et al., 2022b). Macro-plastics have been found recently in skua diet samples collected during the breeding season (Ibanez et al., 2020), which may represent a possible route for chemical pollutants such as Hg (Lim et al., 2022; Hamilton et al., 2023). Overall, there has been little research concerning the physiological effects of Hg contamination on Antarctic seabirds (Caravari et al., 2021; Goutte et al., 2014), and more studies are needed to understand the potential sublethal effects. In the present study, we focus on the sublethal effects of Hg contamination of brown skuas. Total Hg (the sum of inorganic and organic Hg) concentration was measured in red blood cells of brown skuas at Esperanza/Hope Bay. Our aims were to: (i) compare blood THg concentrations between two breeding seasons and sexes; and (ii) relate blood THg concentrations to markers of energy metabolism, immunocompetence and liver function, and to egg volume.

## 2. Materials and methods

### 2.1. Fieldwork and sample collection

Fieldwork for this study was conducted at Bahía Esperanza/Hope Bay, Antarctic Peninsula ( $63^\circ 24' S$ ,  $57^\circ 01' W$ ) (Fig. 1), from November to January during the 2018/19 and 2019/20 breeding seasons. Brown skuas were sampled during the early incubation stage (5–10 days after clutch completion). Blood samples (~2 ml) were obtained from the brachial vein using a 25-G needles ( $n = 49$ ,  $n = 24$  and  $n = 25$  in 2018/19 and 2019/20, respectively). Sampled birds included both individuals from 16 nests and one adult from 17 nests. Once in the laboratory (within 2–6 h after extraction) serum and red blood cells were separated by centrifugation (2000 rpm for 10 min), placed in sterile plastic eppendorf tubes and were stored frozen ( $-20^\circ C$ ) prior to laboratory analyses.

Eggs were measured (length and breadth) using digital calipers and volumes ( $mm^3$ ) calculated as  $0.00048 \times \text{length (mm)} \times \text{breadth (mm)}^2$  (Phillips et al., 2004). We calculated the total clutch volume as the average of the volume of both eggs. The sex of birds was initially assigned morphologically based on body size, and later confirmed by DNA analysis (Fridolfsson and Ellegren, 1999; Phillips et al., 2002).

### 2.2. Total Hg analysis

Studies have demonstrated that Hg in blood is associated predominantly with the cellular fraction (i.e., red blood cells) rather than plasma (Bond and Robertson, 2015; Renedo et al., 2018). THg in seabird RBCs is mostly (>90%) MeHg (Renedo et al., 2018; Albert et al., 2019). Prior to analysis, red blood cells were freeze-dried and homogenized. Blood THg concentrations were measured using an Advanced Mercury Analyser spectrophotometer (Altec AMA 254) (LIENSS, France). For each sample,

a minimum of two aliquots (range: 1.02–1.86 mg dry weight [dw]) were analyzed, and the means and relative standard deviation (RSD) among measurements were calculated (all samples RSD <10%). THg concentrations are presented in  $\mu\text{g g}^{-1}$  dw. Accuracy was tested using certified reference material (CRM; dogfish liver DOLT-5, NRC, Canada; certified Hg concentration:  $0.44 \pm 0.18 \mu\text{g g}^{-1}$  dw) every 10 samples. Recovery of the CRM was  $97.8 \pm 1.7\%$ . Blanks were analyzed at the beginning of each set of samples. The limit of quantification of the AMA was 0.1 ng and the detection limit of the method was  $0.005 \mu\text{g g}^{-1}$  dw.

### 2.3. Hematological determinations

Serum concentrations of five energy metabolism markers (including total proteins, uric acid, triacylglycerol, cholesterol and glucose), albumin, aspartate aminotransferase (GOT) and alanine aminotransferase (GPT) enzymes were measured in each bird using colorimetric commercial kits (Wiener Lab). All assays were conducted using an automatic analyzer (Ibañez et al., 2015).

Serum circulating levels of Immunoglobulin Y (IgY) were determined by direct ELISA using peroxidase conjugated anti-chicken IgY antibodies (Sigma, St Louis, MO, USA, A-9046) (Martínez et al., 2003; Ibañez et al., 2018). For this, 96-well microtiter plates (Nunc PolySorp; Nunc, Roskilde, Denmark) were coated during 1 h at 37 °C with serum samples diluted (1/30 000) in 0.1 M carbonate-bicarbonate buffer (pH = 9.6). Then the plates were washed three times with PBS- 0.05% Tween 20 and incubated with 1% non-fat milk (Nestlé coffee-mate) in PBS-Tween-20 during 1 h at 37 °C to block the free binding sites. After new washing, the wells were incubated with peroxidase-conjugated anti-chicken IgY. Finally, the wells were washed and ABTS [2,2-azino-di (3-ethylbenzthiazoline sulfonate)] was added as substrate. After incubating for 30 min (at room temperature) color development was stopped with oxalic acid 2% and then read as optical density (OD) at 405 nm.

To determine the hematocrit value a heparinized capillary was filled in the laboratory with 100  $\mu\text{l}$  of blood from the Eppendorf tube that contained heparinized blood obtained in the field. The capillary was then centrifuged at 5000 rpm for 15 min (Ibañez et al., 2015), and a digital caliper used to measure the length (mm) of the red blood cell fraction and the total blood volume. Hematocrit values are presented as a percentage of total volume.

### 2.4. Statistical analysis

Data were analyzed using R (R Core Team, 2015). Blood THg concentrations, physiological parameters and egg volumes were checked for normality and homogeneity of variances using Shapiro-Wilk and Levene's tests, respectively. Differences in blood THg between sexes and seasons were tested using general linear mixed models (GLMMs; Gaussian distribution and identity link function), with the individual identity and the breeding pair included as random effects to account for partners potentially being more similar than non-partners in blood THg concentrations. Relationships between physiological markers (total proteins, uric acid, triacylglycerol, cholesterol, hematocrit, albumin, IgY, GOT and GPT), egg volume and blood THg concentrations were also tested using GLMMs, with sex (except in the case of egg volume) and season included as covariates, and individual identity included as a random effect. All GLMMs were fitted using the "nlme" package in R (Pinheiro et al., 2017).

## 3. Results


### 3.1. Sex and annual variation in Hg contamination

Detectable blood THg concentrations were found in all samples from brown skuas at Bahía Esperanza/Hope Bay in 2018/19 (mean  $\pm$  SD,  $0.73 \pm 0.22 \mu\text{g g}^{-1}$  dw) and 2019/20 ( $0.91 \pm 0.44 \mu\text{g g}^{-1}$  dw) (Table 1). There were no significant differences in blood THg concentrations (log-

**Table 1**

Total mercury (THg) concentrations ( $\mu\text{g g}^{-1}$  dw) in red blood cells of adult male and female brown skuas (*Stercorarius antarcticus*) at Esperanza/Hope Bay, Antarctic Peninsula ( $63^{\circ}24'S$ ,  $57^{\circ}01'W$ ), in the 2018/19 and 2019/20 breeding seasons. Data are means  $\pm$  SDs.

Year	N	Sex	THg ( $\mu\text{g g}^{-1}$ dw)	Range ( $\mu\text{g g}^{-1}$ dw)
2018–2019	11	F	$0.66 \pm 0.25$	0.41–1.21
	13	M	$0.79 \pm 0.18$	0.48–1.06
	24	Both	$0.73 \pm 0.22$	0.41–1.21
2019–2020	13	F	$0.82 \pm 0.39$	0.49–1.94
	12	M	$1.02 \pm 0.50$	0.44–2.33
	25	Both	$0.91 \pm 0.44$	0.44–2.33



**Fig. 1.** Location of the study site (black circle), Esperanza/Hope Bay, Antarctic Peninsula ( $63^{\circ}24'S$ ,  $57^{\circ}01'W$ ).

transformed) between sexes (est = 0.23, p = 0.48) or seasons (est = 0.19, p = 0.56) (Table 1).

### 3.2. Physiological markers and egg volume

Physiological parameters, in particular immune and hepatic functions, did not differ between sexes (all p > 0.05), but were negatively associated with blood THg concentrations. Significant negative relationships were found between albumin (est = -0.31, p < 0.0001), hematocrit (%) (est = -9.27, p < 0.0001) and IgY (est = -0.06, p < 0.05) and blood THg concentrations (Fig. 2; Table 2). A significant positive relationship was found between GPT activity and blood THg concentrations (est = 13.10, p < 0.01) (Fig. 2; Table 2). Mean albumin levels differed significantly between seasons (est = 0.31, p < 0.0001). Triacylglycerol levels also varied annually (est = 55.89, p < 0.001) and were unrelated to blood THg concentrations (est = -11.04, p = 0.55) (Table 2). Uric acid, total proteins, cholesterol, glucose and GOT show no relationship with sex, season or blood THg (Table 2). Egg volume ( $\text{mm}^3$ ) showed a significant negative relationship with blood THg concentrations (est = -16.40, p < 0.05) (Fig. 3), and did not differ significantly between seasons (est = 4.25, p = 0.19).


## 4. Discussion

In our study, blood THg concentrations of brown skuas ranged from 0.41 to 2.33  $\mu\text{g g}^{-1}$  dw (equivalent to 0.10–0.58  $\mu\text{g g}^{-1}$  wet weight [ww], assuming a 79% moisture content; Eagles-Smith et al., 2018;


**Table 2**

Serum hematological markers of adult brown skuas (*Stercorarius antarcticus*) at Esperanza/Hope Bay, Antarctic Peninsula ( $63^{\circ}24'S$ ,  $57^{\circ}01'W$ ), in the 2018/19 and 2019/20 breeding seasons. Data are means  $\pm$  SDs. Uric acid (mg/dl), triacylglycerol (mg/dl), total proteins (d/dl), cholesterol (mg/dl), glucose (mg/dl), albumin (g/dl), GOT (U/l) and GPT (U/l). Serum hematological markers that showed an association with season are indicated with <sup>a</sup> (p < 0.05).

Parameter	Season 2018–2019		Season 2019–2020	
	Mean $\pm$ SD	Min-Max	Mean $\pm$ SD	Min-Max
Uric acid	8.126 $\pm$ 2.776	3.87–14.37	6.234 $\pm$ 2.069	2.250–9.56
Triacylglycerols	63.41 $\pm$ 18.40	31–99	118.1 $\pm$ 49.76	53–286
Total proteins	2.809 $\pm$ 0.392	2.1–3.7	2.790 $\pm$ 0.497	1.9–3.5
Cholesterol	245 $\pm$ 58.06	103.6–354.9	244.8 $\pm$ 65.63	113.3–351.5
Glucose	340.6 $\pm$ 43.41	269–420	330.6 $\pm$ 49.95	170–379
Albumin	1.020 $\pm$ 0.135	0.73–1.260	1.282 $\pm$ 0.233	0.87–1.63
GOT	75.95 $\pm$ 27.20	25–160	105.4 $\pm$ 20.79	69–153
GPT	27.70 $\pm$ 12.84	9.1–63	27.27 $\pm$ 7.192	15–39
Hematocrit (%)	45.60 $\pm$ 4.967	35.6–53.5	41.33 $\pm$ 6.204	30.10–30.90
IgY (OD 405 nm)	0.215 $\pm$ 0.069	0.11–0.38	0.237 $\pm$ 0.075	0.104–0.430



**Fig. 2.** Relationships between red blood cells THg concentrations ( $\mu\text{g g}^{-1}$  dw) and (a) albumin (g/dl), (b) hematocrit (%), (c) IgY (OD 405 nm) and (d) GPT (U/l) in the blood of adult brown skuas (*Stercorarius antarcticus*) at Esperanza/Hope Bay, Antarctic Peninsula ( $63^{\circ}24'S$ ,  $57^{\circ}01'W$ ) in the 2018/19 and 2019/20 breeding seasons.



**Fig. 3.** Relationships between red blood cells THg concentrations ( $\mu\text{g g}^{-1}$  dw) and egg volume ( $\text{mm}^3$ ) of female brown skuas (*Stercorarius antarcticus*) at Esperanza/Hope Bay, Antarctic Peninsula ( $63^\circ 24'\text{S}$ ,  $57^\circ 01'\text{W}$ ) in the 2018/19 and 2019/20 breeding seasons.

(Ackerman et al., 2016). These concentrations are comparable to those found in adults of other Antarctic seabirds, including snow petrels (*Pagodroma nivea*) (Tartu et al., 2015), and Antarctic petrels (*Thalassarche antarctica*) (Carravieri et al., 2021). However, they were lower than those reported in skuas at lower latitude breeding colonies in the southwest Atlantic Ocean sector of the Southern Ocean (Mills et al., 2022), but were comparable to those associated with reduced breeding success in south polar skuas at Adélie Land (Goutte et al., 2014b). Despite the low values of blood THg measured here, we found significant negative effects on immunity and hepatic enzymes, and on breeding parameters.

#### 4.1. Impacts of Hg contamination on immunocompetence

Hg is often associated with immunosuppressive effects at sublethal levels, but mostly in captive studies (Fallacara et al., 2011; Kenow et al., 2007; Lewis et al., 2013a). Despite the relatively low concentrations (initially suggesting a low risk of MeHg toxicity; Ackerman et al., 2016), IgY and hematocrit were negatively related to blood THg concentrations in our study (Fig. 2). These results indicate a negative impact of Hg contamination on the immune status of brown skuas. Our results agree with previous studies on captive zebra finches (*Taeniopygia guttata*) that have shown negative effects of Hg exposure on B-cell proliferation (Lewis, 2012; Lewis et al., 2013). Also, impaired macrophage phagocytosis was related to high Hg levels in black-footed albatrosses (*Phoebastria nigripes*) in the North Pacific (Finkelstein et al., 2007). Hematocrit values decreased with blood THg concentrations, which agrees with the hemolytic and anemia-inducing effects of Hg (Zolla et al., 1997). Erythrocytes are an important target of Hg and the majority of Hg in blood is found in the cellular fraction (~90%) (Bond and Robertson, 2015). *In vitro* studies have demonstrated that the exposure of erythrocytes to low concentrations of Hg induce structural changes in the external surface of the membrane. These changes are mediated by the translocation of phosphatidylserine to the external surface of the erythrocyte cell membrane as a signal that may prompt cellular apoptosis (Eisele et al., 2006; Lim et al., 2010). Another possible explanation behind the decrease in hematocrit may be an association with lowered renal function, also linked to hemolytic processes (Rivarob et al., 1983; Chitra and Jayaprakash, 2013).

Serum IgY and albumin are useful indicators of health in birds (Lumeij, 1987; Ibañez et al., 2018). Fitness traits are mediated by hormones including luteinizing hormone, which is a pituitary hormone involved in the onset of breeding (Guillette et al., 1995; Dawson et al.,

2001); corticosterone, an adrenal hormone in the stress response (Wingfield and Sapolsky, 2003); and prolactin, a pituitary hormone involved in the expression of parental care (Angelier and Chastel, 2009). Trace metal pollution may have different effects on hormones of the hypothalamic–pituitary–adrenal (HPA) axis, such as corticosterone (Tan et al., 2009; Tartu et al., 2013). Serum IgY and albumin concentrations were negatively related to blood THg concentrations of brown skuas in our study (Fig. 2). One possible explanation for this may be increasing immunosuppressive effects and catabolic activity associated with endocrine disruption of the HPA axis (Coutinho and Chapman, 2011), as Hg accumulates in the pituitary gland and thyroid in vertebrates (Colborn et al., 1993; Tan et al., 2009; Meyer et al., 2014; Tartu et al., 2013). In this scenario, the levels of Hg found in brown skuas at Esperanza/Hope Bay may induce protein catabolism and immunosuppression (IgY and albumin decline) because corticosterone release is disrupted (Ibañez et al., 2018). Another possible explanation is impaired protein synthesis in the liver. However, these hypotheses would need to be confirmed in future studies on skuas, particularly as a Hg-induced deterioration in immunocompetence is likely to affect disease risk and, ultimately, population dynamics.

#### 4.2. Impacts of Hg contamination on liver function

GPT was positively related to blood THg concentrations of brown skuas. This suggests that Hg contamination impacted the liver function of brown skuas. The liver is the major organ involved in the biotransformation, metabolism, protein synthesis and detoxification processes, which also reflect its susceptibility to pollutants, leading to tissue damage. GOT and GPT enzymes are considered to be biochemical markers of impaired liver function. Liver-cell damage, such as degeneration and necrosis, may increase GOT and GPT levels (Gowda et al., 2009; Marí et al., 2010; Ibañez et al., 2015; Yang et al., 2015; Choi et al., 2017). The biological mechanism of association between Hg exposure and liver dysfunction is mainly explained by oxidative stress, cell death, and impaired metabolism (Malhi et al., 2010). For instance, in male rodents exposed to Hg, levels of GOT, GPT, and gamma glutamyl-transferase activities were elevated, and tissue damage or necrotic changes observed in most livers (Wadaan, 2009). Also, histological analyses described degenerative changes and lysed areas in liver parenchyma in Hg-exposed zebrafish (Maricella et al., 2016). Hg exposure in zebrafish induced deregulation of oxidative stress, intrinsic apoptotic pathways, and resulting hepatotoxicity through cell death, mitochondrial dysfunction, endocrine disruption, and metabolic disorders (Ung et al., 2010; Maricella et al., 2016). In the spectacled caiman (*Caiman crocodilus*), negative associations between Hg and alkaline phosphatase activity (a liver cytoplasmic enzyme involved in the hepatocytic functions) were observed (Lemaire et al., 2021). Therefore, our results suggest that variation in GPT activity in response to elevated blood Hg levels, may indicate hepatotoxicity even at the low levels of contamination found in Antarctica.

#### 4.3. Annual variations in Hg contamination and association with physiology

In a previous study of brown skuas at Hope Bay, we found annual variation in blood THg concentrations was related to trophic ecology, but had no impact on adult body condition (Ibañez et al., 2022a). Blood THg concentrations were higher in 2019/20 than 2018/19 (by ~0.2  $\mu\text{g g}^{-1}$  dw) (Ibañez et al., 2022a). This was potentially due to changes in diet composition, and the consumption of more contaminated prey (Braune et al., 2014). Also, annual fluctuations in environmental conditions may influence Hg transport, MeHg production, and bioavailability to marine predators and their prey (Cossa et al., 2011; Driscoll et al., 2013; Renedo et al., 2020). In the current study, mean albumin and triacylglycerol levels differed between years, but were not linked directly to blood Hg contamination. Albumin and triacylglycerol are

markers of nutritional status (Ibañez et al., 2018). In this scenario, annual dietary differences may affect nutritional status. This may in turn moderate the effects of Hg on metabolism, with potentially limited food resources (thus lower Hg levels) associated with lower albumins and triacylglycerols.

#### 4.4. Impacts of Hg contamination on egg volume

Hg contamination of seabirds may induce changes on body condition that ultimately affect breeding fitness and survival (Evers et al., 2008; Labocha and Hayes, 2012; Ackerman et al., 2016). Effects of Hg on reproduction can be reflected at different levels including egg neglect (Tartu et al., 2015), lower breeding success (Tartu et al., 2016), reproductive failure (Mills et al., 2020), or population dynamics (Goutte et al., 2014b, 2015). However, the effects of Hg contamination during the non-breeding season on subsequent breeding success are poorly known. In little auks (*Alle alle*), Fort et al. (2014) suggested a carry-over effect, in that individuals with the highest Hg concentrations laid smaller eggs. In the present study, egg volume was negatively associated with female blood THg concentrations (Fig. 3). Although not significant, THg concentrations were slightly higher in males than females in both seasons (Table 1; Mills et al., 2022; Ibañez et al., 2022a), possibly because egg production provides a route through which females are able to eliminate Hg (Robinson et al., 2012; Ackerman et al., 2020). An explanation for the association of smaller eggs laid with higher blood THg concentrations of brown skuas relates to the trophic ecology and Hg contamination during female pre-laying exodus or differences in prey consumption prior to sampling. Therefore, our results support a previous study which also suggested carry-over effects of Hg on the reproduction of great skuas (*Stercorarius skua*) with a specific influence of female winter distribution and Hg contamination on egg volume (Albert et al., 2022). However, this would need to be confirmed in future studies by measuring THg in feathers grown during the nonbreeding season to infer the degree of Hg contamination since the last moult (Fort et al., 2014).

The threshold of Hg toxicity in seabirds appears to be related to the latitude of the study site. Toxicity appears to differ in Antarctic compared with lower latitudes such as subantarctic or subtropical environments (Goutte et al., 2014b; Carravieri et al., 2021). Blood THg concentrations of brown skuas in our study would initially suggest a low risk of MeHg toxicity (Ackerman et al., 2016). These sublethal THg concentrations had deleterious effects on physiology and egg volume, but there was no relationship with breeding success (Ibañez et al., 2022a). Goutte et al. (2014b) reported short-term effects of Hg on breeding success in brown skuas and south polar skuas, though concentrations were higher than in brown skuas at our study site. Selenium (Se)-Hg interactions are often observed in the blood and internal tissues of marine predators, and Se has a protective effect against Hg toxicity when Se is in molar excess (Carravieri et al., 2017, 2020; Manceau et al., 2021). Our results may also be explained by the presence of Se, as was reported for brown skua chicks from the Southern Ocean (Carravieri et al., 2017), south polar skuas (Goutte et al., 2014b) and the spectacled caiman (Lemaire et al., 2021). If so, higher concentrations of Se at Antarctic latitudes may reduce the negative effects of Hg on physiology and reproduction; however, this hypothesis requires further investigation.

#### 5. Conclusions

Our study demonstrated negative effects of Hg contamination on physiology and breeding parameters in brown skuas on the Antarctic Peninsula. Despite low blood THg concentrations, which are below or similar to those of other Antarctic seabirds, the association between Hg and the physiological and breeding parameters are of concern. The detrimental impact on egg volume highlights the importance of investigating the relationships between blood THg and the hormones that play a role in stress responses and reproductive decisions, as well as, Hg

concentration in chicks during development. This is particularly as skuas and other predators in the Antarctic may become more susceptible to pollutants and other environmental stressors, given the evidence for rapid, ongoing climatic changes in the region.

#### Credit author statement

**Andrés E Ibañez:** Conceptualization, Methodology, Investigation, Resources, Formal Analysis, Visualization, Writing – Original Draft. **William F Mills:** Methodology, Investigation, Resources, Writing – Review & Editing. **Paco Bustamante:** Investigation, Writing – Review & Editing. **Lara M Morales:** Investigation, Writing – Review & Editing. **Diego S Torres:** Investigation, Writing – Review & Editing. **Beatriz D'Astek:** Investigation, Resources. **Rocío Mariano-Jelicich:** Investigation, Writing – Review & Editing. **Richard A Phillips:** Conceptualization, Methodology, Investigation, Resources, Supervision, Writing – Review & Editing. **Diego Montalti:** Investigation, Resources, Supervision, Writing – Review & Editing.

#### Ethical approval

All applicable international, national, and institutional guidelines for sampling, care and experimental use of animals for the study were followed as established by the Article III, Annex II of the Madrid Protocol, Law 24.216 (Taking, Harmful Intrusion and Introduction of Species) within the framework of the projects evaluated and approved by the Environment Office of the IAA and Dirección Nacional del Antártico (DNA) (permits number 2018-FEAMB-CT-GA-70 and 2019-FEAMB-CT-GA-66). In addition, all the protocols conducted on this project were evaluated and approved by the Institutional Committee for the Care and Use of Study Animals (CICUAE) of Natural Science and Museum Faculty (FCNyM-UNLP).

#### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Data availability

Data will be made available on request.

#### Acknowledgements

The authors are grateful to the Dirección Nacional del Antártico (DNA) and Instituto Antártico Argentino (IAA) for the support of the field-work activities and logistics in the Antarctic. Thanks to Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET) and Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) for the financial support of this project (PIP- CONICET no. 0158 to DM; PICT-2014-3323 and PIP-CONICET no. 0955 to AEI). The authors are grateful to Maud Brault-Favrou for her assistance with Hg analyses. Thanks are also due to the CPER (Contrat de Projet Etat-Région) and the FEDER (Fonds Européen de Développement Régional) for funding the AMA of LIENSs laboratory. Paco Bustamante is an honorary member of the IUF (Institut Universitaire de France). William Mills is supported by a Leverhulme Trust Early Career Fellowship at the University of Reading. This work represents a contribution to the Ecosystems component of the British Antarctic Survey Polar Science for Planet Earth Programme, funded by NERC.

#### References

- Acero, J.M., Agraz, J.L., Aguirre, C.A., 1996. Revisión ambiental de las actividades antárticas en Bahía Esperanza. Instituto Antártico Argentino, pp. 1–35.

- Ackerman, J.T., Eagles-Smith, C.A., Herzog, M.P., Hartman, C.A., Peterson, S.H., Evers, D.C., Jackson, A.K., Elliott, J.E., Vander Pol, S.S., Bryan, C.E., 2016. Avian mercury exposure and toxicological risk across western North America: a synthesis. *Sci. Total Environ.* 568, 749–769.
- Ackerman, J.T., Herzog, M.P., Evers, D.C., et al., 2020. Synthesis of maternal transfer of mercury in birds: implications for altered toxicity risk. *Environ. Sci. Technol.* 54, 2878–2891.
- Albert, C., Renedo, M., Bustamante, P., Fort, J., 2019. Using blood and feathers to investigate large-scale Hg contamination in Arctic seabirds: a review. *Environ. Res.* 171, 108588.
- Albert, C., Strom, H., Helgason, H.H., Bråthen, V.S., Gudmundsson, F.T., Bustamante, P., Fort, J., 2022. Spatial variations in winter Hg contamination affect egg volume in an Arctic seabird, the great skua (*Stercorarius skua*). *Environ. Pollut.* 314, 120322.
- Angelier, F., Chastel, O., 2009. Stress, prolactin and parental investment in birds: a review. *Gen. Comp. Endocrinol.* 163, 142–148.
- Angot, H., Dastoora, A., De Simone, F., Gårdfeldt, K., Gencarelli, C.N., Hedgecock, I.A., et al., 2016. Chemical cycling and deposition of atmospheric mercury in polar regions: review of recent measurements and comparison with models. *Atmos. Chem. Phys.* (16), 10735–10763.
- Arnon, R.B., Thatje, S., McClintonck, J.B., Hughes, K.A., 2011. Anthropogenic impacts on marine ecosystems in Antarctica. *Ann NY Acad Sci* 1223 (1), 82–107.
- Bargagli, R., 2008. Environmental contamination in Antarctic ecosystems. *Sci. Total Environ.* 400, 212–226.
- Bearhop, S., Waldron, S., Thompson, D.R., Furness, R.W., 2000a. Bioamplification of mercury in great skua *Catharacta skua* chicks: the influence of trophic status as determined by stable isotope signatures of blood and feathers. *Mar. Pollut. Bull.* 40, 181–185.
- Bearhop, S., Ruxton, G.D., Furness, R.W., 2000b. Dynamics of mercury in blood and feathers of great skuas. *Environ. Toxicol. Chem.* 19, 1638–1643.
- Bond, A.L., Robertson, G.J., 2015. Mercury concentrations in multiple tissues of Arctic Iceland Gulls (*Larus glaucopterus*) wintering in Newfoundland. *Arctic Sci* 1 (1), 1–8.
- Borghello, P., Torres, D.S., Montalti, D., Ibañez, A.E., 2019. Diet of the Brown skua (*Stercorarius antarcticus lonnbergi*) at Hope Bay, antarctic Peninsula: differences between breeders and non-breeders. *Polar Biol.* 42, 385–394.
- Braune, B.M., Gaston, A.J., Hobson, K.A., Gilchrist, H.G., Mallory, M.L., 2014. Changes in food web structure after trends of mercury uptake at two seabird colonies in the Canadian Arctic. *Environ. Sci. Technol.* 48, 13246–13252.
- Bustamante, P., Carravieri, A., Goutte, A., Barbraud, C., Delord, K., Chastel, O., Weimerskirch, H., Cherel, Y., 2016. High feather mercury concentrations in the wandering albatross are related to sex, breeding status and trophic ecology with no demographic consequences. *Environ. Res.* 144, 1–10.
- Bustnes, J.O., Tveraa, T., Varpe, Ø., Henden, J.A., Ju, Skaare, 2007. Reproductive performance and organochlorine pollutants in an Antarctic marine top predator: the south polar skua. *Environ. Int.* 33, 911–918.
- Carneiro, A.P.B., Manica, A., Trivelpiece, W.Z., Phillips, R.A., 2015. Flexibility in foraging strategies of Brown skuas in response to local and seasonal dietary constraints. *J. Ornithol.* 156, 625–633.
- Calle, P., Alvarado, O., Monserrate, L., Cevallos, JM, Calle, N., Alava, JJ., 2015. Mercury accumulation in sediments and seabird feathers from the Antarctic Peninsula. *Mar. Pollut. Bull.* 91, 410–417.
- Carneiro, A.P.B., Manica, A., Clay, T.A., Silk, J.R.D., King, M., Phillips, R.A., 2016. Consistency in migration strategies and habitat preferences of brown skuas over two winters, a decade apart. *Mar. Ecol. Prog. Ser.* 553, 267–281.
- Carravieri, A., Bustamante, P., Tartu, S., Meillère, A., Labadie, P., Budzinski, H., Peluhet, L., Barbraud, C., Weimerskirch, H., Chastel, O., Cherel, Y., 2014a. Wandering albatrosses document latitudinal variations in the transfer of persistent organic pollutants and mercury to Southern Ocean predators. *Environ. Sci. Technol.* 48, 14746–14755.
- Carravieri, A., Cherel, Y., Blévin, P., Brault-Favrou, M., Chastel, O., Bustamante, P., 2014b. Mercury exposure in a large subantarctic avian community. *Environ. Pollut.* 190, 51–57.
- Carravieri, A., Cherel, Y., Brault-Favrou, M., Churlaud, C., Peluhet, L., Labadie, P., Budzinski, H., Chastel, O., Bustamante, P., 2017. From Antarctica to the subtropics: contrasted geographical concentrations of selenium, mercury, and persistent organic pollutants in skua chicks (*Catharacta spp.*). *Environ. Pollut.* 228, 464–473.
- Carravieri, A., Fort, J., Tarroux, A., Cherel, Y., Love, O.P., Prieur, S., Brault-Favrou, M., Bustamante, P., Descamps, S., 2018. Mercury exposure and short-term consequences on physiology and reproduction in Antarctic petrels. *Environ. Pollut.* 237, 824–831.
- Carravieri, A., Warner, N.A., Herzke, D., Brault-Favrou, M., Tarroux, A., Fort, J., Bustamante, P., Descamps, S., 2021. Trophic and fitness correlates of mercury and organochlorine compound residues in egg-laying Antarctic petrels. *Environ. Res.* 193, 110518.
- Caravari, A., Vincze, O., Bustamante, P., Ackerman, J.T., Adams, E.M., Angelier, F., Chastel, O., Cherel, Y., Gilg, O., Golubova, E., Kitaysky, A., Luff, K., Seewagen, C.L., Strom, H., Will, A.P., Yannic, G., Giraudeau, M., Fort, J., 2022. Quantitative meta-analysis reveals no association between mercury and body condition in birds. *Biol. Rev.* 97 (4), 1253–1271.
- Chetelat, J., Ackerman, J.T., Eagles-Smith, C.A., Hebert, C.E., 2020. Methylmercury exposure in wildlife: a review of the ecological and physiological processes affecting contaminant concentrations and their interpretation. *Sci. Total Environ.* 711, 135117.
- Chitra, S., Jayaprakash, K., 2013. Effect of Mercury on blood components of fresh water edible fish *Labeo rohita*. *J. Acad. Indus. Res.* 1, 12.
- Choi, J., Bae, S., Lim, H., Lim, J.A., Lee, Y.H., Ha, M., Kwon, H.J., 2017. Mercury exposure in association with decrease of liver function in adults: a longitudinal study. *J. Prev. Med. Public Health.* 50, 377–385.
- Colborn, T., Vom Saal, F.S., Soto, A.M., 1993. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ. Health Perspect.* 101, 378–384.
- Core Team, R., 2015. R: A Language and Environment for Statistical Computing. URL R foundation for statistical computing, Vienna, Austria. R-proje ct.org/.
- Corsolini, S., 2009. Industrial contaminants in Antarctic biota. *J. Chromatogr. A* 1216, 598–612.
- Cossa, D., Heimbürger, L.E., Lannuzel, D., Rintoul, S.R., Butler, E.C.V., Bowie, A.R., Avery, B., Watson, R.J., Remenyi, T., 2011. Mercury in the Southern Ocean. *Geochim. Cosmochim. Acta* 75, 4037–4052.
- Cossa, D., 2013. Methylmercury manufacture. *Nat. Geosci.* (6), 810–811.
- Coutinho, A.E., Chapman, K.E., 2011. The anti-inflammatory and immunosuppressive effects of glucocorticoids, recent developments and mechanistic insights. *Mol. Cell. Endocrinol.* 335 (1), 2–13.
- Dawson, A., Thapliyal, J.P., 2001. The thyroid and photoperiodism. In: Dawson, A., Chatruvedi, C.M. (Eds.), *Avian Endocrinology*. Narosa, New Delhi, pp. 141–151.
- de Ferro, A.M., Mota, A.M., Canário, J., 2014. Pathways and speciation of mercury in the environmental compartments of Deception Island, Antarctica. *Chemosphere* 95, 227–233.
- Dietz, R., et al., 2019. Current state of knowledge on biological effects from contaminants on arctic wildlife and fish. *Sci. Total Environ.* 696, 133792.
- Driscoll, C.T., Mason, R.P., Chan, H.M., Jacob, D.J., Pirrone, N., 2013. Mercury as a global pollutant: sources, pathways, and effects. *Environ. Sci. Technol.* 47, 4967–4983.
- Eagles-Smith, C.A., Silbergeld, E.K., Basu, N., Bustamante, P., Diaz-Barriga, F., Hopkins, W.A., Kidd, K.A., Nyland, J.F., 2018. Modulators of mercury risk to wildlife and humans in the context of rapid global change. *Ambio* 47, 170–197.
- Eisele, K., Lang, P.A., Kempe, D.S., Klär, B.A., Niemoller, O., Wieder, T., et al., 2006. Stimulation of erythrocyte phosphatidylserine exposure by mercury ions. *Toxicol. Appl. Pharmacol.* 210, 116–122.
- Evers, D.C., Savoy, L.J., DeSorbo, C.R., et al., 2008. Adverse effects from environmental mercury loads on breeding common loons. *Ecotoxicology* 17, 69–81.
- Fallacara, D.M., Halbrook, R.S., French, J.B., 2011. Toxic effects of dietary methylmercury on immune system development in nestling American Kestrels (*Falco sparverius*). *Environ. Toxicol. Chem.* 30, 1328–1337.
- Finkelstein, M.E., Grasman, K.A., Croll, D.A., Tershy, B.R., Keitt, B.S., Jarman, W.M., Smith, D.R., 2007. Contaminant-associated alteration of immune function in black-footed albatross (*Phoebastria nigripes*), a North Pacific predator. *Environ. Toxicol. Chem.* 26, 1896–1903.
- Fort, J., Robertson, G.J., Gremillet, D., Traisnel, G., Bustamante, P., 2014. Spatial ecotoxicology: migratory arctic seabirds are exposed to mercury contamination while overwintering in the Northwest Atlantic. *Environ. Sci. Technol.* 48, 11560–11567.
- Fridolfsson, A.K., Ellegren, H., 1999. A simple and universal method for molecular sexing of non-ratite birds. *J. Avian Biol.* 30, 116–121.
- Gionfriddo, C.M., Tate, M.T., Wick, R.R., Schultz, M.B., Zemla, A., Thelen, M.P., et al., 2016. Microbial mercury methylation in Antarctic sea ice. *Nature Microbiology* (1), 16127.
- Goutte, A., Barbraud, C., Meillère, A., Carravieri, A., Bustamante, P., Labadie, P., Budzinski, H., Delord, K., Cherel, Y., Weimerskirch, H., Chastel, O., 2014a. Demographic consequences of heavy metals and persistent organic pollutants in a vulnerable long-lived bird, the wandering albatross. *Proc. Roy. Soc. B* 281, 20133313.
- Goutte, A., Bustamante, P., Barbraud, C., Delord, K., Weimerskirch, H., Chastel, O., 2014b. Demographic responses to mercury exposure in two closely related Antarctic top predators. *Ecology* 95, 1075–1086.
- Goutte, A., Barbraud, C., Herzke, D., Bustamante, P., Angelier, F., Tartu, S., Clément-Chastel, C., Moe, B., Bech, C., Gabrielsen, G.W., Bustnes, J.O., Chastel, O., 2015. Survival rate and breeding outputs in a high Arctic seabird exposed to legacy persistent organic pollutants and mercury. *Environ. Pollut.* 200, 1–9.
- Guillette Jr., L.J., Cree, A., Rooney, A.A., 1995. Biology of stress: interactions with reproduction, immunology and intermediary metabolism. In: Warwicker, C., Frye, F., Murphy, J.B. (Eds.), *Health and Welfare of Captive Reptiles*. Chapman and Hall, London, pp. 32–81.
- Gowda, S., Desai, P.B., Hull, V.V., Math, A.A., Vernek, S.N., Kulkarni, S.S., 2009. A review on laboratory liver function tests. *Pan. Afr. Med. J.* 3, 17.
- Graña Grilli, M., Montalti, D., 2015. Variation in diet composition during the breeding cycle of an Antarctic seabird in relation to its breeding chronology and that of its main food resource. *Polar Biol.* 38, 643–649.
- Hamilton, B.M., Baak, J.E., Vorkamp, K., Hammer, S., Granberg, M., Herzke, D., Provencher, J.F., 2023. Plastics as a carrier of chemical additives to the Arctic: possibilities for strategic monitoring across the circumpolar North. *Arctic Sci* 9, 284–296.
- Ibañez, A.E., Najle, R., Larsen, K., Montalti, D., 2015. Hematology, biochemistry and serum protein analyses of antarctic and non-antarctic skuas. *Waterbirds* 38 (2), 153–161.
- Ibañez, A.E., Graña Grilli, M., Figueiroa, A., Pari, M., Montalti, D., 2018. Declining health status of Brown Skua (*Stercorarius antarcticus lonnbergi*) parents and their offspring during chick development. *Polar Biol.* 41, 193–200.
- Ibanez, A.E., Morales, L.M., Torres, D.S., Borghello, P., Haidr, N.S., Montalti, D., 2020. Plastic ingestion risk is related to the anthropogenic activity and breeding stage in an Antarctic top predator seabird species. *Mar. Pollut. Bull.* 157, 111351.
- Ibañez, A.E., Mills, W.F., Bustamante, P., McGill, R.A.R., Morales, L.M., Palacio, F.X., Torres, D.S., Haidr, N.S., Mariano-Jelicich, R., Phillips, R.A., Montalti, D., 2022a. Variation in blood mercury concentrations in brown skuas (*Stercorarius antarcticus*) is

- related to trophic ecology but not breeding success or adult body condition. Mar. Pollut. Bull. 181, 113919.
- Ibañez, A.E., Morales, L.M., Torres, D.S., Borghello, P., Montalti, D., 2022b. Pellet analysis evidences flexibility in the diet of Brown skua (*Stercorarius antarcticus*) during breeding at Esperanza/Hope Bay (antarctic Peninsula). Polar Biol. 45, 419–426.
- Keane, S., Bernaudat, L., Davis, K.J., Stylo, M., Mutemeri, N., Singo, P., Twala, P., Mutemeri, I., Nakafeero, A., Etui, I.D., 2023. Mercury and artisanal and small-scale gold mining: review of global use estimates and considerations for promoting mercury-free alternatives. Ambio 52, 833–852.
- Kenow, K.P., Grasman, K.A., Hines, R.K., Meyer, M.W., Gendron-Fitzpatrick, A., Spalding, M.G., et al., 2007. Effects of methylmercury exposure on the immune function of juvenile common loons (*Gavia immer*). Environ. Toxicol. Chem. 26, 1460–1469.
- Labocha, M.K., Hayes, J.P., 2012. Morphometric indices of body condition in birds: a review. J. Ornithol. 153, 1–22.
- Lemaire, J., Bustamante, P., Mangione, R., Marquis, O., Churlaud, C., Brault-Favrou, M., Parenteau, C., Brischoux, F., 2021. Lead, mercury, and selenium alter physiological functions in wild caimans (*Caiman crocodilus*). Environ. Pollut. 286, 117549.
- Lewis, C.A., 2012. Sublethal Effects of Methyl Mercury on the Songbird Immune Response: an Experimental Study. Master's Thesis. The College of William and Mary, Virginia.
- Lewis, C.A., Cristol, D.A., Swaddle, J.P., Varian-Ramos, C.W., Zwollo, P., 2013. Decreased immune response in zebra finches exposed to sublethal doses of mercury. Arch. Environ. Contam. Toxicol. 64, 327–336.
- Lim, K.M., Kim, S., Noh, J.Y., Kim, K., Jang, W.H., Bae, O.N., Chung, S.M., Chung, J.H., 2010. Low-level mercury can Enhance procoagulant activity of erythrocytes: a new contributing factor for mercury-related thrombotic disease. Environ. Health Perspect. 118, 7.
- Lim, C., Kim, N., Lee, J., Yoon, Y., 2022. Potential of adsorption of diverse environmental contaminants onto microplastics. Water 14, 4086.
- Lumeij, J.T., 1987. The diagnostic value of plasma proteins and non-protein nitrogen substances in birds. Vet. Q. 9, 262–268.
- Guardia, A., Pellegrino, D., Bernabò, I., Tronci, V., Ebbesson, L.O., et al., 2016. Effects of two sublethal concentrations of mercury chloride on the morphology and metallothionein activity in the liver of zebrafish (*Danio rerio*). Int. J. Mol. Sci. 17 (3), 361.
- Manceau, A., Gaillot, A.-C., Glatzel, P., et al., 2021. *In vivo* formation of HgSe nanoparticles and Hg–tetraselenolate complex from methylmercury in seabirds: implications for the Hg–Se antagonism. Environ. Sci. Technol. 55 (3), 1515–1526.
- Malhi, H., Guicciardi, M.E., Gores, G.J., 2010. Hepatocyte death: a clear and present danger. Physiol. Rev. 90 (3), 1165–1194.
- Marí, M., Colell, A., Morales, A., von Montfort, C., Garcia-Ruiz, C., Fernández-Checa, J. C., 2010. Redox control of liver function in health and disease. Antioxidants Redox Signal. (11), 1295–1331.
- Martínez, J., Tomás, G., Merino, S., Arriero, E., Moreno, J., 2003. Detection of serum immunoglobulins in wild birds by direct ELISA: a methodological study to validate the technique in different species using antichicken antibodies. Funct. Ecol. 17, 700–706.
- Matias, R.S., Guiñaro, H.R., Bustamante, P., Seco, J., Chiavez, N., Fragão, J., Tavares, S., Ceia, F.R., Pereira, M.E., Barbosa, A., Xavier, J.C., 2022. Mercury biomagnification in an antarctic food web of the antarctic Peninsula. Environ. Pollut. 304, 119199.
- McKenzie, A.C., Silvestro, A.M., Marti, L.J., Emslie, S.D., 2021. Intraspecific variation in mercury,  $\delta^{15}\text{N}$ , and  $\delta^{13}\text{C}$  among 3 Adélie penguin (*Pygoscelis adeliae*) populations in the northern antarctic Peninsula region. Environ. Toxicol. Chem. 40, 2791–2801.
- Meyer, E., Eagles-Smith, C.A., Sparling, D., Blumenshine, S., 2014. Mercury exposure associated with altered plasma thyroid hormones in the declining Western Pond Turtle (*Emys marmorata*) from California Mountain Streams. Environ. Sci. Technol. 48, 2989–2996.
- Mills, W.F., Bustamante, P., McGill, R.A.R., Anderson, O.R.J., Bearhop, S., Cherel, Y., Votier, S.C., Phillips, R.A., 2020. Mercury exposure in an endangered seabird: long-term changes and relationships with trophic ecology and breeding success. Proc. R. Soc. A B 287, 20202683.
- Mills, W.F., Ibanez, A.E., Bustamante, P., Carneiro, A.P.B., Bearhop, S., Cherel, Y., Mariano-Jelicich, R., McGill, R.A.R., Montalti, D., Votier, S.C., Phillips, R.A., 2022. Spatial and sex differences in mercury contamination of skuas in the Southern Ocean. Environ. Pollut. 297, 118841.
- Phillips, R.A., Dawson, D.A., Ross, D.J., 2002. Mating patterns and reversed size dimorphism in Southern skuas (*Stercorarius lönbergi*). Auk 119, 858–863.
- Phillips, R.A., Phalan, B., Forster, I.P., 2004. Diet and long-term changes in population size and productivity of brown skuas *Catharacta antarctica lönbergi* at Bird Island, South Georgia. Polar Biol. 27, 555–561.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Eispach, S., Heisterkamp, B., Van Willigen, 2017. Package “Nlme”. Linear and Nonlinear Mixed Effects Models.
- Quillfeldt, P., Bedolla-Guzmán, Y., Libertelli, M.M., Cherel, Y., Massaro, M., Bustamante, P., 2023. Mercury in ten storm-petrel populations from the antarctic to the subtropics. Arch. Environ. Contam. Toxicol. 85, 55–72.
- Reinhardt, K., Hahn, S., Peter, H.U., Wemhoff, H., 2000. A review of the diets of Southern Hemisphere skuas. Mar. Ornithol. 28, 7–19.
- Renedo, M., Amouroux, D., Duval, B., Carravieri, A., Tessier, E., Barre, J., Berail, S., Pedrero, Z., Cherel, Y., Bustamante, P., 2018a. Seabird tissues as efficient biomonitoring tools for Hg isotopic investigations: implications of using blood and feathers from chicks and adults. Environ. Sci. Technol. 52, 4227–4234.
- Renedo, M., Amouroux, D., Pedrero, Z., Bustamante, P., Cherel, Y., 2018b. Identification of sources and bioaccumulation pathways of MeHg in subantarctic penguins: a stable isotopic investigation. Sci. Rep. 8 (1), 8865.
- Renedo, M., Bustamante, P., Cherel, Y., Pedrero, Z., Tessier, E., Amouroux, D., 2020. A ‘seabird-eye’ on mercury stable isotopes and cycling in the Southern Ocean. Sci. Total Environ. 742, 140499.
- Ribarov, S.R., Benov, L.C., Benchev, I.C., 1983. On the mechanism of mercury-induced Hemolysis. Gen. Physiol. Biophys. 2, 81–84.
- Ritz, M., Millar, C., Miller, G., Phillips, P., Ryan, P., Sternkopf, V., Liebers-Helbig, D., Hans-Ulrich, P., 2008. Phylogeography of the southern skua complex: rapid colonization of the southern hemisphere during a glacial period and reticulate evolution. Mol. Phylogenet. Evol. 49, 292–303.
- Robinson, S.A., Lajeunesse, M.J., Forbes, M.R., 2012. Sex differences in mercury contamination of birds: testing multiple hypotheses with meta-analysis. Environ. Sci. Technol. 46, 7094–7101.
- Roos, A.M., Backlin, B.M.V.M., Helander, B.O., Rigét, F.F., Eriksson, U.C., 2012. Improved reproductive success in otters (*Lutra lutra*), grey seals (*Halichoerus grypus*) and sea eagles (*Haliaeetus albicilla*) from Sweden in relation to concentrations of organochlorine contaminants. Environ. Pollut. 170, 268–275.
- Tan, S.W., Meiller, J.C., Mahaffey, K.R., 2009. The endocrine effects of mercury in humans and wildlife. Crit. Rev. Toxicol. 39, 228–269.
- Tartu, S., Goutte, A., Bustamante, P., Angelier, F., Moe, B., Clément-Chastel, C., Bech, C., Gabrielsen, G.W., Bustnes, J.O., Chastel, O., 2013. To breed or not to breed: endocrine response to mercury contamination by an Arctic seabird. Biol. Lett. 9, 20130317.
- Tartu, S., Angelier, F., Wingfield, J.C., Bustamante, P., Labadie, P., Budzinski, H., Weimerskirch, H., Bustnes, J.O., Chastel, O., 2015. Corticosterone, prolactin and egg neglect behavior in relation to mercury and legacy POPs in a long-lived Antarctic bird. Sci. Total Environ. 505, 180–188.
- Tartu, S., Bustamante, P., Angelier, F., Lendvai, A.Z., Moe, B., Blévin, P., Bech, C., Gabrielsen, G.W., Bustnes, J.O., Chastel, O., 2016. Mercury exposure, stress and prolactin secretion in an Arctic seabird: an experimental study. Funct. Ecol. 30, 596–604.
- Tavares, S., Xavier, J.C., Phillips, R.A., Pereira, M.E., Pardal, M.A., 2013. Influence of age, sex and breeding status on mercury accumulation patterns in the wandering albatross *Diomedea exulans*. Environ. Pollut. 181, 315–320.
- Ung, C.Y., Lam, S.H., Hlaing, M.M., Winata, C.L., Korzh, S., Mathavan, S., et al., 2010. Mercury-induced hepatotoxicity in zebrafish: *in vivo* mechanistic insights from transcriptome analysis, phenotype anchoring and targeted gene expression validation. BMC Genom. 11, 212.
- Wadaan, M.A., 2009. Effects of mercury exposure on blood chemistry and liver histopathology of male rats. J. Pharmacol. Toxicol. 4 (3), 126–131.
- Whitney, M.C., Cristol, D.A., 2017. Impacts of sublethal mercury exposure on birds: a detailed review. Rev. Environ. Contam. Toxicol. 244, 113–163.
- Wingfield, J.C., Sapolsky, R.M., 2003. Reproduction and resistance to stress: when and How. J. Neuroendocrinol. 15, 711–724.
- Yang, G., Zhou, Z., Cen, Y., Gui, X., Zeng, Q., Ao, Y., Li, Q., Wang, S., Li, J., Zhang, A., 2015. Death receptor and mitochondria-mediated hepatocyte apoptosis underlies liver dysfunction in rats exposed to organic pollutants from drinking water. Drug Des. Dev. Ther. 9, 4719–4733.
- Zolla, L., Lupidi, G., Bellelli, A., Amiconi, G., 1997. Effect of mercuric ions on human erythrocytes. Relationships between hypotonic swelling and cell aggregation. Biochim. Biophys. Acta 1328, 273–280.