

Variation of Total Mercury Concentrations in Different Tissues of Three Neotropical Caimans: Implications for Minimally Invasive Biomonitoring

Jérémy Lemaire^{1,2} · François Brischoux¹ · Olivier Marquis³ · Rosanna Mangione⁴ · Paco Bustamante^{2,5}

Received: 22 January 2021 / Accepted: 8 April 2021 / Published online: 26 April 2021 © The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

Abstract

Mercury (Hg) is a global environmental contaminant that affects ecosystems. It is known to biomagnify through food webs and to bioaccumulate especially in the tissues of top predators. Large-scale comparisons between taxa and geographic areas are needed to reveal critical trends related to Hg contamination and its deleterious effects on wildlife. Yet, the large variety of tissues (keratinized tissues, internal organs, blood) as well as the variability in the units used to express Hg concentrations (either in wet- or dry-tissue weight) limits straightforward comparisons between studies. In the present study, we assessed the moisture content that could influence the total Hg (THg) concentrations measured in several tissues (claws, scutes, total blood, and red blood cells) of three caiman species. We evaluated the moisture content from the different tissues to provide information on THg concentrations of moisture content, with the highest THg values found in keratinized tissues (scute keratinized tissues. In the blood, the relationship between body size and THg concentration in keratinized tissues. In the blood, the relationship between body size and THg concentration and trace elements quantification based on dry weight analytical procedures. In addition, the use of both blood and keratinized tissues offers the possibility to quantify different time scales of THg exposure by non-lethal sampling.

Mercury (Hg) is one of the major contaminants of concern in ecosystems (Ericksen et al. 2003; Fitzgerald et al. 2007; Selin 2009). In addition to naturally present geological Hg, human activities, such as deforestation, fossil fuel combustion, and gold mining activities, have been shown to increase the level of Hg in the environment, particularly in aquatic

Jérémy Lemaire jeremy.lemaire@univ-lr.fr

- ¹ Centre d'Études Biologiques de Chizé (CEBC), UMR 7372 CNRS-La Rochelle Université, 79360 Villiers en Bois, France
- ² Littoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS-La Rochelle Université, 2 Rue Olympe de Gouges, 17000 La Rochelle, France
- ³ Parc Zoologique de Paris, Muséum National d'Histoire Naturelle, 53 Avenue de Saint Maurice, 75012 Paris, France
- ⁴ Haus des Meeres Aqua Terra Zoo GmbH, Fritz-Grünbaum Platz 1, 1060 Vienna, Austria
- ⁵ Institut Universitaire de France (IUF), 1 Rue Descartes, 75005 Paris, France

ecosystems (Ericksen et al. 2003; Scheuhammer and Sandheinrich 2007; Hsu-Kim et al. 2018). In anoxic conditions, aquatic microorganisms can transform inorganic Hg into methylmercury (MeHg), the most bioavailable and toxic form of Hg (Jensen and Jernelöv 1969; Benoit et al. 2003). Importantly, MeHg biomagnifies through food webs and bio-accumulates in the tissues of top predators, which makes them particularly vulnerable to this contaminant (Eagles-Smith et al. 2018).

Crocodilians belong to the world's largest predators, and as such, they have important functions in ecosystems and can constitute indicators of ecosystem health (Somaweera et al. 2020). As apex predators, crocodilians bioaccumulate environmental contaminants that biomagnify across food webs, and thus can be particularly vulnerable to their toxicity (Cook et al. 1991; Camus et al. 1998; Rainwater et al. 2007). Therefore, they are relevant bioindicators regarding environmental contamination (Guillette et al. 1994; Manolis et al. 2002; Campbell et al. 2003; Chumchal et al. 2011; Schneider et al. 2015). In addition, such an evaluation is useful to determine the relatively poorly known detrimental effects of contaminants on this taxon, such as physiological and reproductive impairment and DNA damages (Guillette et al. 2000; Siroski et al. 2016; Burella et al. 2018; Lemaire et al. 2021a).

Crocodilians bioaccumulate Hg at various concentrations, depending on the species and on the location (Elsey et al. 1999; Rumbold et al. 2002; Campbell et al. 2010; Vieira et al. 2011; Nilsen et al. 2017a; Lemaire et al. 2021a). However, interspecific variability across the whole group remains still poorly understood to date as most of the available studies on Hg concentrations have disproportionately focused on two species, the American Alligator, Alligator mississippiensis and the Morelet's Crocodile, Crocodylus moreletii (Yanochko et al. 1997; Jagoe et al. 1998; Elsey et al. 1999; Burger et al. 2000; Rainwater et al. 2007; Horai et al. 2014; Trillanes et al. 2014; Nilsen et al. 2017a, 2019; Buenfil-Rojas et al. 2018, 2020). Some recent studies, however, have focused on several other species (Almli et al. 2005; Vieira et al. 2011; Lázaro et al. 2015; Schneider et al. 2015; Marrugo-Negrete et al. 2019; Lemaire et al. 2021a,b). Furthermore, Hg concentrations in crocodilians have been determined in a variety of tissues (blood, muscles, internal organs, or keratinized tissues), depending on the study, which limits robust comparisons between them. It is worth noting that the associated methodologies (e.g., sample preparation and tissues studied) are variable, which also limits straightforward comparisons between studies (Schneider et al. 2015). Furthermore, total Hg (THg) concentrations in these papers are either presented as the concentration relative to the wet- or dry-weight of sampled tissues. Clearly, such discrepancy prevents direct comparison between studies, especially as the amount of tissue moisture content, and its variation within and across tissues, on Hg concentration has not yet been thoroughly investigated. In addition, one of the most studied tissues in crocodilians is the muscle, most likely because it is used for human consumption (Delany et al. 1988; Elsey et al. 1999; Eggins et al. 2015; Rivera et al. 2016). Although assessing Hg contamination in such tissue can be useful, many species are now classified by the IUCN as in danger of extinction, which supports both the reduction of crocodile hunting per se, and the development of specific, non-lethal sampling methods to assess their Hg contamination levels and its effects (as methods have shown for other reptile species, Day et al. 2005; Lemaire et al. 2018; Beau et al. 2019).

In this context, the goals of the present study were twofold: first, we assessed the moisture content of different tissues (claws, scutes and total blood) and investigated the influence of sample preparation on THg quantification to provide recommendations to harmonize Hg assessment across multiple matrices. Second, we compared THg concentrations between several tissues (full scutes, scute keratin layers, claws, total blood, and red blood cells) to evaluate the use of non-lethal sampling methods to facilitate future comparisons of contamination in crocodilians.

Materials and Methods

Sample Collection

Our study was conducted in French Guiana between April 2016 and February 2020 (Fig. 1). We sampled 51 individuals of smooth-fronted caiman, *Paleosuchus trigonatus* (Schneider, 1801) from 6 different sites, 48 individuals of spectacled caiman, *Caiman crocodilus* (Linnaeus, 1758) from 2 different sites, and 13 individuals of dwarf caiman, *Paleosuchus palpebrosus* (Cuvier, 1807) from 4 different sites. *P. trigonatus* were captured in small forest streams. *C. crocodilus* and *P. palpebrosus* were captured in marsh or stream habitats.

The snout-vent length (SVL) and total length (TL) of each individual were measured. We took claw (randomly at the posterior legs) and scute samples (in all samplings < 1 cm, and never exceeding half of the scute) of all caimans using pliers for clipping the sample. Clipping tail scutes is a common marking method in crocodilians. The scutes which were clipped in order to mark the captured individuals were therefore not the same for all animals as they followed the consecutive identification code. Claw and scute samples were placed in dry plastic containers. Blood was collected on a subsample of individuals depending on the field possibilities (N = 24 for P. trigonatus, N = 40 for C. crocodilus, and N = 7 for P. palpebrosus). Blood samples (0.2-3 ml) were drawn from the lateral tail vein using 27 gauges (25 mm) or 21 gauges (50 mm) (depending on the size of the individual animal) heparinized needles (heparin sodium). Each blood sample was separated into two tubes and kept at cold temperatures $(4 \degree C)$ until being processed at the laboratory (always < 3 h after collection). The first tube, containing total blood, was frozen at -28 °C in the lab. The second was centrifuged at 6,500 rpm for 5 min to separate red blood cells (RBCs) and plasma, after which both fractions were then frozen at -28 °C in the lab until analysis.

After sampling, each individual was released at its capture location. *C. crocodilus, P. trigonatus*, and *P. palpebrosus* are protected by the French law (Ministerial decree NOR: TREL1933710A of 08/10/2018) and a permission to capture individuals, draw blood, and sample claws and scutes was granted by the French authorities (Direction Régionale des Territoires et de la Mer) after evaluation by the CSRPN, the regional scientific committee (Permit: R03-2016-06-21-010; R03-2019-01-09-001; R03-2019-10-24-007, www.guyane. developpement-durable.gouv.fr).



Fig. 1 Location of French Guiana, distribution ranges and capture locations of the spectacled caiman, *Caiman crocodilus* (sites 2, 8), the smooth-fronted caiman, *Paleosuchus trigonatus* (sites 3, 4, 7, 9,

Sample Preparation

Claws and scutes were cleaned for 5 min in an ultrasonic bath of ultrapure water to remove all external dirts and rinsed 3 times, as described in Lemaire et al. (2021b). To assess the wet weight of claws and scutes, we weighed cleaned samples after elimination of water surplus with absorbent paper. Because cleaning procedures were relatively short and because caimans are semiaquatic species that spend most of their lifetime immerged in water, our cleaning procedures were not likely to influence wet weight assessment. In order to assess dry weight, each sample was dried in an oven for 48 h at 45 °C, which was found to be sufficient to reach stable weight (no difference was found when dried at 45 °C for 48 h, 72 h, and 96 h, data not shown). Moisture content was calculated as wet weight minus dry weight and expressed as a proportion of wet weight for further analyses. A subsample of dried scutes (N=26 for C. crocodilus, N=4 for *P. palpebrosus*, and N=4 for *P. trigonatus*) was used to perform comparison of Hg concentrations between these matrices. We separated the external layer, which is composed of keratin (corneoscute), from the underlying connective tissue, which is the link between bone/cartilage (osteoscute) and the keratin layer. For the remaining scute samples, only the keratin layer was analysed. To assess the

10, 11), and the dwarf caiman, *Paleosuchus palpebrosus* (sites 1, 2, 5, 6). (Data for distribution ranges provided by The IUCN Red List of Threatened Species)

moisture of total blood and RBC samples, we weighted the samples before and after the freeze-drying process (48 h). Freeze-dried samples were then ground into a homogeneous powder. The moisture content of total blood and RBC samples was calculated as wet weight minus dry weight and then expressed as a proportion of wet weight.

Instrumental Method and Quality Control

For all samples, total Hg (THg) was determined using an atomic absorption spectrometer AMA-254 (Advanced Mercury Analyser-254; Altec®). At least two replicates of 0.3-1.0 mg dry weight (dw) were analysed for each sample. The reproducibility for duplicate samples was approved when relative standard deviation (RSD) was below 10%. The analyses of certified reference material (CRM) TORT-2 (Lobster hepatopancreas from the National Research Council of Canada; certified Hg concentration: $0.27 \pm 0.06 \,\mu g \, g^{-1} \, dw$) and TORT-3 (Lobster hepatopancreas from the National Research Council of Canada; certified Hg concentration: $0.292 \pm 0.022 \ \mu g \ g^{-1} \ dw$) was performed at the beginning and at the end of the analytical cycle and every 10 samples, for the validation of the method. The TORT-2 measured value was $0.243 \pm 0.015 \ \mu g \ g^{-1} \ dw \ (n = 20), \ giv$ ing a recovery of $90.2 \pm 5.7\%$. The TORT-3 measured value was $0.289 \pm 0.007 \ \mu g \ g^{-1} \ dw \ (n=30)$, giving a recovery of 99.0 ± 2.3%. Blanks were included at the beginning of analytical runs and the limit of quantification of the AMA was 0.05 ng. THg concentrations are expressed in $\mu g \ g^{-1} \ dw$.

Statistical Analysis

All statistical analyses were performed using the software R, v.3.6.1 (R development Core Team 2013). All data were checked for normality and homogeneity of variances. Depending on the results, parametric or nonparametric tests were used. The differences of moisture content in each tissue between species, as well as the differences of moisture content between different tissues for each species, and the differences of moisture content for each species and each tissue between capture sites were all investigated by Kruskal–Wallis test.

Paired *t*-tests were used to compare THg concentrations between full scutes (a combination of keratin, bone and connective tissue) and scute keratin layers for each species. To access the relationship between THg concentrations in full scutes and scute keratin layers, we built a predictive equation for *C. crocodilus*. The predictive equation was built for a significant relationship between both tissues, using slope and intercept derived from the parametric linear regression line. To validate the models, bootstrapping with 1,000 iterations were applied (Harrell 2015). We did not build predictive equations for *P. palpebrosus* and *P. trigonatus* due to the restricted number of samples.

The difference of THg concentrations between tissues was assessed by Friedman ANOVA and Pairwise Bergmann–Hommel comparisons for each species. We performed Spearman rank tests to assess the relationships between body size (SVL and TL) and THg concentrations in the tissues, independently for each species. The significance level for statistical analyses was always set at p < 0.05.

Results

Moisture Content

The results of the moisture content of blood, claws, and scutes for each species are summarized in Table 1. We found significant differences in moisture content between the species for claws (Kruskal–Wallis: $\chi^2 = 22.79$, p < 0.001, n = 57) and full scutes (Kruskal–Wallis: $\chi^2 = 10.66$, p < 0.005, n = 58), but not for total blood (Kruskal–Wallis: $\chi^2 = 1.11$, p = 0.57, n = 31) (Table 1; Fig. 2). We did not find any geographic variation in moisture content of the different tissues for the three species (all p > 0.18).

THg Concentrations in Scutes

Our results show significantly higher concentrations of THg in scute keratin layers than full scutes in two species (Paired *t*-test: *C. crocodilus*, t = -5.44, p < 0.001; *P. trigonatus*, t = -4.71, p = 0.02; *P. palpebrosus*, t = -2.18, p = 0.12; Fig. 3).

Our results do not show significant differences of THg concentration between the two capture sites (site 2, n = 11; site 8, n = 15; Fig. 1) for full scutes and scute keratin layers of C. crocodilus (t-test: t = 0.24, p = 0.815and t = 0.15, p = 0.882, respectively). Because no significant differences were found between sites, all C. crocodilus were pooled. A positive significant relationship was found between THg concentrations in full scutes and the scute keratin layers for C. crocodilus (linear regression: $F_{1-24} = 93.67$, $R^2 = 0.78$, p < 0.001; Fig. 4). The significant level of the predictors and the upper and lower bounds of the 95% confidence interval (CI) are given by the equation ($R^2 = 0.788$; p < 0.001; 95% CI 0.518-0.938): THg scute keratin layers = $0.525 \times \text{THg}$ full scute + 0.321. THg in keratin layers and full scutes are expressed in $\mu g g^{-1} dw.$

Table 1 Proportion of moi	sture (%, mean ± SD (coefficient of variation), [n	nin-max] in claws, scutes and total blood of	of smooth-fronted caiman,			
Paleosuchus trigonatus, spectacled caiman, Caiman crocodilus and dwarf caiman, Paleosuchus palpebrosus from French Guiana						
Species	Moisture content in claws (<i>n</i>)	Moisture content in scutes (<i>n</i>)	Moisture content			

Moisture content in claws (<i>n</i>)	Moisture content in scutes (n)	Moisture content in total blood (<i>n</i>)	
25.7 ± 4.5 (17.5)	39.9±19.3 (48.3)	$72.1 \pm 9.5 (13.2)$	
[15.3–47.1] (29)	[13.5–81.3] (28)	[57.3–88.7] (20)	
20.7±4.1 (19.8)	26.4±4.4 (16.7)	$78.9 \pm 0.2 (0.3)$	
[15.5–26.0] (5)	[20.1–31.5] (5)	[78.8–79.0] (2)	
46.1±15.2 (33.0)	55.3±21.6 (47.2)	72.9 ± 19.2 (26.3)	
[14.8–65.0] (23)	[14.4–90.1] (25)	[39.2–90.3] (9)	
	Moisture content in claws (n) $25.7 \pm 4.5 (17.5)$ [15.3-47.1] (29) $20.7 \pm 4.1 (19.8)$ [15.5-26.0] (5) $46.1 \pm 15.2 (33.0)$ [14.8-65.0] (23)	Moisture content in claws (n) Moisture content in scutes (n) $25.7 \pm 4.5 (17.5)$ $39.9 \pm 19.3 (48.3)$ $[15.3-47.1] (29)$ $[13.5-81.3] (28)$ $20.7 \pm 4.1 (19.8)$ $26.4 \pm 4.4 (16.7)$ $[15.5-26.0] (5)$ $[20.1-31.5] (5)$ $46.1 \pm 15.2 (33.0)$ $55.3 \pm 21.6 (47.2)$ $[14.8-65.0] (23)$ $[14.4-90.1] (25)$	

n number of samples

Deringer

Fig. 2 Moisture content measured in total blood, claws and scutes (%) of spectacled caiman, Caiman crocodilus, dwarf caiman, Paleosuchus palpebrosus, and smooth-fronted caiman, Paleosuchus trigonatus, in French Guiana. Top and bottom of the boxes represent the first and last quartiles. Line across the box represents the median. Whiskers represent the 5th and 95th percentiles. Circles represent outliers



Fig. 3 THg concentrations measured in full scutes and scute keratin layers (µg g^{-1} dw) of spectacled caiman, Caiman crocodilus (a: n=26, paired t-test: t=-5.44, p<0.001), smooth-fronted caiman, Paleosuchus trigonatus (b: n=4, paired t-test: t=-4.71, p=0.02)

and dwarf caiman, Paleosuchus palpebrosus (c: n=4, paired t-test: t=-2.18, p=0.12), in French Guiana. Each pair of connected dots correspond to one individual

Relationship Between Tissues

Our results show a significant difference of THg concentrations between tissues within each of the three species (Friedman ANOVA: P. trigonatus, $\chi^2 = 20$, p < 0.001; C. *crocodilus*, $\chi^2 = 91.33$, p < 0.001; *P. palpebrosus*, $\chi^2 = 19.04$, p < 0.001, respectively; Table 2). The THg concentrations were always higher in claws and scutes than in total blood and RBCs (Fig. 5). The relationships between body size and THg concentrations in the different tissues, and the relationships between tissues were assessed by Spearman rank test for the three species and summarized in Table 3.



Fig. 4 Linear regression between the mercury (THg) concentration in full scutes and scute keratin layers (in $\mu g g^{-1}$ dw) of the spectacled caiman (*Caiman crocodilus*; n=26; $F_{1-24}=93.67$, $R^2=0.78$, p<0.001, in blue). Regression lines (in red) with 95% confidence intervals indicate highly significant relationships between two tissues. Black line represents isometric scaling (ratio 1:1 between the two tissues)

Discussion

Moisture Content

Our results show a strong inter- and intraspecific variability of moisture content between tissues (Table 1; Fig. 2). Indeed, claws contain both alpha- and beta-keratin and present the lowest moisture content, whereas scutes are mainly composed of beta-keratin in association with osteoscutes and connective tissues, which increases water content (Richardson et al. 2002). In contrast to the study of Yanochko et al. (1997), in which study sites were separated by a very large distance, we did not find any geographic differences regarding moisture content in the examined tissues, which may be linked to the more restricted area of our study. Our results suggest that the relatively large variation of moisture content in fresh samples can induce variations in terms of trace element quantification, including THg concentration. This seems particularly the case for keratinized tissues, whereas moisture content in blood samples was less variable. Overall, our assessment of the moisture content in various tissues clearly suggests that the use of dry samples for trace elements quantification should be favoured to facilitate future large-scale comparisons of contamination in crocodilians. Alternatively, the moisture content of the tissues should be specified when trace element concentrations are expressed in wet weight to allow straightforward conversion between units.

THg Concentration: Scute Keratin Layers Versus Full Scutes

Clipping tail scutes is a common sampling method in crocodilians which further allows identification of the animal and can be used for DNA, stable isotopes and contaminant analyses (Jagoe et al. 1998; De Thoisy et al. 2006; Rainwater et al. 2007; Machkour-M'Rabet et al. 2009; Trillanes et al. 2014; Pacheco-Sierra et al. 2016; Santos et al. 2018; Lemaire et al. 2021a). The skin of crocodilians is composed of bony scutes covered by connective tissue and keratin layers (Richardson et al. 2002; Alibardi 2003). Although several studies reported concentrations of Hg in crocodilian scutes (Jagoe et al. 1998; Lázaro et al. 2015; Schneider et al. 2015; Buenfil-Rojas et al. 2018), detailed information on the actual part of the scute which had been used for analyses is generally missing, precluding further comparisons (Schneider et al. 2015). With the exception of two animals, our results show that THg concentrations significantly differ between full scutes (combination of keratin, bone, and connective tissues) and scute keratin layer, which can be explained by the poor affinity of Hg for bone tissue (Schneider et al. 2015),

Table 2 Biometric data (cm) (mean \pm SD, min—max (n)) and THg concentrations (µg g⁻¹ dw; mean \pm SD, min—max (n)) in the tissues of smooth-fronted caiman, *Paleosuchus trigonatus*, spectacled

caiman, *Caiman crocodilus* and dwarf caiman, *Paleosuchus palpebrosus*, in French Guiana. *SVL* Snout Vent Length; *TL* Total Length; *RBCs* red blood cells; *n* number of samples

Species	SVL	TL	THg claws	THg scutes	THg RBCs	THg Total blood
Paleosuchus trigo-	33.32 ± 19.81	$62.11 \pm 36.42,$	$2.420 \pm 1.905^{a},$	$3.332 \pm 3.066^{a},$	0.447 ± 0.270 ^b ,	0.300±0.178 ^b ,
natus	10.90-81 (51)	22.8-143 (51)	0.147-7.509 (50)	0.087-9.859 (48)	0.049–0.774 (11)	0.032–0.738 (24)
Caiman crocodilus	32.70 ± 13.57	66.60±24.11	2.692 ± 1.608^{a}	2.638 ± 1.497^{a}	0.963 ± 0.612 ^b	0.605±0.380 ^b
	14.5–103 (48)	31.0–176.0 (48)	0.321-8.807 (48)	0.307–7.407 (47)	0.145–2.244 (26)	0.089–1.532 (40)
Paleosuchus palpe-	38.12±15.29	79.42±33.22	$\begin{array}{l} 8.351 \pm 4.965^{a} \\ 2.028 - 20.042 \ (13) \end{array}$	7.647±4.742 ^a	2.364 ± 1.884 ^b	1.376±0.986 ^b
brosus	16.5–62 (12)	34.2–150 (13)		0.789–15.628 (13)	0.447–5.775 (6)	0.540–3.415 (7)

THg values marked with the same letter (a or b) are not statistically different (Pairwise Bergmann–Hommel test, p < 0.05), test performed independently for the three species

🖄 Springer

Fig. 5 THg concentrations measured in claws, scutes, red blood cells (RBCs), and total blood ($\mu g g^{-1} dw$) of spectacled caiman, Caiman crocodilus, dwarf caiman. Paleosuchus palpebrosus, and smoothfronted caiman, Paleosuchus trigonatus, in French Guiana. Top and bottom of boxes represent the first and last quartiles. Line across box represents the median. Whiskers represent the 5th and 95th percentiles. Circles represent outlier



 Table 3
 Relationship between
 body size (cm) and THg concentration ($\mu g g^{-1} dw$) in tissues, and THg concentration between tissues of smoothfronted caiman, Paleosuchus trigonatus, spectacled caiman, Caiman crocodilus and dwarf caiman, Paleosuchus palpebrosus

	TL	SVL	THg scutes	THg claws	THg RBCs
Paleosuchus trigonatus					
THg scutes	0.877 (48)	0.885 (48)	_	_	-
THg claws	0.842 (50)	0.850 (50)	0.945 (48)		_
THg RBCs	0.451 (11)	0.623 (11)	0.614 (10)	0.614 (10)	_
THg total blood	0.730 (24)	0.769 (24)	0.897 (21)	0.914 (23)	0.771 (6)
Caiman crocodilus					
THg scutes	0.459 (47)	0.472 (47)	_	_	_
THg claws	0.468 (48)	0.475 (48)	0.909 (47)	_	_
THg RBCs	0.353 (26)	0.388 (26)	0.894 (25)	0.811 (26)	_
THg total blood	0.386 (40)	0.399 (40)	0.882 (39)	0.870 (40)	0.992 (25)
Paleosuchus palpebrosus					
THg scutes	0.831 (13)	0.846 (12)	_	_	_
THg claws	0.945 (13)	0.937 (12)	0.857 (13)	_	_
THg RBCs	0.771 (6)	0.600 (5)	0.829 (6)	0.657 (6)	_
THg total blood	0.750 (7)	0.600 (6)	0.643 (7)	0.607 (7)	0.900 (5)

Values refer to ρ (Spearman rank test) and significant relationships are in bold. Sample sizes are given in parenthesis. TL total length; SVL snout vent length; RBCs red blood cells

which results in the highest concentrations in the keratin layer (Fig. 3). In contrast, keratinized tissues of vertebrates, such as hairs/fur, feathers, and claws, generally display high Hg concentrations due to the affinity of Hg for the sulfhydryl groups contained in keratins (Appelquist et al. 1984; Schneider et al. 2012; Benjamin et al., 2018; Treu et al., 2018; Albert et al., 2019). Standardization of the analysed layers when scutes are used for Hg biomonitoring is therefore required to enable comparison between studies. In this respect, our results suggest that selecting the keratin layer should be favored, because this avoids including an unknown quantity of other tissue types (e.g., bone, connective tissues),

thereby improving the evaluation of the actual environmental contamination status.

By providing a predictive equation of the relationship between THg concentrations found in full scutes and scute keratin layers of C. crocodilus, our study gives an objective tool to compare studies using both tissues (both dry) for this species (Fig. 4).

Relationships Between Tissues

THg concentrations were higher in scutes and claws than in the RBCs and total blood (Table 2; Fig. 5). Blood is involved in the transport of Hg to different organs and represents the recent Hg exposure of the animal via its diet-one source of variation in Hg contamination in crocodilians (Lemaire et al. 2021a). In crocodilians, Hg concentrations in the blood are known to be related to Hg concentrations of internal tissues due to dynamic transfer in tissues involved in elimination (keratinized tissues), excretion (kidneys), detoxification (liver), and storage (muscles) (Eggins et al. 2015; Nilsen et al. 2017b). In contrast to blood, keratinized tissues are known to reflect the long-term exposure due to the nonreversible binding of Hg to the sulfhydryl residues of the keratins (Schneider et al. 2015; Lázaro et al. 2015; Marrugo-Negrete et al. 2019). A combination of non-lethal sampling of several tissues in caimans provides information on the recent (blood) and the long-term (keratinized tissues) Hg contamination of the individual, as well as the dynamics of Hg contamination over time, which had already been shown in other reptiles (Lemaire et al. 2018).

Our results show a positive correlation between the body size (SVL and TL) of the three species included in the study and the THg concentrations in claws and scutes (Table 3). In contrast, the THg concentration of RBCs was positively related to SVL in P. trigonatus solely. The THg concentration in total blood showed a positive relationship with the SVL and TL of *P. trigonatus* and *C. crocodilus*, but not with *P. palpebrosus* (Table 3). The variable relationship between blood Hg and body size is likely related to shortterm variations in the diet of the individuals. Accordingly, such relationship was found in some studies on crocodilians (Eggins et al. 2015; Buenfil-Rojas et al. 2018; Lemaire et al. 2021a) but not in others (Yanochko et al. 1997; Eggins et al. 2015; Lawson et al. 2020). This emphasizes the complementarity of assessing Hg concentrations related to both long-term integration (i.e., keratinized tissues such as claws and scutes) and short-term exposure (i.e., blood) in biomonitoring studies.

Conclusions

Our results highlight the need to standardize the evaluation of Hg concentrations in crocodilians between studies. The variation in moisture content between tissues and individual animals increases the variation in the reported Hg concentrations and hence precludes robust comparison between studies. To avoid such shortcomings, researchers should perform trace elements quantification to report their results based on dry weight analytical procedures or should provide the information on the tissue moisture contents when studies report concentrations based on wet weight. In addition, our results emphasize the need to analyze scute keratin layer rather than full scutes to provide less variable and more reliable values of long-term Hg contamination of caimans. Finally, the simultaneous use of both blood and keratinized tissues in crocodilians offer the possibility to quantify different time scales of Hg exposure by non-lethal sampling.

Acknowledgments The authors are grateful to the authorities of Conseil Scientifique Régional du Patrimoine Naturel (CSRPN) and La Direction Générale des Territoires et de la Mer de Guyane (DGTM), which authorized our fieldwork. They also thank the teams of "Réserve Naturelle des Nouragues," "Réserve Naturelle du Mont-Grand Matoury," "Réserve Naturelle Trésor," "Réserve Naturelle de la Trinité," "Réserve Naturelle de Kaw-Roura," "Pripis de Yiyi," the Conservatoire du Littoral, F. Starace, T. LePape, F. Beau, L. Beau and P. Gaucher for their help in the field, and E. Courtois, M. Dewynter, and Q. Martinez for providing samples. This work was supported by the Office de l'Eau de Guyane; the Office Française pour la Biodiversité, the Direction Générale des Territoires et de la Mer de Guyane, the Parc Zoologique de Paris, the Fondation d'entreprise Hermès and the CNRS. Thanks to C. Churlaud and M. Brault-Favrou from the "Plateforme Analyses Elémentaires" of LIENSs for their assistance during mercury analyses. The Institut Universitaire de France (IUF) is acknowledged for its support to P. Bustamante as a Senior Member. Thanks to the Nouragues Research Field Station (managed by CNRS) which benefits from "Investissement d'Avenir" grant managed by Agence Nationale de la Recherche (AnaEE France ANR-11-INBS-0001; Labex CEBA ANR-10-LABX-25-01), and its team for assistance in the field.

Authors' Contributions JL, FB, OM, RM, PB conceived and designed experiments. JL, FB performed statistical analysis. JL performed chemical analysis. JL, FB, OM, RM, PB wrote the manuscript.

Funding Not applicable.

Code Availability Not applicable.

Declarations

Conflict of interest The authors have no competing interests to declare.

Ethics Approval Permission to capture individuals, draw blood and sample claws and scutes was granted by the French authorities (Direction Régionale des Territoires et de la Mer) after evaluation by the CSRPN, the regional scientific committee (Permit: R03-2016–06-21–010; R03-2019–01-09–001; R03-2019–10-24–007, www.guyane. developpement-durable.gouv.fr)

Availability of Data and Materials Data are available from the corresponding author (jeremy.lemaire@univ-lr.fr).

References

- Albert C, Renedo M, Bustamante P, Fort J (2019) Using blood and feathers to investigate large-scale Hg contamination in Artic seabirds: a review. Environ Res 177:108588
- Alibardi L (2003) Immunocytochemistry and keratinization in the epidermis of crocodilians. Zool Stud 42:346–356
- Almli B, Mwase M, Sivertsen T, Musonda MM, Flåøyen A (2005) Hepatic and renal concentrations of 10 trace elements in crocodiles (*Crocodylus niloticus*) in the Kafue and Luangwa rivers in Zambia. Sci Total Environ 337:75–82

- Appelquist H, Asbirk S, Drabaek I (1984) Mercury monitoring: mercury stability in bird feathers. Mar Pollut Bull 15:22–24
- Beau F, Bustamante P, Michaud B, Brischoux F (2019) Environmental causes and reproductive correlates of mercury contamination in European pond turtles (*Emys orbicularis*). Environ Res 172:338–244
- Benjamin T, Brasso R, Midway S, Thompson D, Harden LA (2018) Mercury (Hg) concentration in gravid Blanding's Turtles (*Emydoidea blandingii*) in Northeastern Illinois. Bull Environ Contam Toxicol 101:295–299
- Benoit JM, Gilmour CC, Heyes A, Mason RP, Miller CL (2003) Geochemical and biological controls over methylmercury production and degradation in aquatic ecosystems. Am Chem Soc Symp Ser 835:262–297
- Buenfil-Rojas AM, Alvarez-Legorreta T, Cedeño-Vázquez JR (2018) Mercury and metallothioneins in blood fractions and tissues of captive Morelet's crocodiles in Quintana Roo, Mexico. Chemosphere 199:630–636
- Buenfil-Rojas AM, Alvarez-Legorreta T, Cedeño-Vazquez JR, Rendónvon Osten J, González-Jáuregui M (2020) Distribution of metals in tissues of captive and wild Morelet's crocodiles and the potential of metallothioneins in blood fractions as a biomarker of metal exposure. Chemosphere 244:125551
- Burella PM, Odetti LM, Simoniello MF, Poletta GL (2018) Oxidative damage and antioxidant defense in *Caiman latirostris* (Broadsnouted caiman) exposed in ovo to pesticide formulations. Ecotox Environ Saf 161:437–443
- Burger J, Gochfeld M, Rooney AA, Orlando EF, Woodward AR, Guillette LJ Jr (2000) Metals and metalloids in tissues of American Alligators in three Florida lakes. Arch Environ Contam Toxicol 38:501–508
- Campbell KR (2003) Ecotoxicology of crocodilians. Appl Herpetol 1:45–163
- Campbell JW, Waters MN, Tarter A, Jackson J (2010) Heavy metal and selenium concentrations in liver tissue from wild American alligator (*Alligator mississippiensis*) liver near Charleston, South Carolina. J Wildl Dis 46:1234–1241
- Camus AC, Mitchell MM, Williams JF, Jowett PLH (1998) Elevated lead levels in farmed American alligators *Alligator mississippi ensis* consuming nutria *Myocastor coypus* meat contaminated by lead bullets. J World Aquacult Soc 3:370–376
- Chumchal MM, Rainwater TR, Osborn SC, Roberts AP, Abel MT, Cobb GP, Smith PN, Bailey FC (2011) Mercury speciation and biomagnification in the food web of Caddo Lake, Texas and Louisiana, USA, a subtropical freshwater ecosystem. Environ Toxicol Chem 30:1153–1162
- Cook RA, Behler J, Breaitis P (1991) Elevated heavy metal concentrations in captive crocodilians-two cases. In: Proceedings annual meeting of American Association of zoo veterinarians, Greensboro, North Carolina, 1989, pp 151
- Day RD, Christopher SJ, Becker PR, Whitakers DW (2005) Monitoring mercury in the Loggerhead Sea Turtle, *Caretta caretta*. Environ Sci Technol 39:437–446
- De Thoisy B, Hrbek T, Farias IP, Vasconcelos WR, Lavergne A (2006) Genetic structure, population dynamics, and conservation of Black caiman (*Melanosuchus niger*). Biol Conserv 133:474–482
- Delany MF, Bell JU, Sundlof SF (1988) Concentrations of contaminants in muscle of the American Alligator in Florida. J Wildl Dis 24:62–66
- Eagles-Smith CA, Silbergeld EK, Basu N, Bustamante P, Diaz-Barriga F, Hopkins WA, Kidd KA, Nyland JF (2018) Modulators of mercury risk to wildlife and humans in the context of rapid global change. Ambio 47:170–197
- Eggins S, Schneider L, Krikowa F, Vogt RC, Silveira RD, Maher W (2015) Mercury concentrations in different tissues of turtle and

caiman species from Rio Purus, Amazonas, Brazil. Environ Toxicol Chem 34:2771–2781

- Elsey RM, Lance VA, Campbell L (1999) Mercury levels in Alligator Meat in South Louisiana. Bull Environ Contam Toxicol 63:598–603
- Ericksen JA, Gustin MS, Schorran DE, Johnson DW, Lindberg SE, Coleman JS (2003) Accumulation of atmospheric mercury in forest foliage. Atmos Environ 37:1613–1622
- Fitzgerald WF, Lamborg CH, Hammerschmidt CR (2007) Marine biogeochemical cycling of mercury. Chem Rev 107:641–662
- Guillette LJ Jr, Gross TS, Masson GR, Matter JM, Percival HF, Woodward AR (1994) Developmental abnormalities if the gonad and abnormal sex hormone concentration in juvenile alligator from contaminated and control lakes in Florida. Environ Health Perspect 102:680–688
- Guillette LJ Jr, Crain DA, Gunderson MP, Kools SAE, Milnes MR, Orlando EF, Rooney AA, Woodward AR (2000) Alligators and endocrine disrupting contaminants: a current perspective. Am Zool 40:438–452
- Harrell FE Jr (2015) Regression modeling strategies: with applications to linear models, logistic regression, and survival analysis, 2nd edn. Springer, New York
- Horai S, Itai T, Noguchi T, Yasuda Y, Adachi H, Hyobu Y, Riyadi A, Boggs ASP, Lowers R, Guillette LJ Jr, Tanabe S (2014) Concentrations of trace elements in American alligators (*Alligator mississippiensis*) from Florida, USA. Chemosphere 108:159–167
- Hsu-Kim H, Eckley CS, Achá D, Feng X, Gilmour CC, Jonsson S, Mitchell CP (2018) Challenges and opportunities for managing aquatic mercury pollution in altered landscapes. Ambio 47:141–169
- Jagoe CH, Arnold-Hill B, Yanochko GM, Winger PV, Brisbin IL Jr (1998) Mercury in alligators (*Alligator mississippiensis*) in the southeastern United States. Sci Total Environ 213:255–262
- Jensen S, Jernelöv A (1969) Biological methylation of mercury in aquatic organisms. Nature 223:753–754
- Lawson AJ, Moore CT, Rainwater TR, Nilsen FM, Wilkinson PM, Lowers RH, Guillette LJ Jr, McFadden KW, Jodice PGR (2020) Nonlinear patterns in mercury bioaccumulation in American alligators are a function of predicted age. Sci Total Environ 707:135103
- Lázaro WL, de Oliveira RF, dos Santos-Filho M, da Silva CJ, Ignácio ÁR, Díez S (2015) Non-lethal sampling for mercury evaluation in crocodilians. Chemosphere 138:25–32
- Lemaire J, Bustamante P, Olivier A, Lourdais O, Michaud B, Boissinot A, Galán P, Brischoux F (2018) Determinants of mercury contamination in viperine snakes, *Natrix maura*, in Western Europe. Sci Total Environ 635:20–25
- Lemaire J, Bustamante P, Marquis O, Caut S, Brischoux F (2021a) Influence of sex, size and trophic level on blood Hg concentrations in Black caiman, *Melanosuchus niger* (Spix, 1825) in French Guiana. Chemosphere 262:127819
- Lemaire J, Marquis O, Bustamante P, Mangione R, Brischoux F (2021b) I got it from my mother: Inter-nest variation of mercury concentration in neonate Smooth-fronted Caiman (*Paleosuchus trigonatus*) suggest maternal transfer and possible phenotypical effects. Environ Res 194:110494
- Machkour-M'Rabet S, Hénaut Y, Charruau P, Gevrey M, Winterton P, Legal L (2009) Between introgression events and fragmentation, islands are the last refuge for American crocodile in Caribbean Mexico. Mar Biol 156:1321–1333
- Manolis S, Webb G, Briton A (2002) Crocodilians and other reptiles: bioindicators of pollution. In the Finnis River. In: Markich SJ, Jeffree RA (eds) A natural laboratory of Moning impact-past, present and future. Australian Nuclear Science and Technology Organisation (ANSTO), Clayton, pp 65–69

- Marrugo-Negrete J, Durango-Hernández J, Calao-Ramos C, Urango-Cárdenas I, Díez S (2019) Mercury levels and genotoxic effect in caimans from tropical ecosystems impacted by gold mining. Sci Total Environ 664:899–907
- Nilsen FM, Dorsey JE, Lowers RH, Guillette LJ Jr, Long SE, Bowden JA, Schock TB (2017a) Evaluating mercury concentration and body condition in American alligators (*Alligator mississippiensis*) at Merritt Island National Wildlife Refuge (MINWR), Florida. Sci Total Environ 607–608:1056–1065
- Nilsen FM, Kassim BL, Delaney JP, Lange TR, Brunell AM, Guillette LJ Jr, Long SE, Schock TB (2017b) Trace element biodistribution in the American alligator (*Alligator mississippiensis*). Chemosphere 181:343–351
- Nilsen FM, Bowden JA, Rainwater TR, Brunell AM, Kassim BL, Wilkinson PM, Guillette LJ Jr, Long SE, Schock TB (2019) Examining toxic trace element exposure in American alligators. Environ Int 128:324–334
- Pacheco-Sierra G, Gompert Z, Dominguez-Laso J, Vazquez-Dominguez E (2016) Genetic and morphological evidence of a geographically widespread hybrid zone between two crocodile species, *Crocodylus acutus* and *Crocodylus moreletii*. Mol Ecol 25:3484–3498
- R Core Team (2013). R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria. http://www.R-project.org/.
- Rainwater TR, Wu TH, Finger AG, Cañas JE, Yu L, Reynolds KD, Coimbatore G, Barr B, Platt SG, Cobb GP, Anderson TA, McMurry ST (2007) Metals and organochlorine pesticides in caudal scutes of crocodiles from Belize and Costa Rica. Sci Total Environ 373:146–156
- Richardson KC, Weeb GJW, Manolis SC (2002) Crocodiles: inside out: a guide to the crocodilians and their functional morphology. Surrey Beatty & Sons, Australia, pp 1–172
- Rivera SJ, Pacheco LF, Achá D, Molina CI, Miranda-Chumacero G (2016) Low total mercury in *Caiman yacare* (Alligatoridae) as compared to carnivorous, and non-carnivorous fish consumed by Amazonian indigenous communities. Environ Pollut 218:233–371
- Rumbold DG, Fink LE, Laine KA, Niemczyk SL, Chandrasekhar T, Wankel SD, Kendall C (2002) Levels of mercury in alligators (*Alligator mississippiensis*) collected along a transect through the Florida Everglades. Sci Total Environ 297:239–252

- Santos X, Navarro S, Campos JC, Sanpera C, Brito JC (2018) Stable isotopes uncover trophic ecology of the West African crocodile (*Crocodylus suchus*). J Arid Environ 148:6–13
- Scheuhammer AM, Sandheinrich MB (2007) Recent advances in the toxicology of methylmercury in wildlife. Ecotoxicology 17:67–68
- Schneider L, Peleja RP, Kluczkovski A Jr, Freire GM, Marioni B, Vogt RC, Da Silveira R (2012) Mercury concentration in the spectacled caiman and black caiman (Alligatoridae) of the Amazon: implications for human health. Arch Environ Contam Toxicol 63:270–279
- Schneider L, Eggins S, Maher W, Vogt RC, Krikowa F, Kinsley L, Eggins SM, Da Silveira R (2015) An evaluation of the use of reptile dermal scutes as a non-invasive method to monitor mercury concentrations in the environment. Chemosphere 119:163–170
- Selin NE (2009) Global biogeochemical cycling of mercury: a review. Annu Rev Environ Resour 34:43–63
- Siroski PA, Poletta GL, Latorre MA, Merchant ME, Ortega HH, Mudry MD (2016) Immunotoxicity of commercial-mixed glyphosate in broad snouted caiman (*Caiman latirostris*). Chem Biol Interact 244:64–70
- Somaweera R, Nifong J, Rosenblatt A, Brien ML, Combrink X, Elsey RM, Grigg G, Magnusson WE, Mazzotti FJ, Pearcy A, Platt SG, Shirley MH, Tellez M, Van der Ploeg J, Webb G, Whitaker R, Webber BL (2020) The ecological importance of crocodylians: towards evidence-based justification for their conservation. Biol Rev 95:936–959
- Treu G, Krone O, Unnsteinsdóttir ER, Greenwood AD, Czirják GÁ (2018) Correlations between hair and tissue mercury concentrations in Icelandic arctic foxes (*Vulpes lagopus*). Sci Total Environ 619–620:1589–1598
- Trillanes CE, Pérez-Jiménez JC, Rosíles-Martínez R, González-Jáuregui M (2014) Metals in Caudal Scutes of Morelet's Crocodile (*Crocodylus moreletii*) from the Southern Gulf of Mexico. Bull Environ Contam Toxicol 93:423–428
- Vieira LM, da Nunes VS, do Amaral MCA, Oliveira AC, Hauser-Davis RA, Campos RC (2011) Mercury and methyl mercury ratios in caimans (*Caiman crocodilus yacare*) from the Pantanal areas, Brazil. J Environ Monit 13:280–287
- Yanochko G, Jagoe C, Brisbin L Jr (1997) Tissue mercury concentrations in alligators (*Alligator mississippiensis*) from the Florida Everglades and the Savannah River site, Carolina. Arch Environ Contam Toxicol 32:323–328

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH ("Springer Nature").

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users ("Users"), for smallscale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use ("Terms"). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

- 1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
- 2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
- 3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
- 4. use bots or other automated methods to access the content or redirect messages
- 5. override any security feature or exclusionary protocol; or
- 6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com