

Mercury contamination in the European green toad *Bufo viridis* in Vienna, Austria.

Jérémy Lemaire^{a,*}, Rosanna Mangione^a, Magdalena Spießberger^b, Stephan Burgstaller^b, Martina Staufer^c, Edith Gruber^b, Johann G. Zaller^b, Paco Bustamante^d, Lukas Landler^b

^a Department of Behavioral and Cognitive Biology, University of Vienna, Djerassiplatz 1, Vienna 1030, Austria

^b Institute of Zoology, BOKU University, Gregor Mendel Straße 33, Vienna 1180, Austria

^c Lindenbaurgasse 13, Vienna 1110, Austria

^d Littoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS-La Rochelle Université, 2 rue Olympe de Gouges, La Rochelle 17000, France

ARTICLE INFO

Keywords:

Trace elements
Amphibians
Urban area
Brain
Toe clip

ABSTRACT

Mercury (Hg) contamination affects all ecosystems worldwide. Its deleterious effects on wildlife and humans encompass a diversity of impacts from individual to population levels. In the present study, we quantified Hg concentration across various tissues (blood, brain, muscle, and toe) of green toads (*Bufo viridis*) and investigated the use of toe clips as a proxy of Hg concentration in internal tissues, including the brain. Our results show distinct patterns of Hg contamination across tissues, with the highest Hg concentration in the blood with $1.496 \pm 0.772 \mu\text{g}\cdot\text{g}^{-1}$ dry weight (dw), followed by muscle tissue with $0.687 \pm 0.376 \mu\text{g}\cdot\text{g}^{-1}$ dw, brain tissue with $0.542 \pm 0.319 \mu\text{g}\cdot\text{g}^{-1}$ dw, and toes with $0.229 \pm 0.143 \mu\text{g}\cdot\text{g}^{-1}$ dw. A strong relationship has been found between toe and brain Hg concentrations ($R^2 = 0.857$, $p < 0.001$). These results emphasize the potential of toe clipping as a reliable, non-lethal method for predicting brain Hg concentrations in the green toad. Further, results open the possibility of assessing the potential association between Hg contamination and the cognitive performance of amphibians.

1. Introduction

Environmental pollution is a global threat to humans and wildlife. Mercury (Hg) is one of the most concerning, prevalent contaminants found in ecosystems worldwide (Chen et al., 2018). While Hg can have natural origins, human activities such as gold mining, fossil fuel combustion, biomass burning, cement production, and agricultural practices have drastically elevated its levels in the environment, surpassing concentrations seen in preindustrial times by large margins (Driscoll et al., 2013; GMA, 2019). These anthropogenic activities have altered the global Hg cycle and have frequently overwhelmed natural cycling (Pirrone et al., 2010; Amos et al., 2013). In anoxic conditions, Hg can be methylated by microorganisms mainly via sulfate-reducing and iron-reducing bacteria into methylmercury (MeHg), the most toxic and bioavailable form of Hg (Compeau and Bartha, 1985; Benoit et al., 2003; Podar et al., 2015). MeHg bioaccumulates in tissues of living organisms during their lifetime and biomagnifies through trophic chains, resulting in elevated Hg contamination levels in top predators (Atwell et al., 1998; Lavoie et al., 2013; Eagles-Smith et al., 2018).

Toxic effects of Hg in humans and wildlife encompass neurological and neurobehavioral alteration, physiological and reproductive impairment, and endocrine disruption (Scheuhammer et al., 2007; Tan et al., 2009; Evers, 2018). While Hg toxicity is well documented in mammals, fishes, and birds, reptiles and amphibians remain understudied. In amphibians, Hg levels and impact remain, for most species, unknown, despite the fact that it has been shown that Hg contamination occurs in this group (Terhivuo et al., 1984; Hothem et al., 2010; Tornabene et al., 2023). Due to the complex life cycle of amphibians, exposure to aquatic contamination, as well as stressors in their terrestrial habitat, increases the risk of toxic effects (Salice et al., 2011). To date, our understanding of the deleterious effects of Hg on amphibians is limited to an increase in larvae mortality (Bergeron et al., 2011), an increase in larval development duration and prevalence of spinal malformation in larvae (Todd et al., 2011), behavioral disruptions (Burke et al., 2010), and reduction of tadpole body condition (Schlippe-Justicia et al., 2024). Notably, most studies have focused exclusively on aquatic habitats and associated Hg contamination. Amphibians are already one of the most endangered taxa worldwide (Stuart et al., 2004; Barnosky

* Corresponding author.

E-mail address: jeremy.lemaire@univie.ac.at (J. Lemaire).

<https://doi.org/10.1016/j.ecoenv.2025.117957>

Received 9 June 2024; Received in revised form 25 November 2024; Accepted 22 February 2025

Available online 1 March 2025

0147-6513/© 2025 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

et al., 2011; IUCN, 2024); understanding the effects of environmental pollutants on physiology, behavior, and ultimately survival is therefore required for risk assessment and to set conservation measures.

The European green toad (*Bufo viridis*) is one example of an amphibian species that used to be very common around Eastern and Central Europe and became locally highly endangered (Sachs et al., 2020; Höglund et al., 2022). Land degradation and loss of primary habitat caused the green toad to increasingly inhabit urban and anthropogenically altered sites (Valkanova et al., 2009; Landler et al., 2023a; Vargová et al., 2023). Often, such habitats provide ponds with stable water levels, allowing regular reproduction. However, they potentially increase toxic loads and lead to physiological alterations as well as lower survival rates. Quantifying Hg concentrations in green toad populations is important to plan future conservation measures and evaluate environmental risk factors. Furthermore, toads, as other amphibians, are potentially reducing pest insects and increase native biodiversity (Wanger et al., 2010; Sathe and Patil, 2014), their decrease may result in increased pest insects in agricultural areas. Also, toads may act as bioindicators due to their permeable skin and complex life cycles, switching from aquatic to terrestrial habitats (Salice et al., 2011).

Most research on Hg contamination in amphibians relies on lethal sampling, which limits the potential to understand chronic Hg effects on adults and the next generations. Recently, studies have validated toe clips as a non-lethal sampling method to quantify Hg contamination (Bergeon et al., 2010; Todd et al., 2012; Pfeeger et al., 2016), which is primordial to evaluating Hg toxicity in threatened species. However, to the best of our knowledge, the use of toe clips as a proxy of Hg concentration in the brain of amphibians has not been yet evaluated. It would be important to understand mercury effects on behavior and neurobiology without sacrificing animals.

The goals of our study were to 1) quantify for the first time Hg concentrations found in green toad tissues, and 2) evaluate the use of toe clips as a proxy for Hg concentration in other tissues, including the brain.

2. Material and methods

2.1. Sample collection

The European green toad (*Bufo viridis*) is a mid-sized amphibian well adapted to semi-arid to arid conditions. It is often referred to as a pioneer species, colonizing anthropogenic environments and human settlements up to larger cities (Mazgajska and Mazgajski, 2020;

Konowalik et al., 2020). Austria's main distribution area is in the Eastern part of the country, including outskirts and even sites close to the center of Vienna (Sistani et al., 2021; Landler et al., 2023b). To date, the largest known population in this city is located at Simmeringer Haide, a traditional farmland south-east of Vienna. Since the twentieth century, however, this green toad habitat is dominated by greenhouses and polytunnels, while open cultivated land is limited to small patches. An industrial area and power plants border the limits of the current core area of this population. Further, a highway runs over parts of it. During seasonal breeding, mass mortality of migrating toads is moderated by a roadkill mitigation project (Stauffer et al., 2023). Between May 5th, 2022, and June 4th, 2022, we collected 11 green toad carcasses (all adult individuals, 10 from the Simmeringer Haide and one from Donauefeld, Fig. 1) in Vienna, Austria, which were as soon as possible frozen at -20°C . Toads were killed by road traffic or of unknown cause. However, this study only used freshly collected carcasses to allow blood and brain tissue analysis. Individual dissections were performed in the laboratory where we extracted brain and muscle tissues, drew blood from the heart, and clipped toes. Toad sampling was performed under permit MA22–230917–2020 attributed by the authority of Vienna, Austria, and collected as part of the roadkill mitigation project n°1509826–2021–25 on behalf of the Vienna Environmental Protection Department – MA 22.

2.2. Mercury analysis

Whole blood, brain, muscle, and toes of each individual were freeze-dried for 48 h, grounded, and homogenized. Total Hg (THg) concentrations were determined by direct measurement using an atomic absorption spectrometer AMA-254 (Advanced Mercury Analyzer-254; Altec®) at La Rochelle University, France, following the instrumental method from Lemaire et al. (2021). For each sample (0.12–1.98 mg dw) at least two replicates were analyzed, and the reproducibility for replicate samples was approved when the relative standard deviation was below 10 %. Certified Reference Material (CRM) TORT-3 (lobster hepatopancreas; certified Hg concentration: $0.292 \pm 0.022 \mu\text{g}\cdot\text{g}^{-1}$ dry weight (dw), NRCC) was analyzed at the beginning and at the end of the analytical cycle to validate the method. Measured values for CRM TORT-3 were $0.297 \pm 0.005 \mu\text{g}\cdot\text{g}^{-1}$ dw ($n = 7$), with a recovery of $101.9 \pm 1.5 \%$. Blanks were included at the beginning of each analytical run and the limit of quantification of the AMA was 0.05 ng. Hg concentrations are further expressed in $\mu\text{g}\cdot\text{g}^{-1}$ dw.

In parallel to carcass collection, we collected water samples (June




Fig. 1. Geographical location of the 11 green toad carcasses collected (red dots) between May 5th 2022 and June 4th 2022. Ten adults from the Simmeringer Haide and one from Donauefeld, Vienna, Austria.

2nd, 2022) from two breeding ponds of green toads in Simmeringer Haide and sent them to the commercial lab GEOTest in Brno, Czech Republic, to measure mercury content in the water (limit of detection < 0.50 $\mu\text{g}\cdot\text{l}^{-1}$).

2.3. Statistical analysis

Statistical analysis was performed using the Software R v.4.2.2 (R Core Team, 2022).

All data were first checked for normality and homogeneity of variances. Analyses of Hg concentrations were performed on log-transformed values. The difference of Hg concentrations between tissues was assessed by ANOVA and Pairwise *t*-test. The relationships between Hg concentrations in different tissues were assessed via linear regressions. The significance level for statistical analyses was always set at $p < 0.05$.

3. Results

Mercury concentrations ranged from 0.109 to 3.549 $\mu\text{g}\cdot\text{g}^{-1}$ and differed between tissues (ANOVA, $p < 0.001$). Mean Hg concentrations were the highest in whole blood with $1.496 \pm 0.772 \mu\text{g}\cdot\text{g}^{-1}$, followed by muscle tissue with $0.687 \pm 0.376 \mu\text{g}\cdot\text{g}^{-1}$, brain tissue with $0.542 \pm 0.319 \mu\text{g}\cdot\text{g}^{-1}$, and toes with $0.229 \pm 0.143 \mu\text{g}\cdot\text{g}^{-1}$ (Table 1).

All tissues showed significant positive relationships between them (Table 2). The strongest relationships were between muscle and brain ($R^2 = 0.913$, $p < 0.001$), muscle and toes ($R^2 = 0.876$, $p < 0.001$), and brain and toes ($R^2 = 0.857$, $p < 0.001$) (Fig. 2).

Mercury concentrations in the water samples from breeding sites were below the limit of detection (< 0.50 $\mu\text{g}\cdot\text{L}^{-1}$).

4. Discussion

In the present study, we demonstrated a strong correlation between Hg concentrations quantified in toe clippings and those found in the brain and muscles of green toad. Environmental contaminants are considered to contribute to the global decline of amphibians significantly (Orton et al., 2023); however, knowledge of bioaccumulation and the specific impacts of Hg on amphibians still lags behind those observed in other vertebrates. Mercury contamination has been identified to occur in most amphibians, which have been evaluated so far, and to date, understanding of its effects on individuals and populations remains largely unknown (but see Schlippe-Justicia et al., 2024). Developing non-lethal sampling methods to quantify Hg concentrations is pivotal to understanding its chronic effects, especially in threatened species. These findings are particularly valuable as they open up new possibilities for understanding the impact of this potent neurotoxin on amphibian cognition.

In this study, we detected Hg contamination in all individuals. The highest Hg concentration in green toad was found in the blood with $1.496 \pm 0.772 \mu\text{g}\cdot\text{g}^{-1}$ dw (ranging from 0.681 to 3.549 $\mu\text{g}\cdot\text{g}^{-1}$ dw) and the lowest in toes clips with $0.229 \pm 0.143 \mu\text{g}\cdot\text{g}^{-1}$ dw (ranging from 0.109 to 0.617 $\mu\text{g}\cdot\text{g}^{-1}$ dw). Blood concentrations are in the range of those found in the American toad *Anaxyrus americanus*, with blood Hg ranging from 0.082 to 4.235 $\mu\text{g}\cdot\text{g}^{-1}$ ww, and Hg concentrations in toe clippings ranging from 0.032 to 0.602 $\mu\text{g}\cdot\text{g}^{-1}$ dw (Todd et al., 2012).

Table 1

Mercury (Hg) concentrations (in $\mu\text{g}\cdot\text{g}^{-1}$ dry weight) in different tissues of the European green toad *Bufo viridis* from Vienna, Austria.

Tissues	n	Hg Mean \pm SD	Hg Min – Max
Brain	11	0.542 ± 0.319	0.273 – 1.341
Muscle	11	0.687 ± 0.376	0.326 – 1.706
Toes	11	0.229 ± 0.143	0.109 – 0.617
Whole Blood	11	1.496 ± 0.772	0.681 – 3.549

Table 2

Relationships between tissue mercury concentrations of the European green toad *Bufo viridis* from Vienna, Austria. Significant relationships are in bold ($p < 0.001$).

	Brain	Muscle	Toe
Brain	-	0.913	0.857
Muscle	-	-	0.876
Whole Blood	0.546	0.696	0.530

Nevertheless, minimum Hg concentrations in our study were surprisingly high and particularly worrying. Blood is known to represent the recent Hg exposure, equal to the lifetime of erythrocytes (Rodnan et al., 1957; Monteiro and Furness, 2001). In predators, diet is the main route of Hg exposure (Chumchal et al., 2011; McGrew et al., 2014; Rumbold et al., 2018; Jackson et al., 2021). In contrast, muscles and the brain are storage tissues known to reflect the long-term assimilation of Hg. However, a potential Hg exposure route through the skin cannot be excluded in amphibians as it allows water and ionic exchange, and the contribution this may represent to the Hg burden is still unknown. The most likely source of mercury contamination in this study are legacy effects from earlier pollution. We did not detect concerning Hg levels in breeding waters, suggesting dietary contamination by contaminated prey. Further studies on Hg contamination throughout the entire trophic chain are necessary to understand better the risks associated with the prey and predators of the green toad in the area.

Methylmercury (MeHg) is the predominant Hg form in blood, brain, and muscles (Faccio et al., 2019; Henny et al., 2002; Oliveira Ribeiro et al., 1999). The elimination rate of MeHg is relatively slow and varies between species and tissues, ranging from 5 days to 3 years for whole-body MeHg burden (reviewed in Chételat et al., 2020). To the best of our knowledge, it has not yet been investigated in amphibians. The high Hg levels in muscles, which represent long-term exposure, are concerning, considering that toads collected in this study are assumed not to be older than 5 years. In addition, the results from the present study are particularly high as they are among the highest found in amphibians studied so far. A recent assessment of MeHg by Tornabene et al. (2023) shows lower concentrations in 14 amphibian species throughout the USA.

All tissues analyzed in the present study have been correlated (Fig. 1, Table 2). Generally, muscles, blood, brain, liver, and kidneys are known to be correlated as they are metabolically active tissues (Cizdziel et al., 2003). The fact that toe clips were positively correlated to internal tissues validates the potential to use this non-lethal sampling method to access internal values, which offers an alternative approach to Hg evaluation in living amphibians. Our application promotes an alternative to lethal sampling for a taxon severely affected by contaminants requiring long-term monitoring without consequences for wild populations and conservation goals.

Mercury concentrations in the brain of green toad were relatively high, $0.542 \pm 0.319 \mu\text{g}\cdot\text{g}^{-1}$ dw (ranging from 0.273 to 1.341 $\mu\text{g}\cdot\text{g}^{-1}$ dw, Table 1). In the Everglades, an ecosystem known to be impacted by Hg contamination, Hg concentration in the brain of pig frog, *Lithobates grylio*, was two-fold lower, with $0.357 \pm 0.08 \mu\text{g}\cdot\text{g}^{-1}$ dw (Ugarte et al., 2005; ww converted to dw assuming 83 % of water in the brain according to Churchill and Storey 1995). For comparison, concentrations we have found in green toad brains exceed Hg concentrations quantified in top predators' brain such as the American alligator, *Alligator mississippiensis*, in Georgia, USA, where values were $0.450 \pm 0.12 \mu\text{g}\cdot\text{g}^{-1}$ dw (Arnold, 2000), and $0.270 \pm 0.043 \mu\text{g}\cdot\text{g}^{-1}$ dw in Louisiana, USA (Moore et al., 2022), or in the polar bear *Ursus maritimus* where brain concentrations were $0.28 \pm 0.07 \mu\text{g}\cdot\text{g}^{-1}$ dw (Krey et al., 2012). However, these values are much lower than those reported in the brains of giant petrels from the Southern Indian Ocean, which ranged between 1.58 and 13.23 $\mu\text{g}\cdot\text{g}^{-1}$ dw (Renado et al., 2021).

Methylmercury, the most toxic form of Hg, generally accounts for




Fig. 2. Linear regressions between mercury (Hg) concentrations (in $\mu\text{g}\cdot\text{g}^{-1}$ dry weight) in **A:** Brain and Muscle ($R^2 = 0.913$, $p < 0.001$), **B:** Brain and Toe ($R^2 = 0.857$, $p < 0.001$), **C:** Muscle and Toe ($R^2 = 0.876$, $p < 0.001$) of the European green toad *Bufo viridis* in Vienna, Austria.

most but not all Hg found in the brain (Basu et al., 2009; Renedo et al., 2021; Manceau et al., 2021). Therefore, high brain Hg levels of green toad raise concern regarding potential neurotoxic effects and cognitive impairment. In amphibians, Hg has been identified to be responsible for the alteration of behavior and performance in the two-lined salamander, *Eurycea bislineata* (Burke et al., 2010). While the brain is a known target organ for Hg accumulation, studies evaluating Hg concentrations in wildlife brains are rare despite the apparent importance of such data. Basu et al. (2009) reported that even at low Hg levels in the brain (approximately $0.1\text{--}1.0 \mu\text{g}\cdot\text{g}^{-1}$ dw), polar bears showed a relationship between brain Hg concentration and NMDA (N-methyl-D-aspartic acid) receptor concentration, indicating neurochemical effects. It is important to emphasize that recent studies on brain toxicity have revealed that levels of Hg in the brain and its impact are species-dependent, and that coprecipitation of Hg with selenium (Se) into tiemannite (Hg:Se) can drastically reduce toxicity of Hg (Manceau et al., 2021). Goutte et al. (2014) have shown that low Hg concentrations ($>2 \mu\text{g}\cdot\text{g}^{-1}$) have strong effects on population dynamics of South Polar Skuas (*Catharacta macrormicki*) from Adélie Land, while no effects were observed at high Hg concentrations ($12 \mu\text{g}\cdot\text{g}^{-1}$) in Brown Skuas (*Catharacta lonnbergi*) from Kergelen Archipelago, highlighting that a direct relationship between high Hg concentrations and toxicity cannot be referred to without assessing the significance of cofactors such as Se. Future studies on Hg contamination in amphibians need to consider Se:Hg ratio to understand potential detoxification processes that can protect populations from Hg toxicity, additionally repercussions of chronic Hg concentration in the brain of amphibians require further evaluation.

Results of the present study show a positive relationship between toe and brain Hg concentrations ($R^2 = 0.857$, $p < 0.001$, Fig. 1). While toe clipping is a common sampling method in amphibian genetic studies with relatively little impact on the fitness of the individual (Ursprung et al., 2011; Fisher et al., 2013; Ginnan et al., 2014), it has further been

used as an appropriate predictor of blood and muscle Hg concentrations in amphibians (Todd et al., 2012), and the results of the present study reinforce these finding (Fig. 1, Table 2).

One of the main results of this study is the confirmation that toe clips are reliable in assessing brain Hg concentrations in green toads. While quantification of Hg concentration in amphibian brain tissue was limited to sampling in dead individuals, the relationship between toes and brain Hg concentrations allows future studies to extend the understanding of Hg neurotoxicity and potential cognitive repercussions on amphibians without lethal sampling. However, future studies should continue investigating species-specific relationships between tissues to facilitate field-based studies providing a non-lethal approach.

5. Conclusion

In the present study, high Hg concentrations have been quantified in different tissues of the green toad, which is worrying in regard of the known deleterious effects of this contaminant on wildlife. These results raise questions about potential sources of Hg at study sites and the extent of trace element contamination in residing wildlife.

Green toad brain Hg concentrations are concerning, especially given the neurotoxicity of Hg. However, the strong relationship between internal organs and toe clippings, where the latter can be collected through non-lethal sampling methods, opens new possibilities to study the potential effects of Hg contamination on the cognitive performance of amphibians, as toe and brain Hg concentrations show a particularly strong relationship.

On a final note, we recommend future studies to consider a potential protective effect of selenium when investigating the effect of Hg at an individual and population level.

CRedit authorship contribution statement

Mangione Rosanna: Writing – original draft, Methodology, Conceptualization. **Spießberger Magdalena:** Methodology. **Burgstaller Stephan:** Methodology. **Stauffer Martina:** Methodology. **Landler Lukas:** Writing – original draft, Methodology, Conceptualization. **Jérémy Lemaire:** Writing – original draft, Visualization, Methodology, Data curation, Conceptualization. **Bustamante Paco:** Writing – original draft, Resources. **Gruber Edith:** Methodology. **Zaller Johann:** Methodology.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors are grateful to the platform Analyses Élémentaires of LIENSs laboratory for access to the AMA. The CPER (Contrat de Projet Etat-Région) and FEDER (Fonds Européen de Développement Régional) are acknowledged for funding the AMA of LIENSs laboratory. PB is an honorary member of the IUUF (Institut Universitaire de France).

Data Availability

Data will be made available on request.

References

- Amos, H.M., Jacob, D.J., Streets, D.G., Sunderland, E.M., 2013. Legacy impacts of all-time anthropogenic emissions on the global mercury cycle. *Glob. Biogeochem. Cy.* 27, 410–421.
- Arnold, B.S. 2000. Distribution of mercury within different trophic levels of the Okefenokee Swamp, within tissues of top level predators, and reproductive effects of methyl mercury in the Nile tilapia (*Oreochromis niloticus*). Ph.D. Dissertation, University of Georgia, Athens, USA. Pp. 187.
- Atwell, L., Hobson, K.A., Welch, H.E., 1998. Biomagnification and bioaccumulation of mercury in an arctic marine food web: insights from stable nitrogen isotope analysis. *Can. J. Fish. Aquat. Sci.* 55 (5), 1114–1121.
- Barnosky, A.D., Matzke, N., Tomiya, S., Wogan, G.O.U., Swartz, B., Quental, T.B., Marshall, C., McGuire, J.L., Lindsey, E.L., Maguire, K.C., Mersey, B., Ferrer, E.A., 2011. Has the Earth's sixth mass extinction already arrived? *Nature* 471, 51–57.
- Basu, N., Scheuhammer, A., Sonne, C., Letcher, R., Born, E., Dietz, R., 2009. Is dietary mercury of neurotoxicological concern to wild polar bears (*Ursus maritimus*)? *Environ. Toxicol. Chem.* 28, 133–140.
- Benoit, J.M., Gilmour, C.C., Heyes, A., Mason, R.P., Miller, C.L., 2003. Geochemical and biological controls over methylmercury production and degradation in aquatic ecosystems. *Am. Chem. Soc. Symp. Ser.* 835, 262–297.
- Bergeron, C.M., Bodinof, C.M., Unrine, J.M., Hopkins, W.A., 2010. Mercury accumulation along a contamination gradient and nondestructive indices of bioaccumulation in amphibians. *Environ. Toxicol. Chem.* 29 (4), 980–988.
- Bergeron, C.M., Hopkins, W.A., Todd, B.D., Hepner, M.J., Unrine, J.M., 2011. Interactive effects of maternal and dietary mercury exposure have latent and lethal consequences for amphibian larvae. *Environ. Sci. Technol.* 45 (8), 3781–3787.
- Burke, J.N., Bergeron, C.M., Todd, B.D., Hopkins, W.A., 2010. Effects of mercury on behavior and performance of northern two-lined salamanders (*Eurycea bislineata*). *Environ. Pollut.* 158, 3546–3551.
- Chen, C.Y., Driscoll, C.T., Eagles-Smith, C.A., Eckley, C.S., Gay, D.A., Hsu-Kim, H., Keane, S.E., Kirk, J.L., Mason, R.P., Obrist, D., Selin, H., Selin, N.E., Thompson, M.R., 2018. A critical time for mercury science to inform global policy. *Environ. Sci. Technol.* 52, 9556–9561.
- Chételat, J., Ackerman, J.T., Eagles-Smith, C.A., Hebert, C.E., 2020. Methylmercury exposure in wildlife: A review of the ecological and physiological processes affecting contaminant concentrations and their interpretation. *Sci. Total Environ.* 711, 135117.
- Chumchal, M.M., Rainwater, T.R., Osborn, S.C., Roberts, A.P., Abel, M.T., Cobb, G.P., Smith, P.N., Bailey, F.C., 2011. Mercury speciation and biomagnification in the food web of Caddo Lake, Texas and Louisiana, USA, a subtropical freshwater ecosystem. *Environ. Toxicol. Chem.* 30 (5), 1153–1162.
- Churchill, T.A., Storey, K.B., 1995. Metabolic effects of dehydration on an aquatic frog, *Rana pipiens*. *J. Exp. Biol.* 198, 147–154.
- Cizdziel, J., Hinners, T., Cross, C., Pollard, J., 2003. Distribution of mercury in the tissues of five species of freshwater fish from Lake Mead, USA. *Environ. Monit.* 5, 802–807.
- Compeau, G.C., Bartha, R., 1985. Sulphate reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* 50, 498–502.
- Driscoll, C.T., Mason, R.P., Chan, H.M., Jacob, D.J., Pirrone, N., 2013. Mercury as a global pollutant: sources, pathways, and effects. 47 (10), 4967–4983. *Environ. Sci. Technol.*
- Eagles-Smith, C.A., Silbergeld, E.K., Basu, N., Bustamante, P., Diaz-Barriga, F., Hopkins, W.A., Kidd, K.A., Nyland, J.F., 2018. Modulators of mercury risk to wildlife and humans in the context of rapid global change. *Ambio* 47, 170–197.
- Evers, D., 2018. The effects of methylmercury on wildlife: A comprehensive review and approach for interpretation. In: Dominick, A., DellaSala, Michael, I., Goldstein (Eds.), *The Encyclopedia of the Anthropocene*, 5. Elsevier, Oxford, pp. 181–194.
- Faccio, S.D., Buckman, K.L., Lloyd, J.D., Curtis, A.N., Taylor, V.F., 2019. Bioaccumulation of methylmercury in wood frogs and spotted salamanders in Vermont vernal pools. *Ecotoxicology* 28, 717–731.
- Fisher, K.J., Guilfoyle, K.J., Hatch, K.A., 2013. Stress induced by toe-clipping in Cane toads (*Rhinella marina*). *Copeia* 2013 (3), 539–542.
- Ginnan, N.A., Lawrence, J.R., Tussell, M.E.T., Eggert, D.L., Hatch, K.A., 2014. Toe clipping does not affect the survival of Leopard frogs (*Rana pipiens*). *Copeia* 2014 (4), 650–653.
- GMA, 2019. Global mercury Assessment 2018. UN Environment Programme, Chemicals and Health Branch, Geneva, Switzerland.
- Goutte, A., Bustamante, P., Barbaud, C., Delord, K., Weimerskirch, H., Chastel, O., 2014. Demographic responses to mercury exposure in two closely related Antarctic top predators. *Ecology* 95 (4), 1075–1086.
- Henny, C.J., Hill, E.F., Hoffman, D.J., Spalding, M.G., Grove, R.A., 2002. Nineteenth century mercury: Hazard to wading birds and Cormorants of the Carson River, Nevada. *Ecotoxicology* 11, 213–231.
- Höglund, J., Bolender, L., Cortazar-Chinarro, M., Meurling, S., Laurila, A., Hermaniuk, A., Dufresnes, C., 2022. Low neutral and immunogenetic diversity in northern fringe populations of the green toad *Bufo viridis*: implications for conservation. *Conserv. Genet.* 23, 139–149.
- Hothem, R.L., Jennings, M.R., Crayon, J.J., 2010. Mercury contamination in three species of anuran amphibians from the Cache Creek Watershed, California, USA. *Environ. Monit. Assess.* 163, 433–448.
- IUCN 2024. The IUCN Red List of Threatened Species. Version 2023-1. <<https://www.iucnredlist.org>>
- Jackson, A.K., Eagles-Smith, C.A., Robinson, W.D., 2021. Differential reliance on aquatic prey subsidies influences mercury exposure in riparian arachnids and songbirds. *Ecol. Evol.* 11, 7003–7017.
- Konowalik, A., Najbar, A., Konowalik, K., Dylewski, L., Frydlewicz, M., Kisiel, P., Starzecka, A., Zalesna, A., Kolenda, K., 2020. Amphibians in an urban environment: a case study from a central European city (Wroclaw, Poland). *Urban Ecosyst.* 23, 235–243.
- Krey, A., Kwan, M., Chan, H.M., 2012. Mercury speciation in brain tissue of polar bears (*Ursus maritimus*) from the Canadian Arctic. *Environ. Res.* 114, 24–30.
- Landler, L., Burgstaller, S., Schweiger, S., 2023a. Land-use preferences of the European green toad (*Bufo viridis*) in the city of Vienna (Austria): the importance of open land in urban environments. *Front. Zool.* 20, 3.
- Landler, L., Burgstaller, S., Spießberger, M., Horvath, A., Zhelev, Z., Mollov, I.A., Sinsch, U., Nepita, J., Schwabel, F., Mazgajski, J., Mazgajski, T., Sistani, A., Schluckebier, R., Andrä, E., Ott, M., Gollman, G., 2023b. A Unified Approach to Analysis of Body Condition in Green Toads. *Diversity* 15 (1), 43.
- Lavoie, R.A., Jardine, T.D., Chumchal, M.M., Kidd, K.A., Campbell, L.M., 2013. Biomagnification of mercury in aquatic food webs: A worldwide meta-analysis. *Environ. Sci. Technol.* 47, 13385–13394.
- Lemaire, J., Brischoux, F., Marquis, O., Mangione, R., Bustamante, P., 2021. Variation of Total Mercury Concentrations in Different Tissues of Three Neotropical Caimans: Implications for Minimally Invasive Biomonitoring. *Arch. Environ. Contam. Toxicol.* 81, 15–24.
- Manceau, A., Gaillot, A., Clatzel, P., Cherel, Y., Bustamante, P., 2021. In Vivo formation of HgSe Nanoparticles and Hg-Tetraselenolate complex from methylmercury in seabirds – Implications for the Hg-Se antagonism. *Environ. Sci. Technol.* 55, 1515–1526.
- Mazgajski, J., Mazgajski, T.D., 2020. Two amphibian species in the urban environment: changes in the occurrence, spawning phenology and adult condition of common and green toads. *Eur. Zool. J.* 87, 170–179.
- McGrew, A.K., Ballweber, L.R., Moses, S.K., Stricker, C.A., Beckmen, K.B., Salman, M.D., O'Hara, T.M., 2014. Mercury in gray wolves (*Canis lupus*) in Alaska: Increased exposure through consumption of marine prey. *Sci. Total Environ.* 468–469, 609–613.
- Monteiro, L.R., Furness, R.W., 2001. Kinetics, dose-response, and excretion of methylmercury in free-living cory's shearwaters. *Environ. Sci. Technol.* 35, 739–746.
- Moore, L.A., Finger Jr., J.W., Haskins, D.L., Elsey, R.M., Castleberry, S.B., Glenn, T.C., Jagoe, C.H., Brisbin Jr., I.L., 2022. Tissue distribution of mercury in the bodies of wild American Alligators (*Alligator mississippiensis*) from a coastal marsh in Louisiana (USA). *Arch. Environ. Contam. Toxicol.* 83, 13–20.
- Oliveira Ribeiro, C.A., Rouleau, C., Pelletier, E., Audet, C., Tjälve, H., 1999. Distribution kinetics of dietary methylmercury in the Arctic charr (*Salvelinus alpinus*). *Environ. Sci. Technol.* 33, 902–907.
- Orton, F., Roberts-Rhodes, B., Whatley, C., Tyler, C.R., 2023. A review of non-destructive biomonitoring techniques to assess the impacts of pollution on reproductive health in frog and toads. *Ecotoxicol. Environ. Saf.* 262, 115163.
- Pfleeger, A.Z., Eagles-Smith, C.A., Kowalski, B.M., Herring, G., Willacker Jr., J.J., Jackson, A.K., Pierce, J.R., 2016. From tails to toes: developing nonlethal tissue indicators of mercury exposure in five amphibian species. *Ecotoxicology* 25, 574–583.

- Pirrone, N., Cinnirella, S., Feng, X., Finkelman, R.B., Friedli, H.R., Leaner, J., Masson, R., Mukherjee, A.B., Stracher, G.B., Streets, D.G., Telmer, K., 2010. Global mercury emissions to the atmosphere from anthropogenic and natural sources. *Atmos. Chem. Phys.* 10, 5951–5964.
- Podar, M., Gilmour, C.G., Brandt, C.C., Soren, A., Brown, S.D., Crable, B.R., Palumbo, A. V., Somenahally, A.C., Elias, D.A., 2015. Global prevalence and distribution of genes and microorganisms involved in mercury methylation. *Sci. Adv.* 1 (9), e1500675.
- R Core Team, 2022. R: A language and environment for statistical computing. R foundation for statistical computing. Vienna, Austria.
- Renedo, M., Pedrero, Z., Amouroux, D., Cherel, Y., Bustamante, P., 2021. Mercury isotopes of key tissues document mercury metabolic processes in seabirds. *Chemosphere* 263, 127777.
- Rodnan, G.P., Ebaugh, F.G., Fox, M.R.S., Chambers, D.M., 1957. The life span of the red blood cell and the red blood cell volume in the chicken, pigeon and duck as estimated by the use of $\text{Na}_2\text{Cr}^{51}\text{O}_4$: with observation on red cell turnover rate in the mammal, bird and reptile. *Blood* 12 (4), 355–366.
- Rumbold, D.G., Lange, T.R., Richard, D., DelPizzo, G., Hass, N., 2018. Mercury biomagnification through food webs along a salinity gradient down-estuary from a biological hotspot. *Estuar. Coast. Shelf Sci.* 205, 116–125.
- Sachs, M., Schluckebier, R., Poll, K., Schulz, V., Sabino-Pinto, J., Schmidt, E., Simon, K., Künzel, S., Ziegler, T., Arndt, H., Vences, M., 2020. Evidence of *Batrachochytrium dendrobatidis* and other amphibian parasites in the Green toad (*Bufo viridis*), syntopic amphibians and environment in the Cologne Bay, Germany. *Salamandra* 56, 275–284.
- Salice, C.J., Rowe, C.L., Pechmann, J.H.K., Hopkins, W.A., 2011. Multiple stressors and complex life cycles: Insights from a population-level assessment of breeding site contamination and terrestrial habitat loss in an amphibian. *Environ. Toxicol. Chem.* 30 (12), 2874–2882.
- Sathe, T.V., Patil, R.N., 2014. Diversity and utility of Toads (Amphibia: Anura) in biological insect pest control in Satara, India. *Zoology* 4 (6), 526–527.
- Scheuhammer, A.M., Meyer, M.W., Sandheinrich, M.B., Murray, M.W., 2007. Effects of environmental methylmercury on the health of wild birds, mammals, and fish. *Ambio* 36 (1), 12–18.
- Schlippe-Justicia, L., Lemaire, J., Dittrich, C., Mayer, M., Bustamante, P., Rojas, B., 2024. Poison in the nursery: Mercury contamination in the tadpole-rearing sites of an Amazonian frog. *Sci. Total Environ.* 912, 169450.
- Sistani, A., Burgstaller, S., Gollman, G., Landler, L., 2021. The European green toad, *Bufo viridis*, in Donauefeld (Vienna, Austria): status and size of the population. *Herpetozoa* 34, 259–264.
- Staufer, M., Burgstaller, S., Horvath, A., Landler, L., 2023. Temporal and spatial variation in local sex ratios in a suburban population of the European green toad *Bufo viridis*. *BMC Ecol. Evol.* 23, 6.
- Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L., Waller, R.W., 2004. Status and trends of amphibian declines and extinctions worldwide. *Science* 306, 1783–1786.
- Tan, S.W., Meiller, J.C., Mahaffey, K.R., 2009. The endocrine effects of mercury in humans and wildlife. *Crit. Rev. Toxicol.* 39 (3), 228–269.
- Terhivuo, J., Lodenius, M., Nuorteva, P., Tulisalo, E., 1984. Mercury content of common frogs (*Rana temporaria* L.) and common toads (*Bufo bufo* L.) collected in southern Finland. *Ann. Zool. Fenn.* 21, 41–44.
- Todd, B.D., Bergeron, C.M., Hepner, M.J., Hopkins, W.A., 2011. Aquatic and terrestrial stressors in amphibians: A test of the double jeopardy hypothesis based on maternally and trophically derived contaminants. *Environ. Toxicol. Chem.* 30 (10), 2277–2284.
- Todd, B.D., Bergeron, C.M., Hopkins, W.A., 2012. Use of toe clips as a nonlethal index of mercury accumulation and maternal transfer in amphibians. *Ecotoxicology* 21, 882–887.
- Tornabene, B.J., Hossack, B.R., Halstead, B.J., Eagles-Smith, C.A., Adams, M.J., Backlin, A.R., Brand, A.B., Emery, C.S., Fisher, R.N., Fleming, J., Glorioso, B.M., Gear, D.A., Campbell Grant, E.H., Kleeman, P.M., Miller, P.A.W., Muths, E., Pearl, D.A., Rowe, J.C., Rumrill, C.T., Waddle, J.H., Winzler, M.E., Smalling, K.L., 2023. Broad-Scale assessment of methylmercury in adult amphibians. *Environ. Sci. Technol.* 57, 17511–17521.
- Ugarte, C.A., Rice, K.G., Donnelly, M.A., 2005. Variation of total mercury concentrations in pig frogs (*Rana grylio*) across the Florida Everglades, USA. *Sci. Total Environ.* 345, 51–59.
- Ursprung, E., Ringler, M., Robert, J., Walter, H., 2011. Toe regeneration in the neotropical frog *Allobates femoralis*. *Herpetol. J.* 21, 83–86.
- Valkanova, M.V., Mollov, I.A., Nikolov, B.N., 2009. Mortalities of the Green Toad, *Epidalea viridis* (Laurenti, 1769) in urban Environment: A Case Study from the City of Plovdiv. *Ecol. Balk.* 1, 21–26.
- Vargová, V., Guziová, D., Balogová, M., Pipová, N., Uhrin, M., Kanuch, P., 2023. Urban environment determines population genetics in the green toad, *Bufo viridis*. *Eur. J. Wildl. Res.* 69, 86.
- Wanger, T.C., Wielgoss, A.C., Motzke, I., Clough, Y., Brook, B.W., Sodhi, N., Tscharntke, T., 2010. Endemic predators, invasive prey and native diversity. *Proc. R. Soc. B.* 278, 690–694.